This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.

+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.

+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.

+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`

LANE MEDICAL LIBRARY STANFORD
2 45 0174 8949

Lane Medical Library
Stanford University Medical Center

Gift

LANE MEDICAL LIBRARY
STANFORD UNIVERSITY
MEDICAL CENTER
STANFORD, CALIF. 94305

Geo. W. Fowler,
Union, Pa. Med. Dept.
Philada.
Pa.

MINOR SURGERY

AND

BANDAGING

INCLUDING THE

TREATMENT OF FRACTURES AND DISLOCATIONS, TRACHEOTOMY, INTUBATION OF THE LARYNX, LIGATIONS OF ARTERIES AND AMPUTATIONS.

BY

HENRY R. WHARTON, M.D.,

DEMONSTRATOR OF SURGERY AND LECTURER ON SURGICAL DISEASES OF CHILDREN IN THE UNIVERSITY OF PENNSYLVANIA, SURGEON TO THE PRESBYTERIAN HOSPITAL, THE METHODIST EPISCOPAL HOSPITAL, THE CHILDREN'S HOSPITAL, AND THE DREXEL HOSPITAL FOR CHILDREN; CONSULTING SURGEON TO THE RUSH HOSPITAL FOR DISEASES OF THE CHEST, ETC.

WITH FOUR HUNDRED AND THREE ILLUSTRATIONS.



PHILADELPHIA:
LEA BROTHERS & CO.
1891.

LANE LIBRARY. STANFORD UNIVERSITY

Entered according to the Act of Congress, in the year 1891, by
LEA BROTHERS & CO.,
in the office of the Librarian of Congress. All rights reserved.

DORNAN, PRINTER.

PREFACE.

THE author has, in this work, endeavored to present, in as concise a manner as possible, a description of the various bandages, surgical dressings, and minor surgical procedures which are employed in the practice of surgery at the present time. The preparation and application of the antiseptic dressings now most commonly used have also received full consideration. The article upon Bandaging is fully illustrated with cuts, mostly new and taken from photographs, which, it is hoped, will prove of value as furnishing an accurate representation of the most important bandages used in surgical practice; the same is in a measure true of the article upon the dressing of Fractures and Dislocations, in which many new cuts of the same kind appear.

The work also contains short articles upon Tracheotomy, Intubation of the Larynx, Ligations of Arteries, and Amputations, and, although these procedures are scarcely to be included with those of Minor Surgery, it is hoped that their description will increase the value of the work to medical students, for whose use it has

been prepared. The author's thanks are due to Dr. Walter D. Green for his kind assistance in revising the proof-sheets, and to Mr. James Wood for the skilful photographic work used in illustrating several of the articles.

112 South Eighteenth St.,
Philadelphia, August, 1891.

CONTENTS.

PART I.

BANDAGING.

	PAGE
Varieties of Bandages	13–32
Bandages of the Head and Neck	32–44
Bandages of the Upper Extremity	44–58
Bandages of the Trunk	58–64
Bandages of the Lower Extremity	65–76
Special Bandages	76–82
Fixed Dressings or Hardening Bandages	82–98

PART II.

MINOR SURGERY.

Theory of Asepsis and Antisepsis in Wound Treatment	99–101
Methods and Dressings Employed in Wounds to Secure Asepsis	101–109
Preparation of Materials and Dressings Used in Aseptic Surgery	109–120
Preparations for and Details of Aseptic Operations	120–129
Materials Used in Surgical Dressings	129–135
Procedures Employed in Minor Surgery	135–192
Anæsthetics	192–201
Trusses	201–206
Use of Catheters and Bougies	206–216
Sutures and Ligatures	217–235

PAGE

Ligatures Employed in the Treatment of Vascular Growths 235–240
Treatment of Hemorrhage 241–262
Treatment of Abscesses 263–266
Dressing of Wounds, Burns and Scalds, Bedsores, Sprains 266–275
Tracheotomy, Laryngotomy, and Laryngo-tracheotomy 275–286
Intubation of the Larynx 286–289

PART III.

FRACTURES . . . 290–302
Treatment of Special Fractures 302–361

PART IV.

DISLOCATIONS . . . 361–364
Treatment of Special Dislocations . . . 364–392

PART V.

LIGATION OF ARTERIES . . 393–395
Ligation of Special Arteries 395–427

PART VI.

AMPUTATIONS . . . 428
Special Amputations 428–480
Index 483

PART I.

BANDAGING.

BANDAGES constitute one of the most widely used and important surgical dressings; they are employed to hold dressings in contact with the surface of the body, to make pressure, to hold splints in place in the treatment of fractures and dislocations, and to restore to their natural position parts which may have become displaced.

Bandages may be prepared of various materials, such as linen, crinoline, flannel, cheese or tobacco cloth, rubber sheeting, or muslin, bleached or unbleached; the latter material is the most commonly employed, by reason of the ease with which it is obtained and its cheapness; flannel, from its elasticity, is sometimes used, but its employment for bandages is now generally limited to its use in dressings for operative work in connection with the eye, and for a primary roller in the application of the plaster-of-Paris dressings.

Bandages are either *simple*, when composed of one piece of material such as the ordinary roller bandage, or *compound* when prepared of one or more pieces, adapted by size and shape to peculiar objects.

Bandages are also described as uniting, dividing, compressing, expelling or retaining bandages, according to the purposes they serve by their application.

The importance of being perfectly familiar with the general rules of bandaging and proficient in the application of the roller bandage cannot be over-estimated, and both the student and general practitioner will never have cause to

regret the time occupied in learning to apply neatly this form of surgical dressing.

A well-applied bandage adds to the comfort of the patient, and the method of its application often secures for the physician the confidence both of the patient and of his friends, while, on the other hand, a badly applied bandage is apt to be uncomfortable and insecure, and to meet with their adverse criticism.

The Roller Bandage.

The roller bandage consists of a strip of woven material, prepared from some of the materials previously mentioned, of variable length and width according to the portion of the body to which it is to be applied; this, for ease of application, is rolled into a cylindrical form.

The material commonly employed for the roller bandage is unbleached muslin, although, for special purposes, linen,

Fig 1.

Bandage winder.

flannel, rubber sheeting, crinoline or cheese-cloth may be used. It is important that the roller bandage should consist of one piece, free from seams and selvage, for if made of a number of pieces sewed together, or if it contains creases or

selvage it cannot be so neatly applied, and it is not so comfortable to the patient, as it is apt to leave creases upon the skin.

In preparing the ordinary muslin bandage the material is torn in strips varying in length and width according to the part of the body to which it is to be applied, and it is then rolled into a cylinder, either by the hand or by a machine constructed for the purpose. (Fig. 1.)

It is important that every student and practitioner should be able to roll a bandage by hand, for in practice the medical

FIG. 2.

Rolling a bandage by hand.

attendant may at any moment be called upon to roll a bandage, in order to apply a dressing, and as the art of preparing a bandage is acquired by a little practice, it should be familiar to every student and physician. To roll a bandage by hand the strip should be folded at one extremity several times until a small cylinder is formed; this is then grasped by its extremities by the thumb and index finger of the left hand; the free extremity of the strip is then grasped between the thumb and index finger of the right hand, and by alternate pronation and supination of the right hand the cylinder is revolved and the roller is formed; the firmness of the

roller will depend upon the amount of tension which is kept upon the free extremity of the strip during the revolution of the cylinder. (Fig. 2.)

Fig. 3.

Single roller.

A bandage rolled in the form of a cylinder is called a *single* or *single-headed* roller (Fig. 3); if rolled from each

Fig. 4.

Double roller.

extremity toward the centre so that two cylinders are formed joined by the central portion of the strip, the *double* or

double-headed roller is formed. (Fig. 4.) Double rollers are not much used, and in practice the single roller will be found to be amply sufficient for the application of almost all the bandages employed in surgical dressings.

The free end of the roller bandage is called the *initial extremity;* the end which is enclosed in the centre of the cylinder is its *terminal extremity;* and the portion between the extremities *the body;* a roller has also two surfaces, *external* and *internal.*

Dimensions of Bandages.

Bandages vary in length and width according to the purposes for which they are employed, and in practice it will be found that a small variety of bandages will be amply sufficient for the application of the ordinary surgical dressings.

The following list comprises those most frequently used and will show their dimensions:

Bandage one inch wide, three yards in length, for bandages for the hand, fingers, and toes.

Bandage two inches wide, six yards in length, for head bandages and for the extremities in children.

Bandage two and a half inches wide, seven yards in length, for bandages of the extremities in adults; a roller of this size is the one most generally used.

Bandage three inches wide, nine yards in length, for bandages of the thigh, groin, and trunk.

Bandages four inches wide, ten yards in length, for bandages of the trunk.

General Rules for Bandaging.

In applying a roller bandage the operator should place the external surface of the free extremity of the roller upon the part, holding it in position with the fingers of the left hand until fixed by a few turns of the roller, the cylinder being held in the right hand by the thumb and fingers; for thus as the bandage is unwound it rolls into the operator's

hand, thereby giving him more control of it; care should also be taken that the turns are applied smoothly to the surface, and that the pressure exerted by each turn is uniform.

If a bandage be applied to a limb the surgeon should see that the part is in the position it is to occupy as regards flexion and extension when the dressing is completed, for a bandage applied when the limb is flexed will exert too much pressure when the limb is extended, and then may, by the pressure it exerts, become a matter of discomfort or even of danger to the patient, or if applied to an extended limb will become uncomfortable upon flexion.

My experience has been that, as a rule, those who have had little experience with the application of the roller bandage are apt to apply their bandages too tightly, and this

FIG. 5.



Bandage scissors.

may lead to disastrous consequences, especially in the dressing of fractures. Professor Ashhurst, in his clinical teaching, advises students to make use of a larger number of turns of a bandage in securing fracture dressings rather than to depend upon a few turns too firmly applied; advice which certainly conduces to the safety and comfort of the patient. When the bandage has been applied the terminal extremity should be secured by a pin or safety-pin applied transversely to the bandage, and if a pin be used its point should be buried in the folds of the bandage; if the bandage is a narrow one, the end may be split and the two tails resulting may be secured around the part by tying. In removing a bandage the folds should be carefully gathered up in a loose mass as the bandage is unwound, the mass being trans-

ferred rapidly from one hand to the other, thus facilitating its removal and preventing the part from becoming entangled in its loops. If it is desirable to cut the bandage to remove it, the use of scissors made for this purpose will be found most satisfactory. (Fig. 5.)

Varieties of Bandages.

The Circular Bandage.

This bandage consists of a few circular turns around a part, each turn covering accurately the preceding turn. This variety of bandage may be used to retain a dressing to a limited portion of the head, neck, or limbs, to make compression upon the veins of the arm before performing venesection. (Fig. 9, *b*.)

The Oblique Bandage.

In this form of bandage the turns are carried obliquely over the part, leaving uncovered spaces between the suc-

Fig. 6.

Oblique bandage.

cessive turns. (Fig. 6.) Its principal use is for the application of temporary dressings.

The Spiral Bandage.

In this bandage the turns are carried around the part in a spiral direction, each turn overlapping a portion of the

preceding one, usually one-third or one-half. (Fig. 7.) This bandage may be used to cover a part which does not increase too rapidly in diameter, for instance the abdomen, chest, or arm.

FIG 7.

Spiral bandage.

The Spiral Reversed Bandage.

This bandage is a spiral bandage, but differs from the ordinary spiral bandage in having its turns folded back or

FIG. 8.

Method of making reverses.

reversed as it ascends a part, the diameter of which gradually increases. By its use it is possible to cover by spiral

turns a part conical in shape, so as to make equable pressure upon all parts of the surface. The reverses are made as follows: After fixing the initial extremity of the roller, as the part increases in diameter the bandage is carried off a little obliquely to the axis of the limb for from four to six inches; the index finger or thumb of the disengaged hand is placed upon the body of the bandage to keep it securely in place upon the limb, the hand holding the roller is carried a little toward the limb to slacken the unwound portion of the bandage, and by changing the position of the hand holding the bandage from extreme supination to pronation the reverse is made. (Fig. 8.) Care should be taken not to attempt to make the reverse while the bandage is tense, for by so doing the bandage is twisted into a cord which is unsightly and uncomfortable to the patient, instead of forming a closely fitting reverse.

The reverse should be completed before the bandage is carried around the limb, and when it has been completed the bandage may be slightly tightened so as to conform to the part accurately.

The reverses should be in line to have the bandage present a good appearance, and care should be taken that the reverses should not be made over salient parts of the skeleton, for if they occupy such positions they cause creases in the skin and become uncomfortable to the patient.

To make reverses neatly and to have them in line requires skill and practice; a well applied spiral reversed bandage is a test of a competent bandager.

Spica Bandages.

When the turns of the roller cross each other in the form of the Greek letter lambda, leaving the previous turn about one-third uncovered, the bandage is known as a spica bandage. (Fig. 9, *a*.) These spica bandages are especially serviceable as a means of retaining surgical dressings upon particular portions of the surface of the body, such as to the shoulder, groin, or foot.

FIG. 9.

a Spica bandage. *b* Circular bandage.

Figure-of-eight Bandage.

This bandage receives its name from the turns being applied so as to form a figure-of-eight. This method of application is made use of in the Barton's bandage, the bandages of the knee and elbow, and many other bandages.

FIG. 10.

Recurrent bandage.

Recurrent Bandage.

This bandage derives its name from the fact that the roller after covering a certain part of the surface is reflected and brought back to the point of starting; it is then reversed

and carried toward the opposite point, and this manipulation is continued until the part is covered by these recurrent turns, which are then secured by a few circular turns. (Fig. 10.) This is the bandage usually employed in the dressing of stumps.

Compound Bandages.

These bandages are usually formed of several pieces of muslin or other material, and are employed to fulfil some special indication in the application of dressings to particular parts of the body. The most useful of the compound bandages are the T-bandages and the many-tailed bandages.

T-bandages.

The single T-bandage consists of a horizontal band to which is attached, about its middle, another having a vertical direction; the horizontal piece should be about twice the

Fig. 11.



Single T-bandage.

length of the vertical piece. (Fig. 11.) The single T-bandage may be used to retain dressings to the head, the horizontal piece being passed around the head from the occiput to the forehead, the vertical piece being passed over the head and secured to the horizontal piece; the shape and width of the two pieces being varied according to the indica-

tions. In applying dressings to the anal region, or perineum, or in securing a catheter in a perineal wound, the single

FIG. 12.

Single T-bandage for chest.

FIG. 13.



T-bandage of groin.

T-bandage will be found most useful. In applying a T-bandage for this purpose the body of the bandage is placed

over the spine, just above the pelvis, and the horizontal portion is tied around the abdomen. The free extremity is split into two tails for about two-thirds of its length, and is carried over the anal region and brought up between the thighs, the terminal strips passing one on each side of the scrotum and being secured to the horizontal strip in front. The single T-bandage may be variously modified according to the indications which are to be met; for instance, in applying a dressing to the breasts the horizontal strip passing around the chest may be made ten or twelve inches in width, the vertical strip, two inches in width, passes from the back over the shoulder and is secured to the horizontal strip in front. (Fig. 12.) The single T-bandage may be variously modified, according to the ideas of the surgeon, so as to meet the indications presented in special cases. For the groin a piece of muslin six inches wide at its base and thirty inches long is sewed to a horizontal strip of muslin one and a half yards long and two inches in width. It may be applied as in Fig. 13 to hold a dressing to this part.

Double T-bandage.

FIG. 14.



Double T-bandage.

The double T-bandage differs from the single bandage in having two vertical strips attached to the horizontal strip,

and it may be used for much the same purposes as the single T-bandage. (Fig. 14.) It may be conveniently used for retaining dressings to the chest, breasts or abdomen; when used for this purpose the horizontal portion should be from eight to twelve inches wide and long enough to pass one and a quarter times about the chest; two vertical strips, two inches wide and twenty inches long, should be attached to the horizontal strip a short distance apart near its middle. In applying this bandage to the chest, the horizontal strip is placed around the chest so that the vertical strips occupy a position on either side of the spine; the overlapping end of the horizontal portion is secured by pins or safety-pins, and the vertical strips are next carried one over either shoulder and secured to other portions of the bandage in front of the chest. (Fig. 15.)

Fig. 15.

Double T-bandage of chest.

The double T-bandage may also be used to secure dressings to the nose, in which event the strips should be quite narrow, about one inch in width, and should be applied as shown in Fig. 16.

Many-tailed Bandages or Slings.

These bandages are prepared from pieces of muslin of various lengths and breadths, which are split at each ex-

tremity into two, three, or more tails up to within a few inches of their centres, their width and length being regulated by the part of the body to which they are applied.

The *four-tailed* bandage may be found useful as a temporary dressing in cases of fracture of the jaw, or to hold

Fig. 16.

Double T-bandage of nose.

Fig. 17.

Four-tailed bandage of chin.

dressings to the chin. It may be prepared by taking a portion of a roller bandage three inches wide and one yard in length, and splitting each extremity up to within two inches of the centre; it is then applied as seen in Fig. 17.

The four-tailed bandage may also be used to retain dressings to the scalp, and can be prepared by taking a piece of muslin one yard and a quarter long and six or eight inches in width, splitting it at each extremity into two tails within three inches of the centre; it may then be applied as seen in Fig. 18.

Fig. 18.

Four-tailed bandage of head.

The four-tailed bandage may also be used in the temporary dressing of fractures of the clavicle—the body of the

bandage being placed upon the elbow of the injured side, two tails passing around the body, fixing the arm to the side, and two tails passing over the sound shoulder.

The many-tailed bandage may also be used for holding dressings in contact with the abdomen or trunk, and is the bandage which many surgeons employ to hold the dressings to a laparotomy wound, and to give support to the abdominal walls after this operation. In preparing this bandage, a strip of muslin, one and a half yards in length and eighteen to twenty inches in width, is required, and the extremities may be split so as to form an eight-tailed bandage. In applying this bandage to the abdomen the body is placed upon the patient's back and the tails are brought around the abdomen and overlap each other, and when sufficiently firmly drawn to make the desired amount of pressure they are secured by means of safety-pins.

Handkerchief Bandages.

The use of handkerchiefs or square pieces of muslin for the temporary or permanent dressing of wounds, fractures,

Fig. 19.



The square.

Fig. 20.

The oblong.

or dislocations was advocated many years ago by M. Mayor, a Swiss surgeon, who wrote an extensive work upon this subject, in which he reduced their application to a system.

He employed a handkerchief or square piece of muslin, and by various modifications in the application of these developed a number of very ingenious bandages.

The various forms which the handkerchief or *square* (Fig. 19) is made to assume are as follows: The *oblong*, made by folding the square once or twice on itself (Fig. 20). The *triangle*, made by bringing together the diagonal angles of the square (Fig. 21). The line of folding is known as

FIG. 21.



The triangle.

the base, the angle opposite the base the apex, and the other angles the extremities.

The *cravat* is prepared from the triangle by bringing the apex to its base, and folding it a number of times upon itself until the desired width is obtained. (Fig. 22.)

FIG. 22.



The cravat.

FIG. 23.

The cord.

The *cord* is formed from the cravat twisted upon itself (Fig. 23). The names of the various handkerchief ban-

dages are derived from the shape of the handkerchiefs used and the parts to which they are applied; the names serve as guides in their application. It is to be remembered that the base of the triangle or the body of the cravat is to be placed upon the portion the designation of which forms the final portion of the name of the bandage; thus, in the *fronto-occipital triangle*, the shape of the handkerchief is given, and we know that the base of the triangle is to be applied to the forehead and then pass to the occiput. In using the cravats the same rule applies; thus, in the *bis-axillary*

FIG. 24.



Bis-axillary cravat.

FIG. 25.



Cruro-pelvic triangle.

cravat, the body of the cravat is to be placed in the axilla of the affected side, the extremities crossed over the corresponding shoulder and carried over the chest, one before, the other behind, to the axilla of the opposite side, where they are secured. To apply the *bis-axillary cravat* (Fig. 24), a piece of muslin a yard and a quarter long and eighteen inches in width folded into a cravat is required; this bandage may be used to hold dressing to the axilla.

The Cruro-pelvic Triangle.

This bandage may be applied with a piece of muslin folded into a triangle a yard and a half long and two feet

deep. It is applied by placing the base of the triangle obliquely across the right groin and conducting the superior extremity around the left side, across the loins to the right groin, when it is secured. The inferior end should be carried around the upper part of the right thigh between it and the scrotum, to a point near the superior extremity, and fastened with a pin (Fig. 25); this bandage may be employed to secure dressings to the groin, hip, and upper portion of the thigh.

Barton's Handkerchief.

This dressing may be employed to make extension in cases of fracture of the leg or thigh. It is applied by taking a handkerchief folded into a narrow cravat and placing the body of it on the extremity of the os calcis below the insertion of the tendo Achillis, so that two-thirds of the cravat comes around under the outer malleolus, and the other third remains on the inside. The inside portion remaining parallel with the sole of the foot, the outside piece is carried over the instep and passed around it so as to form a knot, and also passed under the sole of the foot to be turned around the first turn and to form another knot at the metatarsal articulation, when both ends are carried off perpendicularly from the foot.

I have described a few of the many very ingenious bandages devised by Mayor to substitute the use of the roller bandage, which will give the student some idea of their design and application. It is well to bear in mind this system of dressing, for the occasion might occur in which the other means of bandaging could not be obtained, and the use of handkerchiefs might answer a useful purpose as temporary dressings. I think their principal use is for temporary dressings, and I do not think they will ever take the place of the roller bandage, which can be applied with much greater nicety and exactness, and certainly presents a much neater appearance.

REGIONAL BANDAGING.

Bandages for the Head and Neck.

BARTON'S BANDAGE.

Roller Two Inches in Width, Six Yards in Length.

APPLICATION.—The initial extremity of the roller should be placed on the head just behind the mastoid process and the bandage should then be carried under the occipital protuberance obliquely upward under and in front of the parietal eminence across the vertex of the skull, then downward over the zygomatic arch, under the chin, thence upward over the opposite zygomatic arch and over the top of

FIG. 26.

Barton's bandage.

the head, crossing the first turn, which was made as nearly as possible in the median line of the skull, carrying the turns of the roller under the parietal eminence to the point of commencement. The bandage is then passed obliquely around under the occipital protuberance and forward under the ear to the front of the chin, thence back to the point

from which the roller started. These figure-of-eight turns over the head and the circular turns from the occiput to the chin should be repeated, each turn exactly overlapping the preceding one until the bandage is exhausted. (Fig. 26.) The extremity should then be secured by a pin ; and pins should be introduced at the points where the turns cross each other to give additional fixation to the bandage. In applying the bandage care should be taken to see that the turns overlap each other exactly and that the turns passing over the vertex cross as near as possible in the median line of the skull.

Modified Barton's Bandage.

To obtain additional security in the application of the Barton's bandage a turn of the bandage passing from the occiput to the forehead may be made, this turn being interposed between the turns of the bandage as ordinarily applied.

Fig. 27.

Modified Barton's bandage.

(Fig. 27.) In applying this bandage after the first set of turns has been completed, that is after the bandage has been brought back to the occiput, the bandage is carried forward upon the head just over the ear, around the forehead and backward above the ear on the opposite side to the occiput;

this being done, the ordinary figure-of-eight and circular turns are made, and when these have been completed another occipito-frontal turn may be made as described above, and this may be repeated as often as is desired until the bandage is exhausted, when the extremity is fastened with a pin, and pins are also introduced at all points at which the turns cross.

Use.—This bandage is one of the most useful of the bandages of the head, being employed to secure fixation of the jaw in cases of fracture or dislocation, and for the application of dressings to the chin. I have also employed it in place of the head-gear in slinging patients for the application of the plaster-of-Paris bandage in cases of disease of the spine, a stout cord or a piece of bandage about three inches wide and one yard long being passed under the turns crossing over the vertex; this cord is then secured to the cross-bar of the extension apparatus; this will be found quite as comfortable to the patient as the ordinary head-gear employed and much less likely to slip out of place and interfere with the breathing of the patient.

A firmly applied Barton bandage holds the jaws so closely together that care should be taken in applying it to patients who are under the influence of an anæsthetic, for if vomiting occurs the material may not be able to escape from the mouth and suffocation might occur unless the bandage were promptly removed. This accident I once saw occur and the patient's condition was alarming until the bandage was cut, allowing the jaw to be opened and the contents of the mouth to escape.

Gibson's Bandage.

Roller Two Inches in Width, Six Yards in Length.

Application.—The initial extremity of the roller should be placed upon the vertex of the skull in a line with the anterior portion of the ear; the bandage is then carried downward in front of the ear to the chin, and passed under the chin, and is carried upward on the same line until it

reaches the point of starting. The same turns are repeated until three complete turns have been made; the bandage is then continued until it reaches a point just above the ear, when it is reversed and is carried backward around the occiput, and is continued around the head and forehead until it reaches its point of origin; these circular turns are continued until three turns have been made. When the bandage reaches the occiput, having completed the third turn,

FIG. 28.

Gibson's bandage.

it is allowed to drop down to the base of the skull, and it is then carried forward below the ear and around the chin, being brought back upon the opposite side of the head and neck to the point of origin; these turns are repeated until three complete turns have been made, and upon the completion of the third turn the bandage is reversed and carried forward over the occiput and vertex to the forehead, and its extremity is here secured with a pin. Pins should also be applied at the points where the turns of the bandage cross each other. (Fig. 28.)

Use.—This bandage may be used to fix the lower jaw in cases of fracture or dislocation of the jaw, but is more apt to change its position, and is therefore not so satisfactory as the Barton's bandage for this purpose.

Oblique Bandage of Angle of the Jaw.

Roller Two Inches in Width, Six Yards in Length.

Application.—The initial extremity of the roller is placed just in front of and above the left ear, and if the left angle of the lower jaw is to be covered in, the bandage is to be carried from left to right, making two complete turns around the cranium from the occiput to the forehead. If the right angle of the lower jaw is to be covered in, the turns should be made in the opposite direction.

Having made two turns from the occiput to the forehead, the bandage is allowed to drop down upon the neck, and is carried forward under the ear and under the chin to the angle

Fig. 29.

Oblique bandage of angle of the jaw.

of the jaw; it is now carried upward close to the edge of the orbit, and obliquely over the vertex of the skull, then down behind the right ear, continuing this oblique turn under the chin to the angle of the left jaw, where it ascends in the same direction as the previous turn. Three or four of these oblique turns are made, each turn overlapping the preceding one and passing from the edge of the orbit toward the ear

until the space is covered in; the bandage is then carried to a point just above the ear on the opposite side, is reversed, and finished with one or two circular turns from the occiput to the forehead, the extremity being secured by a pin. (Fig. 29.)

Use.—This will be found to be one of the most useful of the head bandages; it may be used with a compress in treating fractures of the angle of the lower jaw, for holding dressings to the lower part of the chin and to the vault of the cranium, and is especially useful in retaining dressings to the sides of the face and the parotid region. As before stated, it may be applied to cover either the right or left side of the face, and, by reason of the oblique turns, holds its position most securely, having little tendency to become displaced.

Recurrent Bandage of the Head.

Roller Two Inches in Width, Eight Yards in Length.

Application.—The initial extremity of the roller is placed upon the lower part of the forehead and the bandage is carried twice around the head from the forehead to the occiput to secure it. When the bandage is brought back to the median line of the forehead it is reversed and the reversed turn is held by the finger of the left hand while the roller is carried over the top of the head along the sagittal suture to a point just below the occipital protuberance; here it is reversed again and the reverse is held by an assistant while the roller is carried back to the forehead in an elliptical course, each turn covering in two-thirds of the preceding turn. These turns are repeated with successive reverses at the forehead and occiput until one side of the head is completely covered in, and when this is accomplished a circular turn is made from the forehead to the occiput to hold the reverses in place.

The opposite side of the head is next covered in by elliptical reversed turns made in the same manner, and when this has been accomplished two or three circular turns

are carried around the head from the forehead to the occiput to fix the previous turns. Pins should be applied at the forehead and occiput at the points where the reversed turns concentrate. (Fig. 30.)

FIG. 30.

Recurrent bandage of the head.

Use.—This bandage when well applied is one of the neatest of the head bandages, and it will be found useful to retain dressings to the vault of the cranium in the treatment of wounds of the scalp in this region. It will also be found of service in holding dressings to fractures of the cranium and to wounds after the operation of trephining. In restless patients it will sometimes become displaced, and it may be rendered more secure by pinning a strip of bandage to the circular turn in front of the ear and carrying it down under the chin and up to a corresponding point on the opposite side, where it is pinned to the circular turn; or one or two oblique turns passing from the circular turn over the vertex of the skull downward behind the ear, under the chin and up to the circular turn in front of the ear may be applied. The course of these turns is the same as those employed in the *oblique bandage of the angle of the jaw*, the extremity being secured by a pin.

V-BANDAGE OF THE HEAD.

Roller Two Inches in Width, Four Yards in Length.

APPLICATION.—The initial extremity of the roller is secured by two turns of the bandage around the cranium from the forehead to the occiput, and when the roller reaches the occipital protuberance it is allowed to drop slightly a little below this and is carried forward below the ear around the

FIG. 31.

V-bandage of the head.

front of the chin and lower lip, then backward to the point of starting. These turns passing from the occiput to the forehead and from the occiput to the chin are alternately made until a sufficient number have been applied, and the extremity is secured by a pin over the occiput. (Fig. 31.)

This bandage may be modified by carrying the turns from the occiput forward under the ear and around the upper lip and back to the occiput and alternating these turns with the occipito-frontal turns; if employed in this way a bandage of one and one-half inches in width should be used.

Use.—This bandage may be employed to hold dressings to the front of the chin, to the upper and lower lips in cases

of wounds, or to give support to these parts after plastic operations.

Head and Neck Bandage.

Roller Two Inches in Width, Four Yards in Length.

Application.—The initial extremity of the roller is placed upon the forehead and carried backward just above the ear to the occiput and is then brought forward around the opposite side of the head to the point of starting. Two

Fig. 32.

Head and neck bandage.

of these circular turns are made to fix the bandage, and when it is carried back to the occiput it is allowed to drop down slightly upon the neck and is then carried around the neck, the turns around the head alternating with the neck turns until a sufficient number of these have been applied, when the extremity of the bandage is secured by a pin at the point of crossing of the turns at the back of the head. (Fig. 32.)

Use.—This bandage may be found useful in securing dressings to the anterior or posterior portion of the neck or to the region of the occiput.

Care should be taken to apply it in such a manner that too much pressure is not made by the turns around the neck, which would be uncomfortable to the patient, and might seriously interfere with respiration.

Crossed Bandage of One Eye.

Roller Two Inches in Width, Four Yards in Length.

Application.—The initial extremity of the bandage is placed upon the forehead and fixed by two circular turns passing around the head from the occiput to the forehead;

Fig. 33.

Crossed bandage of one eye.

the roller is then carried back to the occiput and passed around this and brought forward below the ear, and passing over the outer portion of the cheek is carried upward to the junction of the nose with the forehead, and is then conducted over the parietal protuberance downward to the occiput; a circular fronto-occipital turn is next made, and when the bandage is brought back to the occiput it is brought forward

again to the cheek and ascends to the forehead, covering in two-thirds of the previous turn, and is again conducted back to the occiput; these turns are repeated, the oblique turns covering the eye alternating with circular turns around the head until the eye is completely enclosed (Fig. 33), and the bandage is finished by making a circular turn about the head and introducing a pin to secure its extremity. It will be found more comfortable to the patient to include the ear on the same side on which the eye is covered in the turns of the bandage.

Use.—This bandage will be found useful in retaining dressings to one eye. It will be more comfortable to the patient if a flannel roller be used to apply this bandage, as well as the bandage which includes both eyes.

Crossed Bandage of Both Eyes.

Roller Two Inches in Width, Six Yards in Length.

Application.—The initial extremity of the roller is placed upon the forehead and secured by two circular turns of the bandage, passing around the head from the forehead to the occiput; the roller is then carried downward behind the occiput and brought forward below the ear to the upper portion of the cheek; it is then carried upward to the junction of the nose with the forehead and conducted over the parietal protuberance to the occiput; a circular turn is now made around the head from the occiput to the forehead, and the roller is carried from the occiput over the parietal protuberance of the opposite side forward to the junction of the nose with the forehead, then downward over the eye and outer portion of the cheek below the ear and back to the occiput; a circular turn around the head is next made, and this is followed by a repetition of the previous turns, ascending over one eye, descending over the other eye, each turn alternating with a circular turn around the head. These turns are repeated until both eyes are covered in, and the bandage is finished by making a circular turn around the head, the ex-

tremity being secured by a pin. (Fig. 34.) In this bandage both ears may be covered in, or left uncovered.

FIG. 34.

Crossed bandage of both eyes.

Use.—This bandage may be used to apply dressings to both eyes, and both of these bandages covering the eyes are used where it is desired to make pressure; but, for the simple application of a light dressing or of a bandage for the exclusion of light, the Liebreich's bandage (Fig. 68) will be found more comfortable to the patient.

OCCIPITO-FACIAL BANDAGE.

Roller Two Inches in Width, Four Yards in Length.

The initial extremity of the roller is placed upon the vertex of the head, and the bandage is carried downward in front of the ear and under the jaw, and upward upon the opposite side in the same line to the vertex; two or three of these turns are made, one turn accurately covering in the other, and a reverse is made just above and in front of the ear, and two or three turns are made around the head from the occiput to the forehead, which completes the

bandage. (*Fig. 35.*) Pins should be inserted at the points where the turns of the bandage cross each other.

FIG. 35.

Occipito-facial bandage.

Use.—This bandage is employed to secure dressings to the vertex, temporal, occipital, or frontal regions.

OBLIQUE BANDAGE OF THE HEAD.

Roller Two Inches in Width, Four Yards in Length.

The initial extremity of the bandage is placed upon the forehead, and is secured by two circular turns passing around the head from the forehead to the occiput. From the occiput the bandage is carried obliquely over the highest part of the lateral aspect of the head, which is to be covered in, and is passed over the forehead and back to the occiput, and is then carried to the forehead by a circular turn, then conducted obliquely over the other side of the head and back to the occiput. These turns are repeated, so that each succeeding turn covers in three-fourths of the preceding turn until the sides of the head are covered in by descending turns, and the bandage is completed by a circular turn

passing around the head from the forehead to the occiput. (Fig. 36.) This bandage may be applied with descending or ascending turns.

FIG. 36.

Oblique bandage of the head.

Use.—This bandage is employed to make pressure upon or to hold dressings to the lateral aspects of the head.

Bandages of the Upper Extremity.

SPIRAL BANDAGE OF THE FINGER.

Roller One Inch in Width, One and a Half Yards in Length.

APPLICATION.—The initial extremity of the roller is secured by two or three turns around the wrist; the bandage is then carried obliquely across the back of the hand to the base of the finger to be covered in, then to its tip by oblique turns; a circular turn is then made and the finger is covered by ascending spiral or spiral reversed turns until its base is reached; the bandage is then carried obliquely across the back of the hand and finished by one or two circular

turns around the wrist; the extremity may be pinned or may be split into two tails, which are tied around the wrist. (Fig. 37.)

FIG. 37.

Spiral bandage of the finger.

Use.—This bandage is employed to retain dressings upon the finger and to secure splints in the treatment of fractures or dislocations of the phalanges.

GAUNTLET BANDAGE.

Roller One Inch in Width, Three Yards in Length.

APPLICATION.—The initial extremity of the roller is fixed at the wrist by one or two circular turns of the bandage; it is then carried down to the tip of the thumb by an oblique turn of the roller, and this is covered in by spiral or spiral reversed turns to the metacarpo-phalangeal articulations; the roller is then carried back to the wrist and a circular turn is made around it, and the bandage is now carried

down to the tip of the next finger by an oblique turn, which is covered-in in the same manner. When all the fingers have been covered in, the bandage is finished by circular turns around the hand and wrist. (Fig. 38.)

Fig. 38.

Gauntlet bandage.

Use.—This bandage may be employed to apply dressings to the fingers and hand in case of wounds or fractures. It was formerly much employed in the treatment of burns of the fingers to prevent the opposed ulcerated surfaces from adhering, but its use for this purpose has been supplanted by wrapping each finger in a separate dressing and applying a dressing over the whole with a few recurrent and spiral turns of a wide roller, the application of this dressing being much less painful to the patient, and being at the same time equally satisfactory.

Demi-gauntlet Bandage.

Roller One Inch in Width, Four Yards in Length.

Application.—The initial extremity of the bandage should be placed upon the wrist and fixed by two circular turns passing from the ulnar to the radial side; then carry the roller obliquely across the back of the hand to the base

Fig. 39.

Demi-gauntlet bandage.

of the index finger, pass the bandage around this and carry the roller back to the wrist, making a circular turn; it should then be carried obliquely across the hand to the base of the next finger, and so successively until the base of each of the fingers and of the thumb has been included; the bandage is then completed by a circular turn around the wrist. (Fig. 39.)

The demi-gauntlet bandage may be also applied in such a manner as to cover the palm of the hand and leave the back of the hand uncovered.

Use.—This bandage may be employed to retain light dressings to the dorsal or palmar surface of the hand.

Spica Bandage of the Thumb.

Roller One Inch in Width, Three Yards in Length.

Application.—The initial extremity of the roller is placed upon the wrist and fixed by two circular turns; then carry the roller obliquely over the dorsal surface of the thumb to its distal extremity; next make a circular or spiral turn around the thumb, and carry the bandage upward over the back of the thumb to the wrist, around which a circular turn should be made. The roller is next carried around the thumb and wrist, making figure-of-eight turns, each turn overlapping the previous one two-thirds as it ascends the thumb, and each figure-of-eight turn alternating with a circular turn about the wrist. These turns are repeated until the thumb is completely covered in with spica turns, and the bandage is finished by a circular turn around the wrist. (Fig. 40.)

Fig. 40.

Spica bandage of the thumb.

Use.—This bandage is employed to apply dressings to the dorsal surface of the thumb, and for the retention of splints in the dressing of fractures or dislocations of the bones of the thumb.

Spiral Reversed Bandage of the Upper Extremity.

Roller Two and a Half Inches in Width, Seven Yards in Length.

Application.—The initial extremity of the roller is placed upon the wrist, and secured by two turns around the wrist; the bandage is then carried obliquely across the back of the hand to the second joint of the fingers, where a circular turn should be made; the hand is covered in by two or three ascending spiral or spiral reversed turns. When the thumb has been reached, its base and the wrist are covered in by

Fig. 41.

Spiral reversed bandage of the upper extremity.

two figure-of-eight turns; the bandage is then carried up the forearm by spiral and spiral reversed turns until the elbow is reached; this may be covered in with spiral reversed turns, and the bandage is next carried up the arm with spiral reversed turns to the axilla. (Fig. 41.) If, on reaching the elbow, the arm is bent or is to be flexed in the subsequent dressing, the elbow should be covered in with figure-of-eight turns, and when this has been done the arm above may be covered in with spiral reversed turns. When properly applied, the reverses should be in line, and should not be made over the prominent ridge of the ulna.

Use.—This is one of the most generally employed of all

the roller bandages; it constitutes the primary roller which is applied in the dressing of fractures of the humerus, and is also the bandage employed in holding dressings to the arm and forearm, and in securing splints to these parts in the treatment of fractures and dislocations.

Figure-of-eight Bandage of the Elbow.

Roller Two Inches in Width, Four Yards in Length.

Application.—The initial extremity of the bandage is placed upon the forearm a short distance below the elbow-joint, and fixed by one or two circular turns, the arm being

Fig. 42.

Figure-of-eight bandage of the elbow.

flexed. The bandage is then carried by an oblique turn across the flexure of the elbow-joint, and passed around the arm a few inches above the elbow; a circular turn is then made, and the roller is next carried across the flexure of the elbow and passed around the forearm. These turns are repeated, the turns from the forearm ascending and those from the arm descending, each set of turns crossing in the flexure

of the elbow until it is covered in, and a final turn is passed circularly around the elbow-joint. (Fig. 42.)

Use.—This bandage is often employed as a part of the spiral reversed bandage of the upper extremity when the arm is to be flexed, and is also used to hold dressings to the region of the elbow-joint. It was formerly much used to hold the compress upon the wound resulting from venesection at the elbow.

Spica Bandage of the Shoulder (Ascending).

Roller Two and a Half Inches in Width, Seven Yards in Length.

Application.—The initial extremity of the roller is placed obliquely upon the outer surface of the arm opposite the axillary fold, and fixed by one or two circular turns. If

Fig. 43.

Spica bandage of the shoulder (ascending).

Fig. 44.

Spica bandage of the shoulder (descending).

the right shoulder is to be covered, the bandage is next carried across the front of the chest to the axilla of the opposite side, then around the back of the chest to the point of starting upon the arm; then conduct the roller around the arm of this side up over the shoulder, across the front of the chest, through the opposite axilla and back over the posterior surface of the chest to the point of starting; continue

to make these ascending turns, each turn overlapping the preceding turn about two-thirds until the shoulder is covered in (Fig. 43), when the extremity of the bandage may be secured by a pin at the point of ending, or the last turn may be carried from the shoulder around the back of the neck and brought forward over the opposite shoulder and pinned to the turns which pass around the axilla It should be remembered that the turns of the roller overlap each other exactly in the opposite axilla, and it will be found more comfortable to the patient to apply a little cotton wadding in the axilla to prevent the bandage from excoriating the skin of this part. Care should be taken to see that the turns are made in such a manner that the spica turns occupy, as nearly as possible, the median line of the shoulder. When this bandage is applied to the left shoulder, after fixing the initial extremity by circular turns around the arm, the roller should be carried over the back of the chest to the axilla of the opposite side and then brought back to the point of starting; the succeeding turns are then applied in the same manner.

Spica Bandage of the Shoulder (Descending).

Roller Two and a Half Inches in Width, Seven Yards in Length.

Application.—The initial extremity of the roller should be fixed upon the arm as near as possible to the axillary line by one or two circular turns, and if it is applied to the right shoulder the bandage should be passed under the axilla and carried obliquely over the shoulder to the base of the neck, then downward across the front of the chest to the axilla of the opposite side; from the axilla the roller is carried over the back of the chest to the base of the neck so as to cross the first turn at this point; it is then carried to the axilla and through this, then back to the neck, the turns descending toward the shoulder. These turns, taking the same course are repeated, each turn overlapping two-thirds of the pre-

vious one until the shoulder is covered in and the circular turn around the arm is reached, at which point the extremity is secured by a pin. (Fig. 44.)

Use.—The spica bandages of the shoulder are employed to hold dressings to the shoulder, to hold compresses over the acromial end of the clavicle in case of dislocation of that portion of the bone, to retain the shoulder-cap used in the treatment of fractures of the upper portion of the humerus, and to retain dressings to the axilla.

Figure-of-eight Bandage of the Neck and Axilla.

Roller Two Inches in Width, Five Yards in Length.

Application.—The initial extremity of the roller is fixed upon the side of the neck and secured by one or two loosely applied circular turns; if applied to the right axilla carry the bandage from left to right over the right shoulder to the posterior part of the axilla under which it passes, to ascend in front over the same shoulder to the back of the neck; these figure-of-eight turns around the neck and axilla, each turn overlapping two-thirds of the previous turn, are repeated until the desired space is covered, and the bandage is completed by a circular turn around the neck. (Fig. 45.)

Fig 45.

Figure-of-eight bandage of the neck and axilla.

Use.—This will be found a useful bandage to secure dressings to the base of the neck, the upper part of the shoulder, and to the axilla, as it does not restrict the motions of the arm unless drawn too tight.

Velpeau's Bandage.

Two Rollers Two and a Half Inches in Width, Seven Yards in Length.

Application.—The patient should place the fingers of the hand of the affected side on the opposite shoulder; the initial end of the roller should be placed on the body of the scapula of the sound side and secured by a turn made by

Fig. 46.

Velpeau's bandage.

carrying the bandage over the shoulder of the affected side, near its outer portion, then conducting it downward over the outer and posterior surface of the arm of the same side, behind the point of the elbow, and obliquely across the front of the chest to the axilla of the opposite side, thence to the point of starting. This turn should be repeated, to fix the initial extremity of the bandage. Having completed the second turn, carry the roller transversely around the thorax, passing over the flexed elbow of the affected side, from this point to the axilla, and through this to the back. From this point the roller is carried over the shoulder and down the

outer and posterior surface of the arm behind the elbow and obliquely across the front of the chest through the axilla to the back, and continuing, passes transversely across the back of the chest to the elbow, which it encircles, then passing to the axilla. These alternating turns are repeated until the arm and forearm are bound firmly to the side and chest. The vertical turns over the shoulder, each turn covering in two-thirds of the previous turn and ascending from the point of the shoulder toward the neck and from the posterior surface of the arm toward the elbow, are applied until the point of the elbow is reached. The transverse turns passing around the chest and arm are so applied that they ascend from the point of the elbow toward the shoulder, each turn covering in one-third of the previous one, and the last turn should pass transversely around the shoulder and chest, covering the wrist. (Fig. 46.)

The extremity of the bandage should be secured by a pin where it ends, and additional fixation will be secured by introducing a number of pins at the points where the turns of the bandage cross each other.

Use.—This bandage is employed to fix the arm in the treatment of certain fractures of the clavicle and scapula, also to secure fixation of the humerus after the reduction of dislocations of the shoulder-joint.

Desault's Bandage.

Three Rollers Two and a Half Inches in Width, Seven Yards in Length.

A wedge-shaped pad to fit in the axilla is also required.

These rollers are known as the *first*, *second* and *third* rollers.

First Roller of Desault's Bandage.

Application.—Before applying the first roller the arm of the patient on the injured side should be elevated and carried off at right angles to the body; the wedge-shaped

pad with its base in the axilla should next be applied to the side of the chest, and the initial extremity of the roller is placed upon the middle of the pad and fixed by two or three circular turns around the chest; the bandage is then carried obliquely across the front of the chest to the sound

FIG. 47.

First roller of Desault's bandage.

shoulder and passed under the axilla, brought over the shoulder and conducted around the chest to pass over the pad, and it is next carried obliquely down to the lower portion of the chest to a point opposite the lower end of the pad; it is now made to ascend the chest by spiral turns until the top of the pad is reached, where it is secured. (Fig. 47.)

Second Roller of Desault's Bandage.

APPLICATION.—The arm should be brought down against the side so as to press upon the pad previously applied, and the forearm should be flexed upon the arm and brought across the lower portion of the chest. The initial extremity of the roller is placed in the axilla of the sound side, and the bandage is carried around the chest and over the arm of

the injured side, making a circular turn around the chest to fix it; then spiral turns are made around the chest from above downward until the elbow is reached, the turns being more

Fig. 48.

Second roller of Desault's bandage.

firmly applied as they descend, and when this point is reached the end of the bandage is secured. (Fig. 48.)

Third Roller of Desault's Bandage.

Application.—The initial extremity of the roller is placed in the axilla of the sound side, and the bandage is carried obliquely over the front of the chest to the shoulder of the injured side, passed over this, and conducted down the back of the arm to the elbow, thence obliquely upward over the upper fifth of the forearm to the axilla of the sound side. From this point it is carried backward obliquely over the back of the chest to the shoulder; crossing the previous shoulder-turn it is conducted down the front of the arm to the elbow, then around this and backward obliquely over the back of the chest to the axilla of the sound side. These turns are repeated until three sets of turns have been applied, and the turns should overlie each other exactly. (Fig. 49.) The course of the turns of the third roller is

considered the most difficult to remember, and the student may be assisted in its correct application by remembering that all the turns start at the axilla, pass to the shoulder, and then to the elbow, and from the elbow always return to

FIG. 49.

Third roller of Desault's bandage.

the starting-point—the axilla. The turns of the third roller make two triangles, one on the anterior surface of the chest, the other upon the back.

After the application of the three rollers the hand and uncovered portion of the forearm should be supported in a sling suspended from the neck.

Use.—This bandage, applied completely, or some one of its various rollers, is employed in the treatment of fractures of the clavicle.

Bandages of the Trunk.

SPIRAL BANDAGE OF THE CHEST.

Roller Three Inches in Width, Nine Yards in Length.

APPLICATION.—The initial extremity of the roller is applied to the anterior portion of the waist, and fixed by one or two circular turns; the bandage is then carried upward,

the injured side, making a circular turn around the chest to fix it; then spiral turns are made around the chest from above downward until the elbow is reached, the turns being more

Fig. 48.

Second roller of Desault's bandage.

firmly applied as they descend, and when this point is reached the end of the bandage is secured. (Fig. 48.)

Third Roller of Desault's Bandage.

Application.—The initial extremity of the roller is placed in the axilla of the sound side, and the bandage is carried obliquely over the front of the chest to the shoulder of the injured side, passed over this, and conducted down the back of the arm to the elbow, thence obliquely upward over the upper fifth of the forearm to the axilla of the sound side. From this point it is carried backward obliquely over the back of the chest to the shoulder; crossing the previous shoulder-turn it is conducted down the front of the arm to the elbow, then around this and backward obliquely over the back of the chest to the axilla of the sound side. These turns are repeated until three sets of turns have been applied, and the turns should overlie each other exactly. (Fig. 49.) The course of the turns of the third roller is

considered the most difficult to remember, and the student may be assisted in its correct application by remembering that all the turns start at the axilla, pass to the shoulder, and then to the elbow, and from the elbow always return to

FIG. 49.

Third roller of Desault's bandage.

the starting-point—the axilla. The turns of the third roller make two triangles, one on the anterior surface of the chest, the other upon the back.

After the application of the three rollers the hand and uncovered portion of the forearm should be supported in a sling suspended from the neck.

Use.—This bandage, applied completely, or some one of its various rollers, is employed in the treatment of fractures of the clavicle.

Bandages of the Trunk.

SPIRAL BANDAGE OF THE CHEST.

Roller Three Inches in Width, Nine Yards in Length.

APPLICATION.—The initial extremity of the roller is applied to the anterior portion of the waist, and fixed by one or two circular turns; the bandage is then carried upward,

encircling the chest by ascending spiral turns, each turn covering in one-half of the previous turn until the axillary line is reached; the roller is next carried around the axilla to the back, and obliquely over this to the base of the neck of the opposite side, and then it may be passed down over the chest and pinned to the spiral turns at several points; a pin should also be inserted at the point where the last turn of the roller leaves the spiral turn upon the back of the chest. (Fig. 50.)

Fig. 50.

Spiral bandage of the chest.

Use.—This bandage is employed to hold dressings to the chest, and may be used as a temporary dressing in fractures of the ribs or sternum. Care should be taken that the bandage be not so tightly applied as to interfere with respiration.

Anterior Figure-of-eight Bandage of the Chest.

Roller Two and a Half Inches in Width, Seven Yards in Length.

Application.—The initial extremity of the roller should be placed in the axilla of one side, and fixed by two or three circular turns around the chest; the bandage is then

carried through the axilla and passed upward over the shoulder of the same side, and obliquely across the anterior portion of the chest to the axilla of the opposite side, then through this to the shoulder of the same side, and obliquely downward to the opposite axilla. These turns should be repeated, ascending from the shoulder toward the neck, each turn overlapping three-fourths of the preceding one, until five or six turns have been applied, the end of the

FIG. 51.

Anterior figure-of-eight bandage of the chest.

bandage being secured by a pin (Fig. 51), or it may be completed by a circular turn around the chest.

Use.—This bandage may be employed to bring the shoulders forward, and to hold dressings to the anterior portion of the chest.

POSTERIOR FIGURE-OF-EIGHT BANDAGE OF THE CHEST.

Roller Two and a Half Inches in Width, Seven Yards in Length.

APPLICATION.—The initial extremity of the roller should be placed upon the outer portion of the left shoulder, and the bandage is carried obliquely backward and downward to the axilla of the opposite side; it is then passed through this

and conducted over the shoulder of the same side and passed obliquely downward to the axilla of the opposite side and carried through this and brought up over the shoulder to fix the initial extremity of the roller. These turns are repeated five or six times, the same precautions being observed in

FIG. 52.

Posterior figure-of-eight bandage of the chest.

covering the turns and in ascending from the shoulder toward the neck (Fig. 52). In applying both of these bandages the crosses of the bandage, either anterior or posterior, should be made in the median line of the chest.

Use.—This bandage may be employed to hold dressings to the posterior portion of the chest and to draw the shoulders backward.

SUSPENSORY AND COMPRESSOR BANDAGE OF THE BREAST.

Roller Two and a Half Inches in Width, Seven Yards in Length.

APPLICATION.—The initial extremity of the roller should be placed upon the scapula of the affected side, and secured by two oblique turns carried over the opposite shoulder and conducted downward under the mamma to be covered in, and then carried to the axilla of the same side. Next carry

the roller transversely around the chest, covering in the lowest portion of the affected mamma. These turns should be repeated, the oblique turns from the axilla over the shoulder

FIG 53.

Suspensory and compressor bandage of the breast.

alternating with the transverse turns around the chest until the breast is covered in, each series of turns ascending, and covering two-thirds of the preceding turn. (Fig. 53.)

Use.—This bandage is employed to support the breast and to make compression at the same time; it may also be employed to hold dressings to the breast.

SUSPENSORY AND COMPRESSOR BANDAGE OF BOTH BREASTS.

Two Rollers Two and a Half Inches in Width, Seven Yards in Length.

APPLICATION.—The initial extremity of the bandage should be secured by oblique turns of the axilla and shoulder as in the preceding bandage; the bandage should next be carried transversely around the back to the breast, then under the breast and upward over the opposite shoulder, then obliquely downward around the chest to the other side,

being carried transversely over the lower portion of both breasts to the point of starting upon the back. Repeat these oblique turns from the shoulder to the axilla and from the axilla to the shoulder, and alternate these turns with a

FIG. 54.

Suspensory and compressor bandage of both breasts.

transverse turn around the chest and over both breasts. Both series of turns should ascend, and each turn should overlap two-thirds of the preceding turn. (Fig. 54.)

Use.—This bandage is employed to support and compress both breasts and to retain dressings to the breasts.

Bandages of the Lower Extremity.

Single Spica Bandage of the Groin (Ascending).

Roller Two and a Half Inches in Width, Seven Yards in Length.

Application.—Place the initial extremity of the bandage upon the anterior portion of the right thigh just below the groin and secure it by one or two circular turns around the thigh, or place the initial extremity of the roller obliquely upon the upper part of the thigh and carry it behind the

Fig. 55.

Ascending spica bandage of the groin.

thigh and upward around the outer side of the thigh to the abdomen, omitting the circular turns; then carry the bandage obliquely across the lower part of the abdomen to a point just below the crest of the left ilium and conduct it transversely around the back of the pelvis to a corresponding point on the opposite side; then bring it obliquely downward to the groin over to the inner portion of the thigh, carrying it around the thigh, crossing the starting-turn in the middle line of the thigh. These turns are

repeated, each turn ascending and covering in two-thirds of the previous turn, until six or eight complete turns have been made, and the bandage is secured at any point where it ends. (Fig. 55.)

Single Spica Bandage of the Groin (Descending).

Roller Two and a Half Inches in Width, Seven Yards in Length.

Application.—Place the initial extremity of the roller obliquely upon the anterior surface of the right thigh and secure it by one or two circular turns around the thigh, or

Fig. 56.

Descending spica bandage of the groin.

start the bandage with an oblique turn, as previously described; then carry the bandage obliquely across the abdomen to a point just below the crest of the ilium, and conduct it transversely around the back of the pelvis to a corresponding point on the opposite side; then bring it obliquely down over the lower portion of the abdomen, crossing the first turn, to the junction of the thigh with the scrotum, pass it under the thigh and bring it up over the lower part of the abdomen, let it follow the course of the first turn. These turns

are repeated, each turn descending and overlapping two-thirds of the previous turn until the groin is covered (Fig. 56). When either of these bandages is applied to the left groin, after the initial extremity of the roller is fixed, it is carried first to the crest of the ilium of the same side, then around the back of the pelvis to a corresponding point on the opposite side, then obliquely across the lower part of the abdomen to the outer aspect of the thigh, being conveyed under this and brought up between the thigh and the scrotum, passing obliquely over the groin to follow the course of the original turn. The turns may be made either to ascend or descend as the bandage is applied.

Double Spica Bandage of the Groins.

Roller Three Inches in Width, Nine Yards in Length.

Application.—The initial extremity of the roller is placed upon the abdomen just above the iliac crests and

Fig. 57.

Double spica bandage of the groins.

secured by one or two circular turns; the bandage is then carried from a point just below the crest of the right ilium

obliquely across the lower portion of the abdomen to the outer portion of the thigh, and is carried around this and brought up between the scrotum and the thigh, and is passed obliquely over the groin, crossing the previous turn in the median line, and is conducted to a point just below the crest of the ilium on the same side. The bandage is then continued around the pelvis to the same point on the opposite side, and from this point is made to pass obliquely over the groin to the inner side of the thigh, passing around this and coming up on its outer side, crossing the previous turn at the middle line of the groin, to be carried obliquely across the groin and lower part of the abdomen to the crest of the ilium on the opposite side. These turns are repeated, each turn covering in two-thirds of the previous turn, until both groins have been covered (Fig. 57). The turns may be so applied as to ascend or descend, forming the ascending or descending double spica bandage of the groin. When properly applied, this bandage presents three sets of crossing turns, one in each groin and one in the median line of the abdomen.

Use.—The spica bandages of the groin, either single or double, are employed to hold dressings to wounds in the inguinal region—for instance, those resulting from herniotomy, or from operation upon the glands of the groin. They are also employed to make pressure upon this region, and will often prove of use in the securing of compresses applied for the temporary retention of herniæ.

Figure-of-eight Bandage of the Knee.

Roller Two and a Half Inches in Width, Five Yards in Length.

Application.—The initial extremity of the roller is placed upon the thigh three inches above the patella and secured by two or three circular turns; then conduct the bandage over the outer condyle of the femur across the popliteal space to the inner border of the tibia and around the

anterior surface below the tubercle and head of the fibula and make one circular turn; the roller should then be carried obliquely across the popliteal space to the inner condyle of the femur, crossing the previous turn; then carry it around the front of the thigh to the outer condyle; repeat these

FIG 58.

Figure-of-eight bandage of the knee.

turns, ascending toward the knee from the leg and descending from the thigh toward the knee, and finish the bandage by a circular turn over the patella (Fig. 58).

Use.—This bandage is employed to hold dressings to the knee-joint either anteriorly or posteriorly. These figure-of-eight turns are often employed in covering the knee in applying the spiral reversed bandage of the lower extremity when it is desired that the patient be allowed to bend the knee.

FIGURE-OF-EIGHT BANDAGE OF BOTH KNEES.

Roller Two and a Half Inches in Width, Seven Yards in Length.

APPLICATION.—Place the knees of the patient together with a compress between them; then place the initial extremity of the roller upon one thigh, about three inches

each other in the median line of the foot, and that the turns are kept parallel to each other throughout their course.

Use.—This bandage will be found very useful when it is desired to make firm compression upon the foot or to retain dressings to it; it is especially useful in the treatment of sprains of the ankle or anterior tarsus.

Bandage of Foot Covering the Heel (American).

Roller Two and a Half Inches in Width, Seven Yards in Length.

Application.—The initial extremity of the roller is placed upon the leg just above the malleoli and fixed by two circular turns around the leg; the bandage is then carried obliquely across the dorsum of the foot to the metatarso-phalangeal articulation, at which point a circular turn is made; two or three spiral or spiral reversed turns are then made ascending the foot; the roller is next carried directly

Fig. 61.

Bandage of foot covering the heel.

over the point of the heel and continued back to the dorsum of the foot; thence beneath the instep around one side of the heel and up over the instep; from this point it is car-

ried beneath the instep around the other side of the heel and up in front of the ankle, from which point it may be continued up the leg. (Fig. 61.)

Use.—This bandage is employed to cover in the foot and retain dressings to the foot and heel.

Bandage of Foot not Covering the Heel (French).

Roller Two and a Half Inches in Width, Seven Yards in Length.

Application.—Fix the initial extremity of the roller upon the leg just above the malleoli and secure it by two circular turns around the leg; the bandage is then carried obliquely across the dorsum of the foot to the metatarso-phalangeal articulation and at this point a circular turn

Fig. 62.

Bandage of foot not covering the heel.

around the foot is made. The roller is now carried up the foot, covering it in with two or three spiral reversed turns, and at this point a figure-of-eight turn is made around the ankle and instep; this should be repeated once, which will cover in the foot with the exception of the heel; the bandage may then be continued up the leg with spiral reversed turns. (Fig. 62.)

Use.—This bandage may be employed to secure dressings to the foot and is the one generally used to cover the foot in applying the spiral reversed bandage of the lower extremity.

Spiral Reversed Bandage of the Lower Extremity.

Roller Two and a Half Inches in Width, Seven Yards in Length.

Application.—The initial extremity of the roller is placed upon the leg just above the malleoli and secured by two circular turns, then carried obliquely over the foot to the metatarso-phalangeal articulation and here a circular turn is made around the foot; the foot is next covered in with two or three spiral reversed turns and two figure-of-eight turns of the ankle and instep, and just above the ankle one or two circular or spiral turns are made around the leg, and as the bandage is carried up the leg, as it increases in diameter, spiral reversed turns are made until it approaches the knee; at this point, if the limb is to be kept straight, the spiral reversed turns may be continued over this region and

Fig. 63.

Spiral reversed bandage of the lower extremity.

up upon the thigh. If the knee is to be bent, figure-of-eight turns may be applied until the knee is covered, and then the thigh may be covered with spiral reversed turns. (Fig. 63.) To cover in the thigh as well as the leg, two bandages of the dimensions before given will be required.

Care should be taken to keep the reverses in a line and not to make them over the spine of the tibia, as they may thus become painful to the patient.

Use.—This is one of the most frequently employed of the roller bandages; it is used to apply pressure to the lower extremity, to retain dressings, and to secure splints in the treatment of fractures and dislocations.

Figure-of-eight Bandage of the Leg.

Roller Two and a Half Inches in Width, Seven Yards in Length.

Application.—This bandage differs from the spiral reversed bandage of the lower extremity only in the fact that

Fig. 64.

Figure-of-eight bandage of the leg.

when the swell of the calf is reached, figure-of-eight turns are made around the leg instead of spiral reversed turns. In applying the roller, when the calf of the leg is reached,

the bandage is carried obliquely around the leg and brought in front of the leg and made to cross the starting-turn in the median line; these turns are repeated until the calf of the leg has been covered in, and the bandage is finished with one or two circular turns just below the knee. (Fig. 64.)

Use.—This bandage holds its place more firmly than the ordinary spiral reversed bandage of the leg, and may be employed in the treatment of ulcers of the leg in conjunction with strapping, where it is desirable to change the dressings at infrequent intervals and to allow the patient to walk about during the course of treatment.

SPECIAL BANDAGES.

Spiral Reversed Bandage of the Penis.

Roller Three-quarters of an Inch in Width, Thirty Inches in Length.

Application.—Fix the initial extremity of the roller by two circular turns around the penis close to the pubis; then

Fig. 65.



Spiral reversed bandage of the penis.

carry the bandage obliquely down to the corona glandis; from this point ascend the body of the penis by spiral reversed turns to the pubis and finish the bandage by two figure-of-eight turns around the neck of the scrotum and root of the penis; split the end of the bandage so as to form two tails and secure it by tying these around the root of the penis. (Fig. 65.)

Recurrent Bandage of Stump.

Roller Two and a Half Inches in Width, Five to Seven Yards in Length.

Application.—Place the initial extremity of the roller upon the anterior or posterior surface of the limb a few inches above the extremity of the stump, and carry the

Fig. 66.

Recurrent bandage of stump.

bandage to the end of the stump, and then conduct it upward or downward on the limb, as the case may be, to a point directly opposite to the point of starting; then bring the bandage back over the face of the stump to the point of starting and continue these recurrent turns, each turn overlapping two-thirds of the previous one, until the face

of the stump is covered; then reverse the bandage and secure the recurrent turns at their points of origin by two or three circular turns. The roller should next be carried obliquely down to the end of the stump and a circular turn should be made around this, and the bandage should next be carried up the limb by spiral or spiral reversed turns beyond the point at which the recurrent turns terminated, and secured by one or two circular turns. (Fig. 66.)

In applying this bandage in very short stumps resulting from amputations at or near the shoulder or hip-joints, after making the recurrent and spiral turns, it will be found necessary to carry the bandage, in the case of the shoulder, across the chest to the opposite axilla and back, and apply several of these turns; so in case of the hip amputations it will be found best to finish the bandage with a few turns about the pelvis.

Bandage for Securing the Hands and Feet in the Lithotomy Position.

Roller Two and a Half Inches in Width, Three Yards in Length.

Application.—The hand of the patient should be brought down and made to grasp the outer side of the foot; the initial extremity of the roller is fixed by two circular turns around the wrist and ankle, and the bandage is then passed around the foot and hand, and these turns are alternated with turns around the wrist and ankle, until the hand and foot are firmly secured. The same procedure is adopted with the hand and foot of the opposite side. (Fig. 67.)

Use.—This bandage is useful in securing the hands and feet while the patient is put in the lithotomy position for that operation, or for perineal section.

Liebreich's Eye Bandage.

This bandage consists of a strip of flannel two and a half inches in width and from six to ten inches in length, to the

extremities of which are sewed tapes. It may be applied transversely so as to cover both eyes, or obliquely so as to cover one eye only, and is secured by the tapes carried around the head and tied over the forehead. (Fig. 68.)

Fig. 67.

Bandage for securing hands and feet for lithotomy.

Fig. 68.

Liebreich's eye bandage.

Use.—This bandage is used to hold compresses or dressings to the eye or eyes, and the elasticity of the flannel permits of its being applied so as to make a variable amount of pressure.

Bandage of Scultetus.

This is a compound bandage, consisting of a number of pieces of muslin, and may be prepared from a two and a half or three-inch roller by cutting off strips long enough to encircle the part about one and one-third times. These strips are placed under the part in such a manner that the first piece shall be overlapped by the second, the second by the third, and so on from below upward; the pieces are then brought around the limb, and the extremities of the

last piece are secured by pins. (Fig. 69.) This bandage was formerly much employed in the treatment of compound fractures to secure dressings to the wound, and possessed the advantage that single strips which became soiled could be removed without disturbing the whole dressing, the new strip to be introduced being pinned to the extremity of the

Fig. 69.



Bandage of Scultetus.

soiled piece to be removed, and then being drawn through by its removal. This bandage will often be found convenient in applying dressings to cases of excision of the joints, where as little disturbance of the parts as possible is important in dressing the wounds. When the strips are attached to each other by a thread passed through each strip in the centre, the bandage is known as *Pott's bandage.*

This bandage is applied and secured in the same manner, but it possesses no advantages over the bandage of Scultetus.

The Rubber Bandage.

This bandage is made from a strip of rubber sheeting, from one inch to four inches in width and from three to five yards in length, which, for convenience of application, is rolled into a cylinder.

Its use was introduced to the profession by Dr. Martin, of Boston, and it will be found a useful form of dressing where it is considered desirable to apply elastic pressure to a part.

It may be employed in the treatment of varicose veins of the legs, in chronic ulcers of those parts where pressure is

Fig. 70.

Martin's rubber bandage.

an important element in the treatment, and may be used as a substitute for strapping to secure this object. Its application has also been recommended in the treatment of swelled testicle in that stage of the affection in which pressure is indicated.

Application.—For application to the leg a rubber bandage two and a half inches in width and three yards in length is required.

The initial extremity of the roller is fixed upon the foot near the toes and secured by a circular turn; the foot is then covered in by spiral turns overlapping each other about two-thirds, and a figure-of-eight turn is made from the ankle

to the instep, and the bandage is then carried up the limb to the knee with spiral turns, where it is secured by two tapes sewed to the terminal extremity of the bandage, which are passed around the leg and tied. The bandage need not be reversed, as its elasticity allows it to conform to the shape of the limb. Care should be taken not to apply these turns with too much firmness; the bandage should be stretched very slightly; if this precaution is not taken, it soon becomes uncomfortable to the patient. A patient using one of these bandages will soon learn to apply it himself, making just the requisite amount of tension to secure its holding its place and to insure a comfortable amount of pressure upon the part. A well-fitting stocking may be placed upon the limb before the bandage is applied, or it may be applied directly to the skin.

The bandage should be removed at night when the patient goes to bed and hung up to dry, as its inner surface becomes moist from the secretions from the skin; it should be reapplied as soon as the patient rises in the morning.

In using it in the treatment of ulcers of the leg no ointments should be applied to the ulcer, as oily dressings soon destroy the rubber; dressings may be made to the ulcer by means of dry powders, such as oxide of zinc, iodoform, or aristol, before the bandage is applied.

In the treatment of swelled testicle the bandage is applied to the testicle by means of recurrent turns not too firmly made, and secured in place by spiral turns, until the whole surface of the organ is covered in; the end of the bandage is secured with tapes tied around the root of the scrotum. The same precaution to apply the bandage so as to make only moderate pressure should here also be observed.

Fixed Dressings or Hardening Bandages.

For the application of these dressings a variety of substances are used which are incorporated in the meshes of some fabric, such as crinoline or cheese-cloth, or painted over its surface to give fixity or solidity to the bandage.

The materials most commonly used in the preparation of

fixed dressings are plaster-of-Paris, starch, silicate of sodium or potassium, paraffine, or a mixture of chalk and gum or of oxide of zinc and glue.

The Plaster-of-Paris Bandage.

The plaster-of-Paris used for the application of surgical dressings should be of the same quality as that which the dental surgeons employ in taking casts for teeth—that is, the extra-calcined variety. If moist or of inferior quality, it will not set rapidly or firmly, and will fail to give sufficient fixation to the dressing.

The plaster-of-Paris dressing may be applied in several ways, either by covering the part to be enclosed with some loose fabric, and rubbing the moist plaster into it, alternating the layers of the fabric with layers of moist plaster, or it may be applied by means of a roller which has been prepared with plaster-of-Paris and is moistened and applied to the part.

To apply a plaster-of-Paris dressing according to the first method, the part to be enclosed—the leg, for instance—should first be covered by a neatly applied flannel bandage or a muslin bandage, which has been shrunken by being washed; new muslin is not satisfactory as a primary application to a limb in applying a plaster-of-Paris dressing, as the moisture from the plaster wets it and causes it to shrink, so that it may exert injurious pressure after the bandage becomes dry.

The limb having been covered by the bandage, and any bony prominences such as the malleoli having been padded with small wads of cotton to prevent undue pressure upon them, the part is next covered by a layer of turns of a crinoline bandage or by strips of cheese-cloth or any other loose material. A small quantity of plaster-of-Paris is next mixed with water until it has the consistence of thick cream, when it is smeared evenly over the whole surface of the previously applied bandage. Another layer of the bandage or of strips is next applied, and the plaster is smeared over

this in the same manner, and so alternate layers of plaster-of-Paris and bandage are applied until a casing of the desired thickness is obtained. If the plaster-of-Paris of the quality previously described be used, it will set or become hard in a few minutes.

The most convenient method of applying the plaster-of-Paris dressing is that employed by Prof. Sayre, which consists in the use of bandages which have been previously prepared with plaster-of-Paris; these are moistened and applied while moist to the part to be encased.

Preparation of the Plaster-of-Paris Bandage.

These bandages are prepared by taking cheese-cloth, mosquito-netting, or crinoline, which latter is by far the best fabric, and cutting or tearing it into strips two and a half to three inches in width and five yards in length. These are laid on a table, and plaster-of-Paris of the quality before mentioned is dusted over them and rubbed into the meshes of the fabric; the material when impregnated with plaster is loosely rolled into a cylinder, and these bandages when prepared should be placed in air-tight jars or tin cans until required.

Bandages thus prepared, which have been exposed to the air or have been kept for a long time, are not apt to set well when applied; but if such bandages are placed in a hot oven and baked for half an hour before being used, they will be found to set as satisfactorily as those freshly prepared.

These bandages may be prepared by a machine made for this purpose, but I do not think that they are apt to have the plaster as evenly distributed through them, and therefore are not as satisfactory, as those prepared by hand.

Application of the Plaster-of-Paris Bandage.

Before applying this dressing, the part to be encased—the leg, for instance—should be covered by a flannel roller,

the bony prominences being protected by pads of cotton, or a closely fitting stocking may be applied to the part.

The bandage should be dipped in warm water and kept covered by water for a few moments; it may be squeezed with the hand, and as soon as bubbles of air cease to escape it is a sign that it is thoroughly soaked and is ready for application.

On removing it from the water the excess of water should be squeezed out by the hand and the bandage should then be evenly applied to the limb with just enough firmness to make it fit the part nicely, and as few reverses as possible should be made. A sufficient number of bandages are applied to make a dressing as firm as may be required; three rollers of the above dimensions are usually quite ample for a dressing for the leg, and when the last roller has been applied some dry plaster should be moistened with water until it has the consistence of thick cream, and it should be rubbed evenly over the surface of the bandage to give it a finish (Fig. 71). If a good quality of plaster has

FIG. 71.

Leg encased in plaster-of-Paris dressing.

been used, the bandage should be quite firm in from ten to fifteen minutes, but the patient should not for a few hours be allowed to put any weight upon the bandage.

An equally firm bandage may be applied with the use of a less number of bandages, if the surgeon rubs over the surface of each layer of bandage applied a little moist plaster, then applying another layer and repeating the same procedure; finishing the dressing by an external coating of moist plaster, as above described.

In applying these dressings a fewer number of bandages will be required if narrow strips of tin, zinc, or binder's board are incorporated in the layers of the bandage, which also increase the strength of the dressing.

Interrupted Plaster-of-Paris Dressing.

This form of plaster-of-Paris dressing is applied by first placing a short iron rod under the extremity some distance above and below the point at which the dressing is to be interrupted; this is fixed by a few turns of the plaster bandage above and below the portion of the limb which is to be left exposed; stout wire is next bent into loops, the

Fig. 72



Interrupted plaster-of-Paris dressing.

extremities of which are incorporated in the subsequent turns of the plaster bandage; three loops thus placed in addition to the posterior iron bar will usually make the dressing sufficiently firm (Fig. 72). A number of turns of the bandage are applied to firmly fix the loops, and the limb is held in the desired position until the plaster has set.

Application of the Plaster-of-Paris Jacket.

The patient's body should be covered with a soft, closely fitting woven shirt without arms, but with shoulder-straps to hold it in position, or an ordinary woven undershirt may be

employed; one or two folded towels, or a pad of cotton folded in a towel, are next placed over the abdomen between the shirt and the skin—this is called, by Prof. Sayre, the dinner pad, and is intended to leave space for the distention of the abdomen after eating. Small pads of raw cotton may

Fig. 73.

Suspensory apparatus.

also be placed over the anterior iliac spines, and, in the case of females, a pad of cotton wrapped in a handkerchief may be placed over each mammary gland.

The patient should next be suspended by the apparatus consisting of a collar and arm-pieces attached to a cross-bar (Fig. 73), which is attached by a cord and pulley to a tripod. If this apparatus is not at hand, a very satisfactory substitute

may be made by folding two towels into cravats and tying together the ends, so as to make two loops, one of which is placed in each axilla; a bar of wood two and a half feet in length is next taken and the loops are secured to the ends of this by stout cords or handkerchiefs; a Barton's bandage

Fig. 74.

Patient suspended for application of plaster jacket.

is next applied to the head, and a strip of bandage is passed under the turns which cross the vertex and is secured to the middle of the cross-bar. The bar is next suspended by a cord passed through a pulley or ring which may be attached to the sill of a door if the ordinary tripod is not to be obtained.

The patient should be slowly raised by the apparatus

until the toes only are in contact with the floor, and the extension should not be carried to the point which makes it uncomfortable to the patient. (Fig. 74.) The shirt should be drawn downward over the hips by an assistant and held in place until a few turns of the bandage have been applied.

The plaster bandage having been soaked and squeezed, a turn should be made around the body above the pelvis, and it should then be carried downward below the iliac spines, and from this point should be made to ascend gradually by spiral turns until it reaches the axillary line. The turns should be applied smoothly and not too tightly. After one or two layers of turns have been applied, the surgeon may rub some moist plaster upon their surface if he desires to use fewer bandages. These turns are repeated until a bandage of the desired thickness is applied, and the surface of the dressing may be finished by rubbing it over with moistened plaster. This jacket for a child will generally require the use of four bandages of the dimensions given; for an adult, six to eight bandages.

The patient should be kept suspended until the bandage has set, usually from ten to fifteen minutes, and then should be carefully lifted so as not to bend the spine, and placed upon his back upon a mattress, until the dressing becomes perfectly hardened. The dinner pad, and mammary pads, if they have been used, should next be removed. In applying this dressing, strips of zinc or tin may be placed between the layers of bandage if it is desired to give more strength to the dressing.

Application of the Jury-mast by Means of Plaster-of-Paris.

In disease of the spine involving the cervical or upper dorsal region the ordinary plaster-of-Paris jacket is not satisfactory, and in such cases the "jury-mast" is employed in connection with the plaster jacket. In applying the "jury-mast" the same steps are taken in the preparation of

the patient as in applying the plaster-of-Paris jacket, with the exception of extension, which need not be used.

After three or four layers of the plaster bandage have been applied to the body, an apparatus made of two bars of metal having two perforated strips of zinc attached to them a few inches apart, which partly encircle the body, is applied and held in position by turns of the plaster bandage. The perpendicular bars have at their upper part a slot, into which the lower end (Fig. 75) of the "jury-mast" fits, and is secured by a screw; to the upper part of this is attached a movable cross-bar, to which are fastened the straps of the collar from which the head is suspended.

FIG. 75.

Head-support and jury-mast.

THE BAVARIAN DRESSING.

To apply this dressing, which is sometimes employed in the treatment of fractures, take two pieces of Canton flannel the length of the part to be enclosed, and more than wide enough to envelope its circumference. In applying it to the leg these pieces should be cut so as to correspond to the outline of the leg and posterior portion of the foot.

These pieces should be placed one over the other and sewed together in the middle line, the seam corresponding to the back of the leg. The leg and foot are then placed upon this, and the inner layer of flannel is brought up in front of the leg and over the dorsum of the foot and made fast with pins. (Fig. 76.) Plaster-of-Paris is next mixed with water and made into a paste, which is rubbed thickly and evenly over the flannel next to the limb until a sufficient thickness is obtained; the outer layer of flannel is then

brought up about the leg and moulded to its surface by the hands. A loosely applied roller may now be used to hold the dressing in place until the plaster has set.

FIG. 76.

Bavarian dressing.

When it is necessary to inspect the parts, the turns of the roller are cut, and upon separating the layers of flannel the two halves can be turned aside, the seam at the back acting as a hinge. Upon reapplying the splints to the leg they may be retained in position by a roller or by one or two strips of bandage.

MOULDED PLASTER SPLINTS.

It is sometimes found difficult to apply the ordinary plaster dressings to parts irregular in their shape, and at the same time to have a splint which can be removed with ease. To accomplish this purpose moulded splints of plaster may be made by cutting a paper pattern of the part to be covered in, and then cutting pieces of crinoline to conform to this pattern; eight or ten pieces will usually form a splint of sufficient thickness. One of these pieces of crinoline is laid upon a table and dry plaster is rubbed into its meshes; another is laid upon this and plaster is applied to it in the same way, and so on until all the pieces have been placed in position, one over the other, with plaster rubbed well into the meshes. The dressing is then folded up and dipped into water, squeezed out, and moulded to the part and held in position, until it sets, by the turns of a bandage. The edges should overlap slightly, and in applying it a strip of waxed paper may be placed under the overlapping edge to prevent

its adhesion to the dressing below, and thus facilitate its removal. Splints prepared in this way can be removed with ease, and are often of service in cases where it is desirable to inspect the parts frequently; I have employed with advantage such splints in making fixation of the hip-joint in cases of coxalgia, and also for the same purpose in affections of other joints. The splints upon being reapplied are secured by a few strips of bandage, or by a roller bandage.

Trapping Plaster Bandages.

In applying the plaster-of-Paris dressing to a part where there is a wound which is covered by the plaster bandage, it is well to make some provision whereby the plaster dress-

Fig. 77.

Plaster-of-Paris bandage trapped. (Esmarch.)

ing over the site of the wound may be cut away, making a trap or window through which the wound may be inspected or dressed, if necessary. (Fig. 77.) To accomplish this, before applying the plaster bandage, a compress of lint or gauze should be placed over the wound, which, when the dressing is completed, forms a projection on its surface, indicating the position of the wound, and also allows the surgeon to cut away the dressing without injuring the skin below. These traps may be cut out after the bandage has partially set, or after it has become hard. In applying the plaster-of-Paris dressing in cases of compound fracture and

after osteotomy, I always make provision for trapping of the bandage if it should become necessary, although in the vast majority of cases it does not have to be done.

Removing Plaster-of-Paris from the Hands.

One objection to the use of plaster-of-Paris dressings is the difficulty of removing it from the hands of the surgeon, and the harsh condition in which the skin of the hands is left after its removal. If, however, the hands are washed in a solution of carbonate of sodium—a tablespoonful to a basin of water—the plaster will be readily removed and the skin will be left in a soft and comfortable condition.

Removing the Plaster-of-Paris Bandage.

The removal of the plaster bandage is sometimes a matter of difficulty, particularly if it has to be removed before the parts below it are consolidated, as it may disarrange them and cause the patient pain if it is not accomplished without much force.

When the bandage is applied to get a cast of the part, a strip of sheet-lead one inch in width is first placed over the flannel bandage, and is allowed to project at each end beyond the dressing; the plaster can then be readily cut through upon this strip with a knife without injury to the parts below.

It may also be removed by means of a saw devised for this purpose (Fig. 78), or by strong cutting shears of various

Fig. 78.

Hunter's saw for removing plaster bandages.

kinds (Fig. 79); or a line may be painted over the dressing with hydrochloric acid or vinegar, which softens the plaster so that it can readily be cut through with a knife. Dr. William

B. Hopkins has devised a vertebrated metal chain which is applied to the part before the plaster is applied and removed when the bandage has set, leaving a hollow longitudinal

Fig. 79.

Shears for cutting plaster bandages.

ridge which can be cut through or divided with a rasp. The saw is, I think, the most satisfactory means of removing these dressings; the only caution to be exercised is to use it carefully, as the final layers of the bandage are divided, to avoid wounding the skin.

Uses of Plaster-of-Paris Dressings.

These dressings are employed to secure fixation, as primary or secondary dressings in the treatment of fractures, and for a like purpose in injuries and diseases of the joints. They are also largely used in the treatment of diseases and deformities of the spinal column, and will also be found most satisfactory applications after osteotomy and tenotomy, to secure immobility and hold the parts in their corrected positions; when employed in dressing cases after tenotomy, they are generally used for a few weeks until the proper mechanical apparatus is applied.

The Starched Bandage.

To apply this bandage starch is first mixed with cold water until a thick creamy mixture results; to this is added boiling water until a clear mucilaginous liquid is produced; if too thin it can be made thicker by heating it upon a stove. The

part to be dressed is first covered with a flannel roller, and over this a few layers of a cheese-cloth or crinoline bandage, which has been shrunken, are applied; the starch is then smeared or rubbed with the hand evenly into the meshes of the material, and the part is again covered with a layer of turns of the bandage, and the starch is again applied; this manipulation is continued until a dressing of the desired thickness is produced. Strips of pasteboard may be applied between the layers of the bandage to give additional strength to the dressing, if desired.

It requires from twenty-four to thirty-six hours for the starched bandage to become dry and thoroughly set, and it may be removed in the same way in which the plaster-of-Paris dressing is removed.

Use. — Before the introduction of the plaster-of-Paris dressing it was formerly much employed in the treatment of fractures and in injuries of the joints. It may be used in such cases, but possesses no advantage over the former dressing and has the disadvantage of setting much less promptly.

Gum and Chalk Bandage.

In applying this dressing equal parts of powdered gum arabic and precipitated chalk are mixed with boiling water until a mass of the consistence of cream results. This is applied to the cheese cloth or crinoline bandage in the same manner as is the starch in the application of the starched bandage; it has the advantage over the latter dressing of setting more promptly, five or six hours only being required for it to become hard. It may be employed for the same purposes as the starched or plaster-of-Paris bandage.

Silicate of Potassium or Sodium Bandage.

In applying this bandage after a flannel roller and several layers of a cheese-cloth or crinoline bandage have been applied to the part, the surface of the latter is coated with silicate of sodium or potassium applied by means of a brush,

then a second layer of bandage is applied and treated in the same manner, and this manipulation is continued until a bandage of the desired thickness is produced. It requires twenty-four hours for this dressing to become firm. In removing the silicate bandage it may be first softened by soaking it in warm water and then it can be readily cut with scissors.

In applying either the starched bandage or the silicate of potassium bandage care should be taken to use cheese-cloth or crinoline which has been shrunken by being moistened and allowed to dry before being employed; otherwise dangerous compression of the part may occur if the bandage has been firmly applied and shrinks after its application.

The Paraffine Bandage.

Paraffine, which melts at from 105° to 120° F., is used in the application of this bandage. The limb being covered by a flannel roller, a vessel containing paraffine is placed in a basin of boiling water. As the roller, which may be either of flannel, cheese-cloth, or crinoline, is unwound it is passed through the melted paraffine and applied to the part, and the turns are repeated until a dressing of sufficient thickness results, and the surface may be brushed over with melted paraffine. This dressing sets very rapidly, being quite firm in from five to ten minutes.

It possesses the advantage of the other fixed dressings in that it does not absorb discharges and become offensive, and for this reason it was formerly recommended in the treatment of compound fractures.

Glue or Glue and Oxide of Zinc Bandage.

Glue or glue combined with oxide of zinc has been employed in the preparation of fixed dressings, but possesses no advantages over those previously mentioned.

Raw-hide or Leather Splints or Dressings.

In moulding raw-hide or leather splints it is necessary, first, to apply a plaster-of-Paris bandage to the part to which the raw-hide splint is to be fitted; as soon as the plaster has set it is removed, and a solid plaster cast is next made by pouring liquid plaster-of-Paris into this mould. When this has become dry a piece of raw-hide, which has been soaked for a time in warm water, is moulded to the cast and held firmly in contact with it by a bandage until it has become perfectly dry. It is then removed, and its surface is covered with several coats of shellac, to prevent its absorbing moisture from the skin when applied, and changing its shape. Eyelets or hooks are fastened to the edges of the splint, through which strings are passed to secure the splint in place.

Fig. 80.

Leather jacket with jury-mast.

Made in this manner raw-hide splints fit the part very accurately, and constitute a very satisfactory dressing for cases of joint-disease, and in the form of leather jackets are often employed in the treatment of disease of the spine in place of the plaster-of-Paris jacket. (Fig. 80.)

In the treatment of high dorsal

Fig. 81.

Leather splint for cervical caries (Owen.)

or cervical caries a leather splint in two sections, which rests upon the shoulders and supports the head, is often used with good results. (Fig. 81.)

Binder's Board or Pasteboard Splints.

This material, which can be obtained in sheets of different thickness, is frequently employed for the manufacture of splints. In moulding these splints a portion of the board of the requisite size and thickness is dipped in boiling water for a short time, and when it has become softened it is removed and allowed to cool; a thick layer of cotton batting is next applied over it, and it is then moulded to the part and held firmly in place by the turns of a roller bandage; in a few hours it becomes dry and hard.

This material, from its cheapness and the ease with which it is obtained, is frequently employed to mould splints for the treatment of fractures, especially in children, and for the fixation of joints in the treatment of acute and chronic joint affections. A moulded pasteboard splint is also often employed to fix the ends of the bones after the excision of a joint.

Porous Felt Splints.

This material is also employed for the manufacture of splints, and is applied by dipping the material in hot water and then moulding it to the part; as it dries it becomes hard.

Hatter's Felt Splints.

Hatter's felt is also frequently employed for the manufacture of splints or dressings. It is softened by dipping it in boiling water or heating it in the flame of an alcohol lamp, and when soft and pliable it is moulded to the part, and as it cools it again becomes hard.

These splints are employed for the same purposes as those made of plaster-of-Paris, leather, or pasteboard.

PART II.

MINOR SURGERY.

Theory of Asepsis and Antisepsis in Wound Treatment.

THE term *Asepsis*, applied to wounds, implies that there is in the wound an absence of those vegetable parasites or microörganisms whose presence sets up fermentative changes, accompanied by suppuration and constitutional disturbance.

Antisepsis, on the other hand, has reference to the means employed to bring about the destruction of microörganisms which may be present in the wound or upon the instruments, dressings, or hands of the surgeon, and which, if not destroyed or rendered inert, will set up fermentative changes in the wound.

It has long been a well-recognized fact that albuminoid substances, such as dead animal tissue, blood, or blood-serum, will, when exposed to moisture, warmth, and the presence of certain living organisms or fungi, bacteria and micrococci, develop putrefactive changes; and if these changes take place in the living body there result certain constitutional disturbances known as symptomatic, inflammatory, or septic fever.

It was also recognized that these putrefactive changes in albuminoid substances could be avoided by their exposure to heat, cold, or by drying—any of these conditions being sufficient to destroy or arrest the development of the micrococci. The microörganisms which set up fermentative and putrefactive changes in animal tissues exist in great variety,

but those which are of most interest to the surgeon belong to the *cocci* and *bacilli*.

Rosenbach's investigations have shown that the most common cause of suppuration in living human tissue is a minute globular micrococcus, to which the name *staphylococcus pyogenes aureus* has been given. This coccus is found in almost all varieties of acute suppuration.

Another form of coccus which may exist alone or in connection with the previously mentioned fungus is the *staphylococcus pyogenes albus*. Both of these varieties of cocci, from the agminated arrangement of the single coccus, are known as grape cocci, and have the peculiarity of causing well-localized foci of inflammation.

The *streptococcus pyogenes*, a pus-generating chain coccus, which extends rapidly along the lymph spaces and lymphatics, and by rapid infiltration of the tissues causes spreading gangrene, is also of especial interest to the surgeon.

Decomposition in tissues, accompanied by the presence of foul odors, is said always to be due to the action of rod-like bodies called bacilli or bacteria, such as the *bacillus pyogenes fœtidus* and *bacillus pyocyaneus*.

In wounds the result of accident or made by the surgeon, we have present conditions most favorable for the entrance and development of these organisms, such as the serum and blood, and the dead or partially devitalized cells of the various tissues which are exposed. We have present also warmth and moisture, and in the air coming in contact with the wound we have vast quantities of dust laden with spores, which under these favoring conditions develop into the organisms before mentioned, which rapidly set up fermentative processes known as decomposition.

The products of this decomposition, carried into the circulation by the lymphatics and veins, set up local changes in the shape of inflammation and at the same time give rise to systemic disturbances which we recognize as septic fever.

Modern wound treatment aims at the prevention of decomposition and suppuration, and accomplishes this purpose by having the wound kept aseptic, by perfect cleanliness of

the region of the wound, the hands and instruments of the surgeon, and by not exposing the wound to an atmosphere which contains dust; as the latter condition is difficult to obtain we secure the destruction of the microörganisms which may be present by heat, as seen in the use of the actual cautery, and by chemical sterilization, which is accomplished by the use of germicides.

Methods and Dressings Employed in Wounds to Secure Asepsis.

To prevent infection of wounds the various chemical sterilizers and dressings are employed in different ways, and the principal types of dressings are as follows:

Method by Simple Drying.

This method is employed in small and not very deep wounds. The edges having been brought together by sutures the surface of the wound is dusted with powdered iodoform, the serum and blood forming with this, as it dries, a scab, which protects the wound from infection from without, and repair takes place promptly under this scab. Iodoform collodion may be employed instead of powdered iodoform in this method of dressing.

Method by Drying and Chemical Sterilization.

The object of this method of dressing is to provide a means of sterilizing the blood or serum which escapes from the wound, and at the same time to insure the sterilization of the air coming in contact with the discharges or the wound.

It is employed in large or deep wounds, where there is always more or less escape of blood or serum, and is accomplished by applying a number of layers of sublimate or iodoform gauze and sublimated cotton over the wound. Evaporation not being interfered with, the whole dressing becomes hardened, and the wound is surrounded by a large

antiseptic crust made up of the dressing and serum or blood.

This method of dressing is the one most generally employed at the present time.

Moist Dressings.

In this method of dressing, the wound is covered by moist gauze dressings and these are kept moist and evaporation is prevented by applying over them some impervious material such as mackintosh or rubber tissue.

Modified Moist Dressing.

In this method of dressing, the wound itself is covered by a piece of protective or rubber tissue; over this is placed the sublimated or iodoform gauze dressing and some layers of bichloride cotton. By this method of dressing, the wound itself is kept in a moist condition favoring particularly the organization of blood clots; the external dressings become dry as the discharges which have escaped into them evaporate, forming an antiseptic crust or covering over the wound.

Surgical Cleanliness.

In the practice of aseptic surgery it is a matter of the first importance that anything coming in contact with a wound should be absolutely clean, such as the hands of the operator, instruments, sponges, towels, ligatures, sutures and dressings.

Antiseptic Substances Employed.

A great variety of substances possessing more or less germicidal properties have been at different times employed in the practice of aseptic or antiseptic surgery; those most employed at the present time are bichloride of mercury, carbolic acid, iodoform, beta-naphthol, chloride of zinc, peroxide of hydrogen, creolin, permanganate of potassium, pyoktanin and boric acid, the double cyanide of mercury and zinc, and aristol.

Bichloride of Mercury.

This is employed as an antiseptic in watery solution, varying in strength from 1 : 500 to 1 : 5000.

The solution 1 : 500 to 1 : 1000 is used only for the irrigation and disinfection of the hands and skin; for the irrigation of wounds, a solution of 1 : 2000 is generally employed. In using the bichloride solution in operations upon children, I am in the habit of using a solution of 1 in 4000, and I find that it produces less irritation of the skin and is equally efficient as a germicide. Where continuous irrigation is kept up or where it is employed in large cavities, a still weaker solution, 1 : 5000 to 1 : 10,000, should be employed.

In using these solutions the surgeon should watch the patient carefully for symptoms of poisoning due to the absorption of the bichloride of mercury; the symptoms denoting this are vomiting, fetid breath, salivation, inflammation of the gums, diarrhœa, blood-stained stools, and bleeding from the mouth and nose.

In preparing the solutions of bichloride of mercury for use, it will be found convenient to have a concentrated solution of the salt in alcohol, one part of the bichloride of mercury to ten parts of alcohol; this can be kept in a well-stoppered bottle, and to this should be added one teaspoonful of common salt, which prevents the disintegration of the mercuric compound. One teaspoonful of this solution added to one quart of water makes a 1:1500 solution.

A 10 per cent. bichloride solution may be made as follows:

Bichloride of mercury	2 parts.
Sodium chloride	1 part.
Dilute acetic acid	1 "
Water	16 parts.

By adding water in an appropriate quantity, 1 : 1000 or 1 : 2000 solution can be made.

Or the solution may be prepared with tartaric acid in the proportion of five parts of the acid to one part of the bichloride of mercury, the following formula being employed:

Hydrarg. chlor. corrosiv.	grs. xv.
Ac. tartaric.	grs. lxxv.
Aquæ dest.	Oij.

Pellets containing a definite amount of bichloride of mercury compounded with a few grains of common salt or muriate of ammonia, which, when dissolved in a definite quantity of water, make a solution of 1 : 1000 or 1 : 2000, will also be found very convenient for the preparation of solutions.

These bichloride or sublimate solutions are also employed to sterilize the gauze and cotton which are largely employed in antiseptic dressings.

Carbolic Acid.

This drug is employed in solutions of 1 : 20 or 1 : 40. The stronger solution, 1 ; 20, is usually employed to sterilize the instruments, the latter being allowed to remain in this solution for thirty minutes before being used. As a carbolic solution of this strength benumbs and cracks the hands of the operator, it should be diluted just before the instruments are required, by adding an equal quantity of water, making it a 1 : 40 solution.

The 1 : 40 solution is used for the irrigation of wounds and the washing of sponges. Carbolic acid is also employed in the preparation of gauze. A ready method of making a 5 per cent. carbolic solution is to add one tablespoonful of carbolic acid to one quart of water.

In using carbolic acid solutions the surgeon should be on the watch for the symptoms of poisoning, which will show itself by dark-colored urine, headache, dizziness, vomiting, and in severe cases bloody diarrhœa, hæmoglobinuria, and death from collapse. Carbolic acid solutions should be used with great caution in young children, as they seem to be more susceptible than adults to the constitutional effects of this drug. I have seen the use of quite dilute solutions produce the characteristic symptoms of poisoning in such patients.

Iodoform.

Iodoform has been shown by experimental research to possess little germicidal action, but in spite of this fact clinical experience has proved that it possesses powerful antiseptic properties, due not to the destruction of germs, but to its undergoing a decomposition in their presence, and thus rendering the ptomaines which have resulted from the germ-growth inert. Iodoform may be rendered absolutely sterile by washing it in a 1 : 1000 bichloride solution, which destroys all microörganisms; it should then be dried, and kept for use in closely stoppered bottles.

Iodoform is very extensively employed as an application to wounds; it is especially useful as a dressing to infected wounds, and to tubercular or syphilitic ulcers. It is also employed in the preparation of iodoform gauze, and may be combined with collodion to form *iodoform collodion*, which is a useful dressing in superficial wounds:

Iodoform	grs. xlviij.
Collodion	℥j.

An ethereal solution of iodoform (iodoform grs. xv, ether ℥j) is also used as an application to chronic ulcers.

An *emulsion of iodoform* in glycerin (iodoform ℥j, glycerin ℥x) is much employed at the present time as an injection in the treatment of chronic or tubercular abscesses.

Elderly persons are more prone to the toxic action of iodoform than young persons. These symptoms are manifested by sleeplessness, debility, headache, delirium, and death may result from meningitis or cardiac depression.

Beta-naphthol.

Beta-naphthol, in a 1 : 2500 solution, is employed for much the same purposes as the bichloride of mercury solution; it is not, however, so powerful a germicide. It is employed in irrigating large cavities because it is not a poisonous agent, but is especially useful as a bath for instru-

ments, as it does not corrode them, as does the sublimate solution. It also possesses the advantage over a carbolic acid solution of not irritating the skin of the surgeon's hands.

Chloride of Zinc.

Chloride of zinc, in a solution of 30 to 40 grains to water f℥j, is a very powerful antiseptic. When employed upon raw surfaces it produces marked blanching of the tissues; it is especially useful in wounds which are infected or which have been exposed to infection. I have found it by all means the best application to the poisoned wounds which are received in dissecting dead bodies and in operating. In such cases the whole cavity or surface of the wound should be washed with a 30-grain solution of the chloride of zinc, and then the wound should be dressed with a bichloride dressing.

Sulpho-carbolate of Zinc.

This drug has been found to possess more decided antiseptic properties than the chloride of zinc, and is much less irritating. It may be used in the same strength and for the same purposes as the former drug.

Peroxide of Hydrogen.

This drug is employed in what is known as a 15-volume solution, which may be diluted from 10 per cent. upward or used in full strength. It is employed in the sterilization of sinuses or suppurating cavities, such, for instance, as often result from diseases of or operations upon bone. It seems to have a direct action upon pus generation by destroying the microörganisms of pus. It is injected into sinuses and cavities by means of a syringe, or may be applied to open wounds in the form of a spray; its action is shown by the escape of bubbles of air, and it should be used as long as these continue to escape.

Kreolin or Creolin.

This substance is obtained from English coal by dry distillation, and has been found to possess powerful germicidal properties; it is non-irritating and practically non-toxic. It is insoluble in water, but forms an emulsion with it which possesses marked germicidal properties. It is employed for the same purposes as carbolic acid, and has the advantage over the latter drug that it is not irritating to the skin, and is almost devoid of toxic properties.

It is used in an emulsion, in strength from two to five per cent., and is employed in the irrigation of large wounds or cavities of the body, and has been most favorably recommended in gynecological practice. As a bath for instruments to render them sterile during operations it is useful, but the opacity of the emulsion makes it difficult to find the instruments and interferes with its efficiency.

Boric Acid.

This drug has not very marked antiseptic qualities, but is unirritating even in saturated solutions. It is frequently employed in a 5 to 30 per cent. solution to cleanse and disinfect mucous surfaces and large cavities. It is often employed to wash out the bladder before the operation for the removal of calculi or growths from that organ.

In the dressing of superficial wounds, or in wounds in which the bichloride or carbolic acid dressings produce irritation, an ointment of boric acid, made by taking boric acid 1 part, vaseline 5 parts, will be found very satisfactory.

Boro-salicylic Lotion.

This lotion is prepared by adding 2 parts of salicylic acid and 12 parts of boric acid to 1000 parts of hot water. This forms a very bland solution, which can be used where there is danger in using bichloride or carbolic solutions—as, for instance, in the bladder or peritoneal cavity.

Permanganate of Potassium.

This drug, owing to its rapid absorption of oxygen, acts as an antiseptic, and is often employed for the disinfection of foul wounds and ulcers. It is also employed in solution for washing the operator's hands, and for the washing of sponges. It is pratically non-irritating, and may be used in quite concentrated solutions, but is usually employed in the following solution: Permanganate of potash ℥j, water f℥j. One fluid drachm of this solution to a pint of water makes a 1 : 1000 solution.

Pyoktanin.

Methyl-violet, known in commerce under the name of pyoktanin, has been recommended as a drug possessing marked antiseptic powers. It is said to prevent suppuration by destroying the organisms which are active in its production, which are said to have an affinity for and are killed by aniline colors. It has been claimed that it sterilizes the pus of suppurating wounds and ulcers, and it is recommended as an injection in the treatment of large suppurating cavities for this purpose, as it is practically non-poisonous.

It is employed in a solution of a strength of 1 : 1000 or 1 : 2000, and for the sterilization of surgical instruments a 1 : 10,000 solution may be employed. When employed as a means of irrigating wounds, it should be used until the tissues are of a deep-blue color. Recent investigations have shown that it is, as a germicide, much less reliable than bichloride of mercury.

Aristol.

Aristol, which is a compound of iodine and thymol, possesses germicidal properties and has been introduced as a substitute for iodoform. It has the advantage over iodoform of not being poisonous and is also without disagreeable odor. It may be employed for the same purposes as iodoform and

it seems to be particularly useful as a dressing to chronic and specific ulcers.

Double Cyanide of Mercury and Zinc.

Cyanide of potassium, cyanide of mercury, and sulphate of zinc are mixed together in solution, in quantities proportioned to the atomic weights of 2KCy, $HgCy_2$ and $ZnSO_4 + 7H_2O$; the cyanide of potassium and cyanide of mercury being dissolved together in one and a half ounces of water for every 100 grains of potassium cyanide, are added to the sulphate of zinc dissolved in three times that amount of water. The precipitate is collected and washed in two successive portions of water equal in quantity to that used for the solutions, that is six ounces of water for every 100 grains of the potassium cyanide, to free the precipitate from the irritating salts associated with it in its formation. The precipitate being well washed, is next mixed with distilled water containing one part of hæmatoxylin for every 100 parts of the cyanide salt; this, when it precipitates the cyanide salt, changes its color to a pale bluish tint.

Ammonia is next added in such a proportion to the mixture that one fluidrachm of the ammoniacal liquid shall correspond with one grain of hæmatoxylin, and the ammoniacal mixture is allowed to stand for three or four hours, when it is filtered and the dyed salt is drained and dried at a moderate heat, is next levigated and may then be kept for any length of time for use. When employed for charging gauze it is mixed with a 1 : 4000 bichloride solution in the proportion of four pints of the solution to 100 grains of the salt.

Preparation of Materials Used in Aseptic Surgery and Dressings.

Sponges.

Sponges, while dry, should be beaten to free them from calcareous matter, then placed in a 15 per cent. solution of hydrochloric acid for thirty minutes to dissolve any lime

which may remain in them; they should then be removed from this solution and washed, and should next be well washed with green or castile soap and warm water for a few minutes and then thoroughly rinsed and placed in a 1 : 1000 bichloride solution or in a 5 per cent. carbolic solution in closely covered jars until required for use.

Or, after beating the sponges to remove any sandy matter, they may be placed for twenty-four hours in a solution of hydrochloric acid—hydrochloric acid ℥iv, water four pints—then removed and washed until free from acid, then steeped for half an hour in a solution of permanganate of potassium, 180 grains to six pints of water. Next wash them and place them in the following solution: hyposulphite of sodium, ℥x; hydrochloric acid, f℥v; water, f℥lxviij; and allow them to remain in this solution for four hours; remove them from this and place them in running water for six hours; they should then be placed in jars and covered either by a 5 per cent. carbolic acid solution or a 1 : 1000 bichloride solution. The carbolic acid solution is better for keeping the sponges than the sublimate solution, as it does not decompose.

They may be prepared also by beating and washing them, and then soaking them for twelve hours in a solution of chlorinated soda — chlorinated soda 1 part, water 5 parts. They are then removed and well rinsed and placed in a 5 per cent. carbolic solution, or they may be placed in a moderately warm oven until thoroughly dry, and then placed in air-tight jars, if it is desired to keep them dry.

It is better to use a cheaper grade of sponges, and to use them only once, but if the same sponges are to be used again, they should be well washed in a solution of carbonate of soda, 1 ounce to the quart, and then placed in a 1 : 1000 bichloride solution.

Silk.

Silk for sutures or ligatures should be sterilized by boiling for thirty minutes in a 5 per cent. solution of carbolic acid or water, then placed in stoppered bottles and covered with a 5 per cent. solution of carbolic acid or with absolute alcohol.

Silkworm-gut.

Silkworm-gut is an excellent material for sutures, and is much easier to thread than the silk or catgut. It may be kept dry in glass jars, or preserved in alcohol, and should be placed in a 5 per cent. carbolic solution for a few minutes before being used, as this renders it more supple.

Catgut Ligatures or Sutures.

Juniper Catgut.

Catgut, varying in size from No. 0, which is very fine, to No. 4, which is quite thick, is placed in oil of juniper berries for one week, and is then transferred to absolute alcohol, in which it should be kept until required for use.

No. 1 catgut is the size usually employed for ligatures and sutures.

Alcohol is the best material in which to preserve the catgut, as it keeps it firm, and does not interfere with its flexibility, while both carbolic acid and bichloride solutions render it brittle and weak.

Chromic Acid Catgut.

The catgut is first washed in alcohol and placed in 1 quart of a 5 per cent. solution of carbolic acid, containing 30 grains of bichromate of potassium, and is allowed to remain for forty-eight hours. This immersion should be longer when large-sized varieties of catgut are used; but for the sizes of catgut which are ordinarily used, this time of immersion will prepare the gut to resist the action of the living tissues for a week or more. Catgut thus prepared may be dried and placed in closely stoppered jars, or may be kept in alcohol.

Catgut may also be prepared by soaking it in alcohol for a short time, and then placing it in the following solution for forty-eight hours: Chromic acid, 1 grain; carbolic acid,

200 grains; alcohol, 2 drachms; water, 2½ ounces. It is then removed and placed in glass jars for use.

Before being used it should be soaked for thirty minutes in a 5 per cent. carbolic acid solution, or in a 1 : 1000 bichloride solution.

The chromic acid catgut is by far the best variety of gut to use for sutures and for the ligation of the larger vessels in their continuity.

Drainage-tubes.

The drainage-tubes usually employed are prepared from rubber tubing of different sizes perforated at short intervals;

Fig. 82.

Rubber drainage-tube.

the black rubber tubes are softer and more pliable than the red or white rubber tubes, and should be preferred. (Fig. 82.) Drainage-tubes are also made of glass (Fig. 83), which

are almost exclusively used in abdominal surgery, and also of decalcified bone. Drainage-tubes should be kept in a 5 per cent. solution of carbolic acid, or, if kept dry, they

Fig. 83.

Glass drainage-tube.

should be well washed and placed in a carbolic or bichloride solution for thirty minutes before being used.

Horse-hair and Catgut for Drainage.

Catgut as ordinarily prepared for ligatures may be used to secure drainage in small and superficial wounds; a number of strands of catgut are placed in the bottom of the wound, and the end or ends are allowed to project from one or both extremities of the wound.

Horsehair may be employed for the same purpose, a number of strands of the hair being placed in the wound in the same manner. Before being used it should be well washed with soap and water and then soaked in a 5 per cent. carbolic solution or 1 : 1000 bichloride solution for thirty minutes.

Protective.

Protective is employed to prevent the wound from being irritated by the antiseptic substances with which the gauze is impregnated or by its irregular surface.

Various materials are employed as protectives, the principal requirement being that it is some tissue which can be readily rendered aseptic, and does not absorb any irritating materials from the dressings.

The protective first employed by Mr. Lister, which is still generally employed, is prepared by coating oiled silk with

copal varnish, and when this is dry a mixture of 1 part of dextrine, 2 parts of powdered starch, and 16 parts of a 1 : 20 carbolic acid solution is brushed over its surface.

Rubber tissue may be employed very satisfactorily as a substitute for this protective.

Before applying the protective to the wound, it is dipped into a solution of bichloride of mercury or carbolic acid.

Mackintosh.

This consists of cotton cloth, with a thin layer of India-rubber spread on one side. It is employed in antiseptic dressings as a layer of dressing outside of the gauze, and should be applied with the rubber surface toward the wound, to prevent the discharge from the wound from soaking directly through the dressing.

The mackintosh cloth is not at the present time as much employed as formerly, unless the moist method of dressing is adopted.

Rubber Tissue.

This consists of a very thin sheet of India-rubber with glazed surfaces, which can be obtained from the rubber manufacturers; it is employed for the same purposes as the mackintosh, is much less expensive, and, as previously stated, may be used instead of protective for covering the wound.

Parchment Paper.

This consists of a very tough paper material which can be soaked in a watery solution of corrosive sublimate or carbolic acid without becoming so much softened as to tear upon handling. It is prepared by the manufacturers of surgical dressings, and is employed for the same purposes as mackintosh.

Gauze Dressings.

The most convenient and cheapest material for wound-dressing is a sheer material known in the trade as cheese or tobacco cloth. By reason of having a very open mesh it absorbs well either the materials with which it is prepared or the discharges from the wound when applied as a dressing. It can be readily obtained anywhere, is inexpensive, and is soft and pliable, so that it is a comfortable form of dressing to the patient. The gauze is impregnated with different materials to render it antiseptic, and its preparation is a matter of little difficulty.

Preparation of Gauze Dressing.

Bichloride of Mercury or Corrosive Sublimate Gauze.

In preparing bichloride or corrosive sublimate gauze, thirty yards of cheese-cloth are placed in a wash-kettle and covered with water, to which is added two pounds of washing soda or a pint of lye, and boiled for an hour; the soda or lye is added to remove any oily matters which the cheese-cloth contains, and thus make it more absorbent. The gauze is next removed from the water and washed in clear water, and passed through a clothes-wringer, and then immersed in a 1 : 1000 bichloride of mercury solution for twenty-four hours. It is then dried and cut into pieces several yards in length, and packed in closely covered glass jars or tin boxes and put away for use. Or it may be preserved as moist gauze by packing it in air-tight jars. If gauze has been prepared for some time, it is well to soak it for a short time in a 1 : 1000 bichloride of mercury solution before using it.

In using the sublimate gauze on delicate skins there will sometimes result a dermatitis which is known as mercurial eczema; this is particularly apt to occur if the gauze is

moistened or covered with rubber tissue or mackintosh. If this condition develops, the parts covered by the gauze should be rubbed over with boric acid ointment or vaseline before it is reapplied, or another variety of dressing may have to be substituted, such as the iodoform or carbolized gauze.

Iodoform Gauze.

Iodoform gauze may be prepared by sprinkling cheese-cloth, which has been boiled in soda solution, with powdered iodoform and rubbing it well into its meshes; it should then be dried and packed in glass jars for use.

It may also be prepared by rubbing an emulsion of iodoform, made by adding 3 drachms of iodoform to 6 ounces of Castile soap-suds, into 18 ounces of moist gauze; this should be dried and packed in glass jars for use.

Double Cyanide of Mercury and Zinc Gauze.

The preparation of this gauze is much more difficult than that of the other varieties of gauze, requiring the following:

Potass. cyanide	130 grains.
Mercuric cyanide	251.7 "
Zinc sulphate	268 9 "
Hæmatoxylin	1.3 "
Sal ammonia (gas NH_3 1 per cent.)	6 minims.
Gauze (previously boiled and dried)	10 ounces.
Bichloride of mercury solution	7.6 pints.
Distilled water	q. s.

In charging gauze with this substance, 100 grains of the salt are dissolved in 4 pints of a 1 : 4000 bichloride solution, which will give from 2 to 3 per cent. of the cyanide to the dry gauze. The gauze should be freshly prepared and used moist, or if allowed to become dry it should be moistened again with a weak bichloride solution before being used. The advantages claimed for this gauze are that it is not irritating to the skin, and as the antiseptic is not soluble it is not washed out by the discharges from the wound.

Prof. J. William White, who has used it extensively in his practice, considers that it possesses decided advantages over the bichloride gauze.

Carbolized Gauze.

The carbolized gauze which is used in the University Hospital is prepared in the following manner: Cheese-cloth, which has been previously boiled and dried, is soaked for a few hours in the following solution:

Resin	1 pound.
Alcohol	5 pints.
Castor oil	24 ounces.
Carbolic acid	12 "

The gauze is next removed from this solution and passed through a clothes-wringer, and is then cut in pieces four to six yards in length, which are folded and packed in air-tight tin boxes for use.

Sawdust Dressing.

Sawdust is impregnated with a 1 : 1000 bichloride of mercury solution for twenty-four hours, and then spread out to dry; after it has become sufficiently dry it is enclosed in bags made of cheese-cloth of various sizes. This will be found to be a satisfactory substitute for the ordinary gauze. In using this dressing the wound should be covered with a piece of protective or a few layers of gauze, and the bags are then packed over this and held in place by a bandage.

Moss Dressing.

Different species of *sphagnum* or moss, on account of their cheapness, elasticity, and great absorbing power, have been found a very satisfactory material with which to make dressing bags.

Clean moss is soaked for twenty-four hours in a 1 : 1000 bichloride of mercury solution and then dried; cheese-cloth

bags are filled with this material and may be used dry or may be moistened with 1 : 3000 bichloride solution before being applied in the dressing of wounds. They are much employed in the hospitals of Germany, and have largely superseded the gauze dressings.

Improvised Antiseptic Dressings.

In cases of emergency, when the ordinary gauze dressings cannot be obtained, it is well to remember that old muslin or linen, which can usually be obtained, will serve for a temporary dressing if properly sterilized, until a more elaborate dressing can be applied. Old sheets either of muslin or linen should be torn into pieces half a yard square and thrown into boiling water; after remaining for a few minutes in this they should be removed and soaked for a few minutes in a 1 : 1000 or 1 : 2000 bichloride solution, or a 5 per cent. carbolic solution, and applied to the wound, a number of layers of this material being applied and held in position by a bandage. This dressing will keep the wound aseptic until a more elaborate dressing is obtained.

Antiseptic Bandages.

These bandages are prepared by tearing or cutting bichloride or carbolized gauze into strips two to three inches in width and five yards in length, and forming the strips into rollers and packing them in air-tight vessels.

The bandages may also be prepared from boiled dry gauze in the same manner, and are kept in air-tight boxes or jars until required for use, when they are soaked for a few minutes in a 1 : 1000 bichloride or 5 per cent. carbolic solution.

They may also be prepared from crinoline, the same material which is used for the plaster-of-Paris bandage; as this material is quite stiff, the bandage should be soaked in a bichloride or carbolic solution before being applied, and as the material contains a certain amount of starchy matter, it becomes firm as it dries and makes a very secure dressing.

For this reason it is often applied over the ordinary antiseptic bandage.

Bichloride Cotton.

This material, which is an important part of most antiseptic dressings, is prepared by soaking absorbent cotton in a 1 : 1000 bichloride of mercury solution for twenty-four hours, and then allowing it to dry. When dry, it is packed in jars or air-tight boxes. Its great absorbing power and

Fig. 84.



Sterilizing oven.

its elasticity make it, when properly prepared, a most valuable dressing; it is generally employed to cover the gauze dressing, a number of layers being applied.

Borated, carbolized, and salicylated cotton, prepared in the same manner, are also frequently employed for a similar purpose.

Dry Sterilized Dressings.

These dressings are prepared by sterilizing ordinary gauze with steam. Gauze cut into proper lengths is placed in wire cases and exposed to super-heated steam in an oven for a few hours, and is then dried in another oven, removed, and placed in air-tight jars or boxes The apparatus required for perfect sterilization of dressings is expensive, and is not likely to be employed by practitioners, but is used in hospitals where a large number of dressings are constantly required. A convenient form of sterilizing oven is shown in Fig. 84. Unless the sterilization is perfect, these dressings should not be employed; the same method is employed in the sterilization of instruments.

Preparation for Aseptic Operation.

Surgical Cleanliness.

The Hands.

The hands and forearms of the surgeon and of his assistants should be well washed in hot water with soap for a few minutes, and a nail-brush should be used to cleanse the region of the finger-nails; rings with irregular surfaces which might retain filth should be removed. After the hands and forearms have been thoroughly cleansed, they should be immersed in 1 : 1000 bichloride solution for a short time.

The same precautions should be taken with the hands and forearms of nurses who handle the instruments and dressings. If in any manner the hands of the surgeon or of his assistants come in contact, during the operation, with any objects which have not been disinfected, such as the clothing

of the patient, the operating-table, etc., it is a matter of the first importance that they should be thoroughly washed and disinfected before again being brought in contact with the wound.

Sterilizing of Instruments.

The instruments should be carefully scrubbed with warm water and soap, care being taken to see that all joints and roughened surfaces are freed from any dry matter which may adhere to them; after being thoroughly cleansed in this manner, they should be placed in a metal or porcelain tray and covered by a 5 per cent. carbolic solution for fifteen minutes before being used.

The instruments which are now constructed with metal handles may be sterilized by placing them in boiling water, or by boiling them; where instruments are employed which have wooden handles this method of sterilization cannot be employed, and here it will be found necessary to resort to the first method of sterilization.

Instruments which fall upon the floor or come in contact with the clothing of the surgeon or the patient during the operation, should be washed and placed in the carbolic solution before again being brought in contact with the wound.

Preparation of the Patient for Operation.

The patient having been prepared for the operation by whatever constitutional treatment the surgeon considers necessary, the region of the proposed wound is first rubbed over with cotton saturated with spirits of turpentine, and next is thoroughly washed with soap and water; if hairs are present in the region they should be shaved off; after a careful washing with soap and water, the skin is carefully washed with a 1 : 1000 bichloride or 5 per cent. carbolic solution, and is then covered with a towel wrung out in a 1 : 2000 bichloride solution until the surgeon is ready to begin the operation.

This cleansing of the region of the proposed wound in

hospital practice is generally made a few hours before the operation, but in private practice it has to be done just before the operation is undertaken; but if carefully done the results will be in no wise less satisfactory. In private practice the operation may have to be performed while the patient is in his bed, or, if an ordinary kitchen-table is at hand, it will be found more convenient to place him upon this; this should be prepared by placing upon it a folded quilt or blanket, and over this a sheet of rubber cloth, and upon this is laid a clean, folded, linen or muslin sheet. The surgeon should carry with him a sheet of this rubber cloth, three by four feet, which he will find most useful in preparing the table for the operation, or in protecting the bed of the patient if he is not placed upon the table; a rubber cloth of this size takes up little space if carefully folded, and can be easily packed in the instrument-bag.

Details of an Aseptic Operation.

The patient being anæsthetized and placed upon the table, the clothing is so arranged as to expose freely the part to be operated upon; the clothing or the skin surrounding this region is next covered with towels wet with a 1 : 1000 bichloride solution. If any considerable surface of the patient's body is covered by these towels, to avoid chilling the surface and adding to the shock which naturally follows the operation, they should be wrung out in a hot bichloride or carbolic solution, and should be replaced as they become cold by hot towels prepared in the same manner. The patient being ready for operation, the surgeon should assign the assistants and nurses their duties, and having again immersed their hands and forearms in the bichloride solution the operation is begun.

During the operation the wound is irrigated frequently with a 1 : 2000 or 1 : 3000 bichloride solution, which may be allowed to run over the wound, or be applied by means of a syringe or irrigating apparatus (Fig. 85), and the hands of the surgeon and assistants should also be washed in this solution at not too long intervals. In prolonged operations,

or in those in which a large wound is made, I think it is especially important that the irrigating solutions should be used as warm as can be comfortably borne by the hands of the surgeon; warm solutions, it has been shown by recent

FIG. 85.

Irrigating apparatus. (ESMARCH.)

investigations, possess a greater germicidal power than those of the same strength when used cold, and they also possess the advantage of preventing the chilling of the patient, and thus diminish the shock of the operation.

Hemorrhage during the operation is controlled by the use of hæmostatic forceps, which are applied to the bleeding vessels, or the vessels may be ligatured as they are divided. After the operation has been completed, and all hemorrhage has been controlled, the wound is thoroughly irrigated with a 1 : 2000 or 1 : 3000 bichloride solution.

The next step is to provide for drainage; this may be disregarded in small or superficial wounds, but in a wound of any considerable size or depth it is safer to provide free drainage. This is accomplished by the use of perforated rubber drainage-tubes, or a number of strands of catgut or horsehair, or by decalcified bone or glass drainage-tubes.

The rubber tube in large wounds will be found most comfortable to the patient and satisfactory as regards drainage; it may be laid in the wound, the ends being allowed to extend from the extremities of the wound, or it may be so introduced that one end of the tube rests in the deepest part of the wound and the other extremity is brought out of the wound at its most dependent portion; in large or irregularly shaped wounds a number of tubes may be required to secure free drainage. The ends of the drainage-tubes are transfixed with safety pins which have been sterilized and allowed to remain in a 5 per cent. carbolic solution until required, and the ends of the tube should next be cut off close to the pins so as to be as nearly as possible flush with the skin.

The wound is next closed by the introduction of sutures, which may be of silkworm-gut, chromicized catgut, silk, or silver wire; the needles and sutures should be soaked in a 5 per cent. carbolic solution for thirty minutes before being used. The wound being closed, a final irrigation of its deepest parts should be made, by injecting a stream of bichloride solution, 1 : 2000 or 1 : 3000, into the end of the drainage-tube; if through-and-through drainage has been employed, one end of the tube should be closed and the solution should be injected into the wound through the other end of the tube by means of a syringe or irrigating-tube, until the wound is slightly distended with the solution, which allows the latter to find its way to all parts of the cavity of the wound. The external surface of the wound and the skin for some distance surrounding it should next be washed with a 1 : 2000 bichloride solution, and a piece of protective, a little longer and wider than the wound, is next dipped in a bichloride or carbolic solution and placed over it, and over this is laid the deep dressing, which consists of a pad of bichloride gauze from eight to sixteen layers in thickness and large enough to overlap the wound two or three inches in all directions. This should be dipped in a 1 : 2000 bichloride solution, and wrung out as dry as possible before being applied. The superficial gauze dressing is next applied, and consists of sixteen layers of gauze, which should be large enough to extend from three to

six inches beyond the wound in all directions; this gauze is applied dry. Over the superficial gauze dressing there is next applied a number of layers of bichloride cotton, so arranged as to extend a little beyond the margin of the superficial gauze dressing. These dressings are now secured in position by the application of a gauze bandage, which is prevented from slipping by the introduction of a few safety-pins.

The dressing being completed, the patient is moved from the operating-table to his bed, and care should be exercised to see that the dressings do not become soiled if the patient vomits upon coming up from the anæsthetic.

In this method of dressing no mackintosh or rubber tissue is employed, outside of the superficial gauze dressing; the discharges from the wound are disseminated through the dressing and become dry by evaporation, and the dressing forms an antiseptic scab which covers and surrounds the wound.

Moist Method of Dressing.

If, for any reason, it is desired to adopt the moist method of dressing, a piece of mackintosh or rubber tissue larger than the superficial gauze dressing is placed over it, and over this is placed a few layers of bichloride cotton, care being taken to see that the layers of cotton overlap the mackintosh or rubber tissue by a few inches; the application of an antiseptic gauze bandage then completes the dressing. On the removal of this dressing the gauze will be generally found to be soaked with the discharges from the wound, and in a moist condition. The disadvantage of this variety of dressing is that there is apt to be more irritation of the skin set up by the bichloride gauze when kept moist than when applied in the manner of a dry dressing.

Reapplication of Dressings.

The re-dressing of a wound which remains aseptic need not be made for some days; if the temperature remains

normal or a little above this point, and the patient exhibits no unfavorable constitutional symptoms, and the dressing is comfortable to the patient, it need not be disturbed for a week or ten days; at the expiration of this time it is well to examine the wound and to remove the drainage-tube if a drainage-tube has been used, and to remove a portion or all of the sutures if the superficial parts of the wound are firmly healed.

In re-dressing an aseptic wound at the end of a week or ten days, to prevent any possible infection, as much care should be exercised as in the original dressing of the wound. The patient's clothes should be removed so as to freely expose the dressing, and a rubber cloth should be placed under the patient so as to protect the bed, and the clothing and skin in the region of the wound should be protected by towels wrung out in a 1 : 1000 bichloride solution. The surgeon should wash his hands and immerse them in a 1 : 1000 bichloride solution before removing the dressings. The bandage retaining the dressing should be divided with bandage scissors and the dressings should be removed layer by layer, and when the deep dressing is removed care should be taken to see that the drainage-tubes are not pulled upon if they are adherent to the dressing; the protective should next be removed, and the surface of the wound should be irrigated with a 1 : 2000 bichloride solution; the drainage-tubes should next be inspected to see that they are free, and a stream of bichloride solution may be passed through them by means of a syringe. If the wound is found aseptic the drainage-tube may be removed, and the wound should next be irrigated through its track by a stream of bichloride solution. If the wound is healed the sutures may be removed at this dressing; but if the wound has been an extensive or deep one it may be well to remove only a portion of the sutures; if animal sutures have been employed, they need not be removed. The surface of the wound is next washed with a 1 : 2000 bichloride solution and a piece of protective is placed over the line of incision, and the deep and superficial dressings are applied as previously described and covered with layers of bichloride cotton, and

the whole dressing is secured by the application of an antiseptic bandage. If the wound remains aseptic after this dressing, the dressings need not be changed for a week or ten days, and at this time the wound will usually be found healed, so that further dressings are not required.

If, however, the wound is not running the typical course of an aseptic wound, constitutional symptoms will be developed, as evidenced by a rise in the temperature and pulse-rate and other constitutional disturbances. In this event the wound should be re-dressed as soon as possible, and if the cause of the disturbance can be found it should be removed; for instance, hemorrhage may have taken place into the wound, and the blood not being able to escape through the drainage-tubes may have caused so much distention of the wound that the vitality of the skin covering the wound is threatened, or the sutures may be found to be causing irritation, or suppuration may be found to be present.

If, on exposure of the wound, it is found that it is distended with blood-clots, and blood is escaping from the wound, the sutures should be removed, the clots should be turned out, and the bleeding vessel or vessels should be sought for and ligatured, and the wound, after a thorough irrigation with 1 : 2000 bichloride solution, should be drained and closed with sutures, and dressed as previously described.

If, however, on exposure of the wound, and upon the removal of a portion or all of the sutures, the wound is found distended with a blood-clot, and no evidence of hemorrhage at the time exists, or of suppuration in the wound, the clot may be allowed to remain in place, and the wound should be re-dressed as in the original dressing, trusting to the organization of the blood-clot if it has remained aseptic. If the patient's condition improves after the dressing, and the temperature and pulse-rate become normal, it is an indication that the wound is still aseptic, and it need not be re-dressed for some days.

If, on the other hand, examination of the wound shows that the drainage is insufficient, or that the drainage-tubes are occluded by blood-clots, these should be removed by washing out the tubes with a 1 : 2000 bichloride solution by

means of a syringe, and introducing additional drainage-tubes, if it is deemed necessary; the wound should then be re-dressed.

When it is found on examination of the wound that suppuration is present, the surgeon may adopt one of two methods of treatment: he may thoroughly wash out the wound through the drainage-tubes with a 1 : 2000 bichloride solution, and after thorough irrigation of the wound re-dress it, and, if the patient's constitutional symptoms improve, he may be assured that the wound has been rendered aseptic, and is running an aseptic course.

If he does not feel that this method of treatment is sufficient, he may open the wound and wash it thoroughly with a 1 : 2000 bichloride solution, and next apply to its surface a 15-volume solution of the peroxide of hydrogen, which may be diluted with water one-third or one-half, or a 30-grain solution of chloride of zinc may be used; and after this application a final irrigation with the 1 : 2000 bichloride solution shall be made, and it should then be drained, closed, and dressed, as previously described.

If the treatment instituted to render the wound aseptic has been successful, the patient's constitutional condition will improve, and it will heal as an aseptic wound.

Dressing of Septic Wounds.

It often happens that patients suffering from wounds which have been improperly treated, or have had no treatment, come under the care of the surgeon; such wounds are already infected, and to render them aseptic, if possible, the following treatment should be adopted: The skin surrounding the wound should be carefully washed with spirits of turpentine, and then with soap and water, and finally with a 1 : 2000 bichloride solution. The wound itself should be next exposed as fully as possible, and any foreign bodies which are found in it, or dirt, should be removed with forceps and a stream of water; it should next be thoroughly irrigated with a 1 : 2000 bichloride solution, and then should be drained, closed, and dressed as an operation wound.

If suppuration is already present, after cleansing the region of the wound it should be irrigated with a 1 : 2000 bichloride solution, and should next be washed with a 30-grain solution of chloride of zinc, or a 15-volume solution of the peroxide of hydrogen diluted with water one-half, one-third, or used in its full strength; and finally thoroughly irrigated with a 1 : 2000 bichloride solution.

The introduction of drainage-tubes and sutures will depend upon the character of the wound. Sutures cannot often be used with advantage if much retraction of the skin has occurred, and a drainage-tube is not required if the wound is left open. A piece of protective is next applied over the surface of the wound, and the deep gauze dressing, wrung out in a 1 : 2000 bichloride solution, is applied over this, and the superficial gauze dressing and bichloride cotton are next applied and secured by a bandage.

Infected wounds which are treated in this manner will often be rendered aseptic, and in their subsequent course will be perfectly satisfactory, both to the patient and to the surgeon.

Materials Used in Surgical Dressings—*Continued.*

Lint.

This material is employed in surgical dressings, and is of two varieties; the domestic lint, which consists of pieces of old linen or muslin which have been thoroughly washed or boiled and then dried, or the surgical lint which is manufactured by machinery, and resembles Canton flannel in appearance; the latter is the best material, as it has a greater absorbing capacity.

Lint is used as a material on which unctuous preparations are spread in the dressing of wounds, and is also employed as a material for saturating with the various solutions which are used in wet dressings, such as lead-water and laudanum, or dilute alcohol; the lint, after being saturated with these solutions, is covered with rubber tissue or oiled silk when

applied, to prevent too rapid evaporation of the solution. It is also one of the best materials from which to construct compresses employed in the treatment of fractures, to control hemorrhage, or to make pressure for any purpose

Paper-lint, made from old rags or wood pulp, has great absorbing power for fluids, and may be used as a substitute for surgical lint in the application of wet dressings to surfaces when the skin is unbroken.

Oakum.

This material, made from old tarred rope, was formerly much employed in the dressing of wounds before the introduction of the antiseptic method of wound-treatment; it was supposed to possess some antiseptic properties due to the tar with which it was impregnated. From its elasticity it is found to be an excellent material for padding splints or other surgical appliances. It is also employed in the form of pads to place under patients to relieve portions of the body from pressure, or to absorb discharges which soak through the dressings. A mass of oakum which has been well teased out and wrapped in a towel forms an excellent pillow on which to support a stump. The *oakum seton* is highly recommended by Dr. Sayre as a means of making a direct application of ointments to sinuses of bone; the oakum is loosely twisted into a cord and covered with any ointment desired and is passed through the sinuses in the bone; the position of the seton is changed from time to time, fresh ointment being applied before it is drawn through; resin cerate is a favorite application to these sinuses made in this manner.

Cotton.

Cotton is now employed in surgical dressings principally as a material to pad splints or to relieve salient parts of the skeleton from pressure in the application of splints or bandages; for instance, in the application of the plaster-of-Paris bandages, the bony prominences are generally covered by

small masses of cotton; it possesses but little absorbent power unless used in the form of absorbent cotton, and is not much employed in surgical dressings, except for the purposes mentioned above.

Absorbent Cotton.

This material is prepared from ordinary cotton, which is boiled with a strong alkali to remove the oily matter which it contains. When so prepared it absorbs liquids freely, and by reason of its great absorbing capacity it is largely employed in surgical dressings. A small mass of absorbent cotton wrapped upon the end of a probe or stick is now generally employed to make applications to wounds, and has taken the place of the sponge or brush which was formerly employed for this purpose. From its cheapness, after one application it can be thrown away and a new piece can be used, and thus the danger of carrying infection from one wound to another by the applicator is abolished. It is largely employed in gynecological practice for making applications to the female genital organs.

It is impregnated with various antiseptic substances, such as the bichloride of mercury, carbolic acid, boric acid, and salicylic acid, and, when thus treated, forms the *bichloride*, *carbolized*, *borated*, and *salicylated cotton* so much employed in antiseptic dressings.

Jute.

This substance is made from the fibre of the *Corchorus capsularis*, which, on account of the character of its fibre, possesses both elasticity and absorbing qualities; it has been employed for much the same purposes as oakum and cotton, such as the padding of splints, and is also used as an external absorbing dressing.

Oiled Silk or Muslin.

These materials are employed as an external cover for moist dressings to prevent rapid evaporation from the dress-

ings; they form excellent materials for this purpose, but as they are quite expensive their use is limited.

Waxed or Paraffine Paper.

This dressing is prepared by passing sheets of tissue-paper through melted wax or paraffine, and then allowing them to dry for a few minutes. Paper thus treated forms an excellent and cheap substitute for oiled silk or muslin, and may be employed for the same purpose for which the latter materials are used.

Rubber Tissue.

This material, which is prepared by rubber manufacturers, consists of rubber run out into very thin sheets; it has a glazed surface, is very pliable and strong at the same time, and forms a cheap and satisfactory substitute for oiled silk, and is employed for the same purposes. In the moist method of antiseptic dressing it may be used in place of the mackintosh, and indeed I prefer it to the latter in this method of dressing.

Parchment Paper.

This paper is prepared so as to render it water-proof; it is employed in surgical dressings for the same purposes as oiled silk and rubber tissue.

Compresses.

Compresses are prepared by folding pieces of lint, muslin, linen, or flannel upon themselves so as to form firm masses of variable sizes; oakum or cotton may also be used to form compresses. Compresses are employed to make pressure over localized portions of the body, as in the treatment of fractures, or to make pressure upon vessels for the control of hemorrhage.

TENT.

This consists of a small portion of lint, oakum, or muslin rolled in a conical shape, which is employed to keep wounds open and facilitate discharges. This dressing is not much employed at the present time, its use being largely superseded by the drainage-tube.

RETRACTORS.

Retractors are made by taking a piece of muslin four inches wide and twelve to eighteen inches in length and

FIG. 86.

Two-tailed retractor.

FIG. 87.



Three-tailed retractor.

splitting it as far as the centre, thus making a *two-tailed retractor*. (Fig. 86) A *three-tailed retractor* is made in

the same way, except that the muslin is slit twice instead of once. (Fig. 87.) Retractors are used to retract the soft parts in amputation, to prevent their injury by the saw in the division of the bones. When one bone is sawed a two-tailed retractor is used, and when two bones are sawed a three-tailed retractor is employed.

Plasters.

The varieties of plaster which are most commonly employed in surgical dressings are *adhesive* or *resin plaster*, *isinglass plaster* and *rubber adhesive plaster*.

Resin Plaster.—This plaster, which is machine-spread, is one of the most widely employed plasters in surgical dressings; the spread surface is covered with a layer of tissue-paper, which should be removed before it is used; it is cut into strips of the required width and length, and the strips should be cut lengthwise from the roll of plaster, as the cloth upon which it is spread stretches more transversely than in a longitudinal direction. When heated and applied to the surface it holds firmly; it is prepared for application by applying the unspread side to a vessel containing hot water, or it may be passed rapidly through the flame of an alcohol lamp.

This is the variety of plaster which is generally used in making the extension apparatus for the treatment of fractures, for strapping the chest in fractures of the ribs and sternum, for strapping the pelvis in cases of fractures of the pelvic bones, or for strapping the breast, the testicle, ulcers, or joints.

Rubber Adhesive Plaster is made by spreading a preparation of India-rubber on muslin, and has the advantage over the ordinary resin plaster that it adheres without the application of heat. It is employed for the same purposes as resin plaster, but when applied directly to the skin it is apt to produce a certain amount of irritation, and for this reason when it is to be continuously applied for some

time, as in the case of an extension apparatus, it is not so comfortable a dressing as that made from resin plaster.

Isinglass Plaster, which is made by spreading a solution of isinglass upon silk or muslin, will be found to be a most useful dressing in the treatment of superficial wounds. It is made to adhere to the surface by moistening it, and when used in the treatment of wounds it should be moistened with an antiseptic solution; it is in this way rendered antiseptic and may be used with safety in connection with other antiseptic dressings. The best form of this plaster is spread on muslin, and when properly applied adheres as firmly and possesses as much strength as the ordinary resin plaster.

Before using any of these plasters, if the part to which they are to be applied contains hairs, these should be shaved off, otherwise traction upon these, if the plaster is used for the purpose of extension, or in its removal, will cause the patient discomfort or pain.

Soap Plaster. — Soap plaster for surgical purposes is prepared by spreading *emplastrum saponis* upon kid or chamois. It is not employed for the same purposes as the resin or rubber plaster, as it has little adhesive power, and is used simply to give support to parts or to protect salient portions of the skeleton from pressure. It is found a most useful dressing when applied over the sacrum in cases of threatened bedsores, and may be applied for the same purpose to other parts of the body where pressure sores are apt to occur.

In the treatment of sprains of joints a well-moulded soap-plaster splint secured by a bandage will often be found a most efficient dressing, and in the treatment of fractures the comfort of the patient is often materially increased by applying small pieces of soap plaster over the bony prominences, upon which the splints, even when well padded, are apt to make an undue amount of pressure.

Strapping, or applying pressure to parts by means of strips of plaster firmly applied, is a procedure often employed in surgical practice.

Strapping the Testicle.

In strapping the testicle strips of resin plaster are usually employed; a dozen or more strips one-half an inch wide and twelve inches in length will be required.

The scrotum should be first washed and shaved, and the surgeon next draws the skin over the affected organ tense by passing the thumb and finger around the scrotum at its upper portion, making circular constriction; a strip of plaster which has been heated is passed in a circular manner around the skin of the scrotum above the organ, and is tightly drawn and secured; this isolates the part and prevents the other strips from slipping. Strips are now applied in a longitudinal direction, the first strip being fastened to

Fig. 88.

Strapping the testicle. (Smith.)

the circular strip and carried over the most prominent part of the testicle, and is then carried back to the circular strip and fastened. A number of these strips are applied in an imbricated manner until the skin is covered (Fig. 88), and the dressing is completed by passing transverse strips around the testicle from its lowest portion to the circular strip; care should be taken to see that no portion of the skin is left uncovered.

Strapping the testicle is employed with advantage in the subacute stage of orchitis or epididymitis, and it will be found a useful means of applying pressure to the scrotum after the injection treatment of hydrocele. As the swelling of the testicle diminishes the strips become loose, and the part will require re-strapping.

Strapping of the Breast.

To strap the breast, strips of resin plaster two inches wide and long enough to pass from the opposite shoulder under the breast to the point of starting are required. In applying the strips the end of the strip is placed on the spine of the scapula of the side opposite the diseased breast and is carried forward over the shoulder and obliquely downward under the breast and axilla, and then over the back to the point of starting; the first strip being applied in this manner, the next one is applied in the same direction, overlapping about one-third of the previous strip (Fig. 89). These oblique strips are applied in an imbricated manner until a sufficient number have been used to cover in the breast, or the oblique strips may be alternated with circular strips passing from the sternum over the breast to the spine. A sufficient number of strips are used to cover the breast and to make firm compression upon it. Strapping of the breast in this manner will be found a satisfactory method of treatment in chronic inflammatory conditions of the breast, where it is of advantage to support the breast and make compression at the same time; it has the advantage over the use of a bandage to support and compress the breast, that it does not interfere with the chest motions upon the opposite side of the body.

Fig. 89.

Strapping the breast. (Smith.)

Strapping of the Chest.

To strap one-half of the chest, strips of resin plaster two and a half inches wide, and long enough to extend from the spine to the median line of the sternum are required—eighteen to twenty inches in length. The first strip is heated and one

extremity is placed upon the spine opposite the lower portion of the chest; it is then carried over the chest and its other extremity is fixed upon the skin in the median line of the sternum. Strips are next applied from below upward in the same manner, each strip overlapping one-third of the preceding one, until the axillary fold is reached (Fig. 90); a second layer of strips may be applied over the first, if additional fixation is desired, or a few oblique strips may be employed.

Fig. 90.



Strapping of the chest.

Adhesive straps applied in this manner very materially limit the motion of the chest-wall upon the affected side, and are frequently employed in the treatment of fractures and dislocations of the ribs, in contusions of the chest, and in cases of plastic pleurisy when the motions of the chest-walls are extremely painful to the patient.

Strapping of Ulcers.

To strap ulcers of the leg, strips of resin plaster one and a half inches wide, and long enough to extend two-thirds around the limb, are required. The ulcer should be thoroughly cleansed, and the skin surrounding it should be well dried; the first strip, being heated, is applied obliquely to the long axis of the leg about two inches below the ulcer, and is carried two-thirds around the limb; another strip is applied to a corresponding point of the skin on the opposite side of the limb, and is carried obliquely over the limb, crossing the first strip in the median line, and is carried two-thirds of the way around the limb; alternate strips are thus applied until the ulcer is covered in, and the strips are carried several inches above the ulcer (Fig. 91). Care should be taken to see that the strips are so applied as not to meet or cover the entire circumference of the limb, as by so doing injurious circular compression may result. Chronic ulcers

upon other portions of the body may be strapped in the same manner.

Strapping of leg ulcers is usually reinforced by the application of a firmly applied spiral reversed or spica bandage of the lower extremity.

FIG. 91.

Strapping of ulcer of leg. (LISTOR.)

Strapping of ulcers of the leg applied in the manner described will be found a most satisfactory method of treating chronic ulcers in this location in patients who have to work during the course of treatment; the strips need only be removed at intervals of a week, and, if well applied, the dressing is generally a comfortable one to the patient.

STRAPPING OF JOINTS.

Strips of resin plaster two inches in width and long enough to extend two-thirds around the joint are required. The first strip is applied a few inches below the joint, and strips are then applied over this, each strip covering in two-

thirds of the preceding one until the joint is covered in and the strips extend a few inches above the joint.

The ankle-joint is strapped by taking strips of resin plaster one inch in width; the first strip is placed over the heel, and its ends are brought forward until they meet over the dorsum of the foot; a second strip encircles the foot and secures the ends of the first strip. These strips are alternately applied, each strip covering in one-half of the previous strip until the foot and ankle are covered in.

Strapping of joints will be found a satisfactory dressing in the treatment of sprains of joints in their chronic state.

Strapping of a Carbuncle.

To strap a carbuncle strips of resin plaster one to one and a half inches in width are required; these strips are applied at the margin of the swelling and are laid on concentrically until all except the central portion is covered. If a number of openings exist, the strips are so placed as not to cover these. Strapping applied in this manner in the treatment of carbuncle is often a comfortable dressing for the patient, and at the same time the concentric pressure favors the extrusion of the slough.

Poultices.

This form of dressing was formerly much employed in the treatment of inflammatory conditions and injuries as a means of applying heat and moisture to the part at the same time, and although the use of poultices is now very much restricted since the introduction of the antiseptic method of wound treatment, yet I think there are still conditions in which their employment is both useful and judicious.

They are often employed with advantage in inflammatory affections of the chest and of the abdominal organs, and in inflammatory affections of the joints and of bone, combined with rest, their action is often most satisfactory; in cases of

gangrene their employment hastens the separation of the sloughs.

They constitute a form of dressing which conduces much to the comfort of the patient in cases of deep suppuration by their relaxing effect upon the tissues, and their previous use does not prevent the surgeon from using all antiseptic precautions in the opening and drainage of these abscesses and the employment of antiseptic dressing in their subsequent treatment.

Flaxseed Poultice.

This poultice is prepared by adding first a little cold water to ground flaxseed and then adding boiling water, and stirring it in until the resulting mixture is of the consistency of thick mush. A piece of muslin is next taken which is a little larger than the intended poultice, and this is laid upon the surface of a table and the poultice mass is spread evenly upon it with a spatula or knife from one-quarter to one-half an inch in thickness; a margin of the muslin of one or one and a half inches is left, which is turned over after the poultice is spread, and serves to prevent it from escaping around the edges when applied. The surface of the poultice may be thinly spread over with a little olive oil, or may be covered with a layer of thin gauze to prevent the mass from adhering to the skin.

It is now applied to the surface of the skin and is covered with a piece of oiled silk, rubber tissue, or waxed paper, and held in position by a bandage or a binder.

Bread Poultice.

This poultice is prepared from stale wheaten bread, the crusts being discarded and the crumb only being used; this is moistened with boiling water and allowed to soak for a few minutes, when the excess of water is poured off and the mass is spread upon a piece of muslin or linen, as before described.

Charcoal Poultice.

In preparing this poultice flaxseed-meal and powdered charcoal in equal parts are mixed together, and by adding boiling water a poultice mass is produced, which is spread upon muslin, as previously detailed. It is better to use animal charcoal in making this poultice, as it possesses greater disinfecting power than vegetable charcoal. This poultice is used as an application to gangrenous parts, as it possesses marked disinfecting properties.

Fermenting Poultice.

This poultice may be prepared by adding yeast, two tablespoonfuls, to a mixture of flaxseed with hot water, making a thin poultice mass, and allowing it to stand for a few hours in a warm place; it rises and becomes light, and is then spread upon muslin and applied as required. A few ounces of porter or a piece of yeast cake may be used as a substitute for the yeast in preparing this poultice; charcoal may also be added to it to increase its disinfectant power. This poultice was formerly and is still used as an application to gangrenous parts to hasten their separation and to diminish the odor arising from the necrotic tissues.

Oakum Poultice.

This is prepared by soaking a mass of loosely picked oakum in hot water, wringing it out and covering it with a layer of cheese-cloth or antiseptic gauze. It is next applied to the part and covered with oiled silk or rubber tissue, and held in place by a bandage; it has a large capacity for the absorption of discharges.

It may be wrung out in a warm bichloride solution, or carbolic solution, and thus form an *antiseptic* poultice.

Hot Fomentations.

Hot fomentations are employed to keep up the vitality of parts which have been subjected to injury, as seen in severe

contusions resulting from railway or machinery accidents; also to combat inflammatory action. Flannel cloths, several layers in thickness, or surgical lint should be soaked in water having a temperature of 120°; these are wrung out and placed over the part and covered with waxed paper or rubber tissue; a second cloth should be in the hot water, ready to apply as soon as the first-applied cloth begins to cool, and so by continuously reapplying them the part is kept constantly covered by a hot dressing. The use of these hot fomentations may in many cases have to be continued for hours before the desired result is obtained. *Hot compresses* applied in this manner are frequently employed in treating inflammatory conditions of the eye, and are also of the greatest service in keeping up the vitality of parts which have been subjected to severe injury interfering with their blood-supply. I have seen contused limbs, which were cold and seemed to be doomed to gangrene by reason of diminished blood-supply, have their temperature and circulation restored by the patient and persistent use of this dressing. After the vitality of such a part is restored it should be covered with cotton and a flannel bandage and surrounded by hot-water bags or hot-water cans.

Irrigation.

This may be accomplished by allowing the irrigating fluid to come in contact with the wound or inflamed part, or by allowing the cold or warm fluids to pass through rubber tubes which are in contact with or surround the part; the latter method is known as *mediate irrigation*.

In employing irrigation in the treatment of wounds or in inflammatory conditions, a funnel-shaped can with a stop-cock at the bottom, or a bucket is suspended over the part at a distance of a few inches (Fig. 92), or a jar with a skein of thread or lamp-wick arranged to act as a siphon may be employed. (Fig. 93.) The can or jar is filled with water, and this is allowed to fall drop by drop upon the part to be irrigated, which should be placed upon a piece of rubber sheeting

so arranged as to allow the water to run off into a receptacle so as to prevent the wetting of the patient's bed. The water employed may be either cold or warm, and this is decided by the indications in special cases, and if it is

FIG. 92.



Apparatus for continuous irrigation. (ESMARCH.)

desired to make use of antiseptic irrigation the water is impregnated with carbolic acid or bichloride of mercury; a 1:5000 to 1:10,000 bichloride solution, or a 1:60 carbolic acid solution, is frequently employed with good results.

Antiseptic irrigation employed in this manner will be found a most useful method of treating lacerated and con-

tused wounds of the extremities in which the vitality of the tissues is much impaired; and in such cases the warm water should be preferred to cool water, the temperature being from 100° to 110°.

FIG. 93.



Irrigating apparatus. (ERICHSEN.)

Under the use of warm irrigation it is surprising to see how tissues apparently devitalized regain their vitality; the absence of tension from the non-introduction of sutures and firm dressings, and the warmth and moisture kept constantly in contact with the wound by this method of irrigation, are the important factors in the attainment of this favorable result.

Mediate Irrigation.

In this method cold or warm irrigation is applied to the part by means of cold or warm water passing through a rubber tube in contact with the part. A flexible tube of India-rubber half an inch in diameter, with thin walls, and sixteen or twenty feet in length, is applied to the limb like a spiral bandage, or is applied in a coil to the head, breast, or joints and held in place by a few turns of a bandage; the end of the tube is attached to a reservoir filled with cold or warm water above the level of the patient's body, and

the water is allowed to flow constantly through the tubing and escape into a receptacle arranged to receive it. (Fig. 94.)

Fig. 94.



Cold coil applied to arm. (Esmarch.)

Cold Water Dressings.

These dressings are applied by bringing the cold water either directly in contact with the part or by applying it by means of a rubber bag or bladder.

The temperature of the water may vary from cool water to that of ice-water.

These dressings are employed in local inflammatory conditions; a favorite method of employment of this dressing is by means of *cold compresses*, which are made of a few layers of surgical lint, dipped in water of the desired temperature and applied to the part; they are renewed as soon as they become warm. When it is desirable to have the compresses very cold, they may be laid upon a block of ice or in a basin with broken ice; to obtain the best

results from their employment they should be renewed at very short intervals.

A convenient method of applying cold without moisture is by the use of the *ice-bag*. This is either a rubber bag or bladder, which is filled with broken ice and applied to the part. In using an ice-bag it is better to cover the part first with a towel or a few layers of lint or muslin, which prevents the surface from becoming wet by absorbing the moisture which condenses upon the surface of the bag or bladder, and thus renders the dressing more comfortable to the patient. The ice-bag is often employed as an application to the head in inflammatory conditions of the brain or membranes, and is also used upon the surface of the body to control internal hemorrhage.

Counter-irritation.

Counter-irritants are substances employed to excite external irritation, and the extent of their action varies according to the material used and duration of its application; superficial redness or complete destruction of the vitality of the parts to which they are applied may result.

The use of counter-irritants under favorable circumstances is found to have a decided effect in modifying morbid processes, and they are widely employed as local revulsants in cases of congestion or inflammation, and in cases of collapse for their stimulating effect.

Rubefacients.

These agents, by reason of their irritating properties when applied to the skin, produce intense redness and congestion.

Hot Water.—When it is desired to make a quick impression upon the skin, the application of muslin or flannel cloths wrung out in hot water and renewed as rapidly as they become cool will soon produce a superficial redness of the integuments.

Spirits of Turpentine.—This drug applied to the skin is a very active counter-irritant; it may be rubbed upon the surface of the skin until redness results. When used in patients whose skin is very delicate, its action may be modified by mixing it with equal parts of olive oil before applying it; this will be found a useful precaution in applying it as a rubefacient to the tender skins of young children.

When redness of the skin has resulted from the application, the skin should be wiped dry by means of a soft towel or absorbent cotton to remove any turpentine from the surface, which by its continued contact may cause vesication.

Turpentine is often employed as a rubefacient in the form of the *turpentine stupe*, which is prepared by sprinkling spirits of turpentine over flannel cloths which have been wrung out in hot water, or by dipping hot flannel in warm spirits of turpentine; prepared in either way the stupe should be squeezed as dry as possible to remove the excess of turpentine before being applied to the surface of the body. A turpentine stupe may cause vesication if allowed to remain for too long a time in contact with the skin; its application for from five to ten minutes will usually produce the desired effect; it should be removed after this time, and it can be reapplied if desired.

If the patient complains of severe burning of the skin after the use of turpentine, the painful surface should be freely smeared with vaseline or lard, which will relieve the uncomfortable symptom.

Chloroform.—A few drops of chloroform applied to the surface of the body by means of a piece of lint, muslin, or flannel, and covered by oiled silk or rubber tissue, will excite a rapid rubefacient effect.

Mustard.—Ground mustard or mustard flour prepared from either *Sinapis alba* or *Sinapis nigra* is one of the most commonly used substances to produre rubefacient action. It is generally employed in the form of the *mustard plaster* or *sinapism*, which is prepared by mixing equal parts of mustard flour with wheat flour or flaxseed meal, and adding to this enough warm water to make a thick paste; this is

spread upon a piece of old muslin, and the surface of the paste should be covered with some thin material, such as gauze, to prevent the paste from adhering to the skin. In making a mustard plaster for application to the tender skin of a child, 1 part of mustard flour should be mixed with 3 parts of wheat flour or flaxseed meal.

A mustard plaster or sinapism may be allowed to remain in contact with the skin for a period varying from fifteen to thirty minutes, the time being governed by the sensations of the patient; if it is allowed to remain longer it may cause vesication, which is to be avoided, as ulcers produced by mustard are very painful and extremely slow in healing. After removing a sinapism the irritated surface of the skin should be dressed with a piece of muslin or lint spread with vaseline, boric acid or oxide of zinc ointment.

To excite a rapid revulsive action the *mustard foot-bath* is often employed; it is prepared by adding two tablespoonfuls of mustard flour to a bucket or foot-tub of water at a temperature of 100° to 110°; in this the patient is allowed to soak his feet for a few minutes.

Mustard Papers—Charta Sinapis, which can be obtained in the shops ready for use, are a convenient means of obtaining the rubefacient action of mustard. They are dipped in warm water, and as they are generally very strong, it is well to place a layer of muslin between the surface of the plaster and the skin before applying it to the surface.

Capsicum or *Cayenne pepper* is also sometimes employed as a rubefacient, but it is generally employed in combination with spices, forming the well-known *spice plaster*; this is prepared by taking equal parts of ground ginger, cloves, cinnamon, and allspice, and adding to them one-fourth part of Cayenne pepper; these are thoroughly mixed, enclosed in a flannel bag, and evenly distributed; a few stitches should be passed through the bag at different points, to prevent the powder from shifting its position; before applying it, one side of the bag should be wet with warm whiskey or alcohol. Capsine plasters are also employed to obtain the rubefacient effect of Cayenne pepper.

Aqua ammonia may also be employed for its rubefacient

action. A piece of lint saturated with the stronger water of ammonia, placed upon the skin and covered with waxed paper, and allowed to remain for one or two minutes, will produce a marked rubefacient effect.

Caution should be exercised in applying counter-irritants to patients who are comatose or under the influence of a narcotic, for here the sensations of a patient cannot be used as a guide to their removal, and their too long continued application when the vitality of the tissues is impaired may result in serious consequences.

Vesicants.

Where it is desirable to make a more permanent counter-irritant effect than that produced by rubefacients, substances are employed which by their action on the skin cause an effusion of serum, or of serum and lymph, beneath the cuticle, thus giving rise to vesicles or blisters; they are known as vesicants.

The substance most commonly employed to produce vesication is *Cantharis*, or Spanish fly, and the preparation commonly used is the *Ceratum cantharidis*, which is spread upon adhesive plaster, leaving a margin one-half an inch in width uncovered, which will adhere to the skin and hold the blister in position. The time required for a fly blister to produce vesication is from four to six hours; it should then be removed and the surface should be covered with a flax-seed-meal poultice, or with a warm water dressing. When the blister or vesicle is well developed, it may be punctured at its most dependent part, to allow the serum to escape, and it should be dressed with vaseline or boric ointment. If, for any reason it is desired to keep up continued irritation, after allowing the serum to escape, the cuticle should be cut away and the raw surface should be dressed with some stimulating material, such as the compound resin cerate.

Cantharidal Collodion may also be employed to produce vesication; it is applied by painting several layers upon the skin with a brush over the part on which the blister is to be produced. It is a convenient preparation to use when the

patient would disturb the ordinary blister, as in the case of an insane patient, or where the surface is so irregular that the ordinary blister cannot be well applied. The after-treatment of blisters produced by cantharidal collodion is similar to that previously described.

In the treatment of chronic inflammation it is often better to apply a number of small blisters at intervals than one large blister producing an extensive vesication of the surface. Caution should be observed in using blisters upon the tender skins of children; if employed, they should be allowed to remain in contact with the skin for a short time only. They are contra-indicated in patients in whom the vitality of the tissues is depressed by adynamic diseases, and in aged persons.

A complication which sometimes occurs from the use of cantharidal preparations as blisters is *strangury*, which is shown by frequent and painful micturition, the urine often containing blood. This accident should be treated by the use of opium and belladonna by suppository, demulcent drinks, and warm sitz-baths, and by leeches to the perineum if the symptoms are very severe.

To avoid the development of strangury small blisters should be employed, and should not be allowed to remain too long in contact with the surface, and cantharidal preparations should not be employed in cases where renal or vesical irritation has existed or is present. It is said that strangury may also be avoided by incorporating opium and camphor with the cantharidal cerate.

Aqua ammonia fortior and *chloroform* may be employed to produce rapid vesication, a few drops being placed upon the surface of the body and covered by an inverted watch-glass for a few minutes, or lint saturated with aqua ammonia or chloroform may be placed upon the skin and covered with waxed paper or oiled silk. Either of these agents applied in this manner, and allowed to remain in contact with the skin for fifteen minutes, will produce marked vesication. The blisters resulting from these agents are painful, and they are only used where a rapid result is desired.

Nitrate of silver, in a strong solution or in the form of

the solid stick, may be applied to the surface of the skin to produce a counter-irritant effect. Nitrate of silver, applied by drawing the moist stick across the skin of the scrotum at a number of points, was formerly a popular form of treatment for acute epididymitis.

Acupuncture.

Counter-irritation is effected by this method by thrusting steel needles deeply into the subcutaneous tissues. The needles employed should be of steel, from two to four inches in length, strong, highly polished, and sharp-pointed, and should have round metallic heads or be fixed in handles. (Fig. 95.) Before being used they should be allowed to remain for a few minutes in boiling water or in a carbolized solution to thoroughly sterilize them. In performing the operation of acupuncture, localities containing important organs, large bloodvessels, the joints, and viscera, should be avoided. When introduced the needles should be passed through the skin with a rotary motion, the skin being rendered tense between the thumb and fingers, and pushed into the deep-seated structures. They are allowed to remain in position for a few moments and are then withdrawn, the skin being supported by the thumb and fingers.

Fig. 95.

Acupuncture needles.

Acupuncture has been found of service in cases of deep-seated neuralgia, obstinate rheumatic affections, and sciatica.

Issues.

Issues are ulcers made intentionally by the application of caustics, the moxa, or the knife. They are not often em-

ployed at the present time, but were formerly a popular means of causing long-continued counter-irritation. In making an issue, a region was selected where the subcutaneous cellular tissue was abundant, and which was free from large bloodvessels and nerves, and not near the joints. The plan usually adopted was to apply over the surface of the skin a piece of adhesive plaster perforated in the centre. A small piece of caustic potash or Vienna caustic, mixed with water to make it a paste, was placed in the hole in the adhesive plaster, and held in position by a strip of adhesive plaster. In one or two hours the plaster should be removed and the part should be washed with dilute acid to prevent further action of the caustic; a poultice of flaxseed should next be applied, to hasten the separation of the slough. The ulcer remaining after the removal of the slough may be kept from healing by introducing into it a small wooden ball known as an *issue pea*, or a glass bead or pebble held in place by a compress and adhesive strap.

The Moxa was formerly used to make an issue; it consisted of a small mass of some combustible material, such as punk, cotton, or lint, rolled into pyramidal shape, which was placed upon the surface of the body and ignited so as to produce an eschar upon the skin. To facilitate the application of the moxa an instrument called the porte-moxa (Fig. 96) is employed. The treatment of the eschar resulting from the moxa is the same as that resulting from the use of caustic potash.

Fig. 96.

Porte-moxa.

The knife was also employed to establish an issue, a crucial incision being made through the skin and cellular tissues into the deep tissues; the objection to the use of the knife in forming an issue was the difficulty in preventing the wound from healing.

The Seton.—A seton is a subcutaneous sinus, or an issue with two openings upon the surface, which is prevented from healing by the introduction of a foreign body. It is established by introducing a few strands of silk, a narrow strip of linen, or a rubber

ligature, by means of a seton-needle (Fig. 97), or by means of a sharp-pointed bistoury and an eyed probe. The seton-

Fig. 97.



Seton-needle.

needle should be passed deeply into the superficial fascia, care being taken to avoid important veins and nerves.

A seton may also be established by pinching up a fold of skin and transfixing its base with a narrow, sharp-pointed bistoury (Fig. 98), and passing through the wound thus

Fig. 98.



Method of forming a seton.

made an eyed probe armed with a few strands of silk, a strip of muslin, or an elastic ligature; the probe is then removed and the ends loosely tied together. The wound should be dressed, and at each change of the dressing the strip should be removed, or it may be smeared with some stimulating

ointment, which can thus be brought in contact with the granulating surface by drawing it through the wound.

Actual Cautery.

This method of counter-irritation is accomplished by bringing in contact with the skin some metallic substance brought to a high degree of temperature. This constitutes one of the most powerful means of counter-irritation and revulsion; it is rapid in its action, and is not more painful than some of the slower methods. The cauteries generally employed are made of iron, and are fixed in handles of wood or other non-conducting material, and have their extremities fashioned in a variety of shapes (Fig. 99). The irons are

Fig. 99.



Cautery irons.

heated by placing their extremities in an ordinary fire, or by holding them in the flame of a spirit-lamp until they are heated to the desired point, either white or dull-red heat. They are then applied to the surface of the skin at one point, or drawn over the skin in lines either parallel to or crossing one another. The intense burning which follows the use of the cautery may be allayed by placing upon the cautery-marks compresses wrung out in ice-water or saturated with equal parts of lime-water and sweet oil.

Where the ordinary cautery irons are not at hand, a steel knitting-needle or iron poker heated in the flame of a spirit-

lamp or in a fire may be employed with equally satisfactory results. Where the cautery iron is held in contact with the surface for some time to make a deep burn, the pain of its application may be allayed by placing a mixture of salt and cracked ice upon the spot to be cauterized for a few minutes immediately before its application. The cautery iron should not be placed over the skin covering salient parts of the skeleton or over important organs.

Actual cautery thus applied, in addition to its use in producing counter-irritation and revulsion, is often employed to control hemorrhage and to destroy morbid growths.

Paquelin's Thermo-cautery.

A very convenient and efficient means of using thermo-cautery is the apparatus of Paquelin, which utilizes the property of heated platinum-sponge to become incandescent when

Fig. 100.

Paquelin's cautery.

exposed to the action of the vapor of benzole or rhigolene. (Fig. 100.) The cautery is prepared for use by attaching the gum tube to the receiver containing benzole and heating the platinum knife or button, which is also attached to the benzole receiver by a rubber tube, in the flame of the alcohol lamp for a few moments, and then passing the vapor of benzole through the platinum-sponge, which is enclosed in the knife or button, by compressing the rubber bulb. The points may be brought to a high degree of heat, or may be brought only to a dull-red heat.

This form of cautery may be employed for the same purposes as that previously mentioned; its great advantage consists in the ease with which it can be prepared for use. The knives heated to a dull-red heat will be found of great service in operating upon vascular tumors, where the use of an ordinary knife would be accompanied by profuse or even dangerous hemorrhage. Wounds made by the actual cautery are aseptic wounds, and when dusted with iodoform will generally heal promptly under the scab without suppuration.

Bloodletting.

This procedure is often resorted-to to obtain both the local and the general effects following the withdrawal of blood from the circulation. Local depletion is accomplished by means of some one of the following procedures: *scarification*, *puncturation*, *cupping* and *leeching*, and general depletion is effected by means of *venesection* or *arteriotomy*.

Scarification.

Scarification is performed by making small and not too deep incisions into an inflamed or congested part with a sharp-pointed bistoury; the incisions should be in parallel lines and should be made to correspond to the long axis of the part, and care should be taken in making them to avoid wounding superficial veins and nerves. Incisions thus made relieve tension by allowing blood and serum to escape from

the engorged capillaries of the infiltrated tissue of the part. Warm fomentations applied over the incisions will increase and keep up the flow of blood and serum. Scarification is employed with advantage in inflammatory conditions of the skin and subcutaneous cellular tissue and in acute inflammatory swelling or œdema of the mucous membrane; for instance, of the conjunctiva, and in acute inflammation of the tonsils, tongue, and epiglottis it is an especially valuable procedure. A modification of scarification known as *deep incisions* is practised in urinary infiltration to establish drainage and relieve the tissues of the contained urine and to prevent sloughing; in threatened gangrene and phlegmonous erysipelas the same procedure is adopted to relieve tension by permitting of the escape of blood and serum, and its employment is often followed by most satisfactory results.

Puncturation.

This procedure consists in making punctures, which should not extend deeper than the subcutaneous tissue, into inflamed tissues with the point of a sharp-pointed bistoury; it is an operation similar in character to that just described, its object being to relieve tension and bring about depletion. It is employed in cases similar to those in which scarification is indicated, and is resorted to in cases of diffuse areolar inflammation or erysipelas.

Cupping.

Cupping is a convenient method of employing local depletion by inviting the blood from the deeper parts to the surface of the skin. Cupping is accomplished by the use of *wet* or *dry* cups. When the former are used, no blood is abstracted and the derivative action only is obtained; when wet cups are employed there is an actual abstraction of blood or local depletion as well as the derivative action.

Dry Cupping.

Dry cups as ordinarily applied consist of small cup-shaped glasses, which have a valve and stop-cock at their summit; these are placed upon the skin and an air-pump is attached, and as the air is exhausted in the cup the congested integument is seen to bulge into the cavity of the cup. When the exhaustion is complete the stop-cock is turned and the air-pump is removed, the cup being allowed to remain in position for a few minutes, and is then removed by turning the stop-cock and allowing the air again to enter the cup. This procedure is repeated until a sufficient number of cups have been applied.

FIG. 101.

Cupping-glass and air-pump.

In cases of emergency, when the ordinary cupping-glasses and air-pump cannot be obtained, a very satisfactory substitute may be obtained by taking a wineglass and burning in it a little roll of paper, or a small piece of lint or paper wet with alcohol, and before the flame is extinguished rapidly inverting it upon the skin, or the air may be exhausted by the introduction, for a moment or two, of the flame of a spirit-lamp into the cup. Applied in this manner cups will draw as well as when the more complicated apparatus is used, and when they are removed it is only necessary to press the finger on the skin close to the edge of the cup until air enters the cup, when it will fall off. Although dry cups do not remove blood there is often an escape of blood from the capillaries into the skin and cellular tissue, as evidenced by the ecchymosis which frequently remains at the seat of the cup-marks for some days.

Dry cups, as previously stated, are employed for their derivative action in cases in which depletion is not indicated.

Wet Cupping.

When the abstraction of blood as well as the derivative action is desired wet cups are resorted to, and here it is necessary to have a scarificator as well as the cups and air-pump. (Fig. 102.)

Before applying wet cups the skin should be washed carefully with a carbolic solution, and the scarificator should also be dipped in a carbolic solution. A cup is first applied to produce superficial congestion of the skin; this is removed and the scarificator is applied, and the skin is cut by springing the blades, and the cups are immediately applied and exhausted, and they are kept in place as long as blood continues to flow. When the vacuum is exhausted and blood ceases to flow, they should be removed and emptied, and can be reapplied if it is desirable to remove more blood. A sharp-pointed bistoury which has been sterilized may be employed to make a few incisions into the skin instead of the scarificator, and the improvised cups may be employed if the ordinary cupping apparatus cannot be obtained.

FIG 102.



Scarificator.

After the removal of wet cups the skin should be washed carefully with a bichloride or carbolic solution, and an antiseptic dressing should be placed over the wounds and held in place by a roller bandage.

LEECHING.

In the abstraction of blood by leeching two varieties of leeches are used—the *American* leech, which draws about a teaspoonful of blood, and the *Swedish* leech, which draws three or four teaspoonfuls.

Before applying leeches the skin should be carefully

washed, and the leech should be placed upon the part from which the blood is to be drawn, and confined to this place by inverting a tumbler or glass jar over him; if he does not bite or take hold, a little milk or blood should be smeared upon the surface, which will generally secure the desired result. As soon as the leech has ceased to draw blood he is apt to let go his hold and fall off; if, however, it is desired to remove leeches, they may be made to let go their hold by sprinkling them with a little salt. After the removal of leeches bleeding from the bites may be encouraged if desirable by the application of warm fomentations. Leech-bites should be washed with a bichloride or carbolic solution, and a compress of bichloride or iodoform gauze should be placed over them and secured by a bandage.

It sometimes happens that free bleeding continues from the leech-bite after the removal of the leeches; in this event, if a compress does not control the hemorrhage, the bleeding-point should be touched with a stick of nitrate of silver or with the point of a steel knitting needle heated to a dull-red heat, and if this fails to control the bleeding a delicate harelip pin should be passed through the skin under the bite and a twisted suture should be thrown around this; the wound should then be washed and dressed as previously described.

In applying leeches in or near mucous cavities, care should be taken to see that they do not escape into the cavities and pass out of reach. Leeches should not be employed directly over inflamed tissue, but should be applied to the parts surrounding it; they should not be allowed to take hold directly over a superficial artery, vein, or nerve, and should never be applied to a part where there is delicate skin and a large amount of loose cellular tissue, as in the eyelid or scrotum, as unsightly ecchymoses will result, which will persist for some time. Leeches should not be used a second time.

FIG. 103.



Mechanical leech.

Mechanical Leech.

The mechanical leech is an apparatus which has been constructed to take the place of the leech; it consists of a scarificator, cup, and exhausting syringe or air-pump. (Fig. 103.) In using this apparatus, after the scarificator has been used the piston of the exhausting instrument should be drawn out slowly, which secures a better flow of blood than if a sudden vacuum is made.

The mechanical leech may be employed when the natural leech cannot be obtained, but possesses no advantages over the latter, and is apt to get out of order if not in constant use.

Venesection.

Venesection, as its name implies, consists in the division of a vein, and it is the ordinary operation by which general depletion or bleeding is accomplished. Venesection at the bend of the elbow is the operation which is now usually resorted to for general bloodletting; the vein selected is the median cephalic, which is further from the line of the brachial artery than the median basilic vein.

Fig. 104.

Venesection. (Heath.)

To perform venesection the surgeon requires a bistoury or lancet—the spring lancet was formerly much used, but it is not employed at the present time—several bandages, a small antiseptic dressing, and a basin to receive the blood.

The patient's arm should be carefully cleansed, washed

over with a bichloride solution, and a few turns of a roller bandage should be placed around the middle of the arm, being applied tightly enough to obstruct the venous circulation and make the veins below become prominent, but not to obstruct the arterial circulation. The patient at the same time should be instructed to grasp a stick or a roller bandage and work his finger upon it. The surgeon should next assure himself that there is no abnormal artery beneath the skin, and having selected the vein, the median cephalic by preference, he then steadies the vein with his thumb and passes the point of the bistoury or lancet beneath it and cuts quickly outward, making a free skin opening. The blood usually escapes freely, and the amount withdrawn is regulated by the condition of the pulse and the appearance of the patient. For this reason it is better to have the patient sitting up or semi-reclining when venesection is performed, as the surgeon can judge better as to the constitutional effects of the loss of blood while the patient is in this position.

When a sufficient quantity of blood has been removed, the thumb is placed over the wounded vein and the bandage is removed from the arm above. The wound is next washed with a bichloride solution, and a compress of antiseptic gauze is applied over the wound and held in position by a bandage, which should be so applied as to envelop the limb from the fingers to the axilla. The dressing need not be disturbed for five or six days, at which time the wound is usually found to be healed.

Wounds of the brachial artery have occurred in opening the vein at the bend of the elbow, but if care is taken this accident should not take place.

Venesection may be practised on the *external jugular vein* when, from excess of fat or in the case of children, the veins at the bend of the elbow cannot be easily found. The vein is rendered prominent by placing the thumb or a pad over the vein at the outer edge of the sterno-cleido-mastoid muscle just above the clavicle. The vein is next opened over this muscle by an incision parallel to its fibres. After a sufficient quantity of blood has escaped, the wound is

washed with an antiseptic solution and closed by a compress of antiseptic gauze held in position by a bandage carried around the neck.

Bleeding from this vein has been advocated in cases of apoplexy and cerebral inflammation, but it is questionable whether any advantage is gained by opening the external jugular vein over the vein at the bend of the elbow.

The *internal saphena vein* is also sometimes selected for venesection, and here care should be taken not to wound the accompanying nerve which lies directly behind the vein.

Arteriotomy.

This operation is now scarcely ever performed, but if done the vessel generally selected is the anterior branch of the temporal artery. The position of the vessel is fixed by the finger and thumb, and it is opened by a transverse incision with a bistoury. After a sufficient quantity of blood has escaped the wound is inspected, and if the vessel is not completely divided, its division is completed; the wound should be washed out with an antiseptic solution and a gauze compress should next be applied and held in position by a firmly applied bandage.

Transfusion of Blood.

This operation may be employed to introduce a certain quantity of blood into the circulation of a patient who has suffered from profuse hemorrhage. There are two methods by which transfusion may be effected: the *direct*, by which the blood is conveyed directly and without exposure to the air from the vessel of one person to that of another, and the *indirect*, in which the blood is first drawn from one person and is then injected into the vein of another, being first deprived of its fibrin before being injected.

Direct Transfusion of Blood.

This is best accomplished by using Aveling's apparatus, which consists of a rubber tube, about eighteen inches in length, with a small bulb in the centre, having metallic extremities provided with stop-cocks, and two bevel-pointed metallic canulæ to be used to connect the tube with the bloodvessels. In performing the operation of direct transfusion the bulb and tube are first placed in a shallow basin containing warm normal saline solution (0.7 per cent.), and the bulb and tube are filled with this solution to displace any air which they contain. The person supplying the

FIG. 105.



Apparatus for the direct transfusion of blood.

blood places his arm near the arm of the patient, and the operator exposes a prominent vein on the patient's arm at the bend of the elbow and opens it, and inserts into it one of the canulæ filled with saline solution, with the point directed toward the body, and at the same time an assistant should introduce the other canula into a vein at the bend of the elbow of the party who supplies the blood.

The canulæ are held in position by assistants, and the

tube is quickly connected with them, the stop-cocks being closed before it is taken out of the saline solution, to prevent the entrance of air; then upon opening the stop-cocks a direct communication is established between the circulation of the patient and the donor. (Fig. 105.) The introduction of the contents of the bulb into the vein of the patient is effected by the operator slowly compressing the bulb with one hand, while he keeps the tube closed on the side of the donor with the finger and thumb of the other hand. By relaxing the pressure on the tube on the donor's side of the bulb and closing it on the patient's side, blood will flow from the donor's vein into the bulb as it slowly expands, and when filled the communication with the patient's circulation is again made, and the manipulation is repeated until a sufficient quantity of blood has been introduced as indicated by the condition of the patient's pulse. The quantity of blood or saline solution introduced can be calculated by remembering that at each emptying of the bulb two drachms of fluid are introduced into the circulation. When a sufficient quantity has been introduced the canulæ are removed and the wounds are dressed as ordinary venesection wounds.

Indirect Transfusion of Blood.

Indirect transfusion of blood is accomplished by withdrawing from a vein of the donor by venesection about ten ounces of blood, which is received in a clean glass or porcelain vessel, which is placed in water at a temperature of 110°. The blood thus kept warm is next defibrinated by whipping it with a bundle of broom straws or a wire brush, and after being filtered through a fine linen cloth or wire strainer, it is injected by means of an ordinary syringe attached to a canula which has previously been inserted into a vein of the patient; care should be taken that no air is introduced with the blood. When a sufficient quantity of blood has been introduced, the canula is removed and the wound is dressed in the usual manner. The success of this operation largely depends upon the expedition with which

it is performed; to prevent the coagulation of the blood not more than two minutes should be allowed to intervene between the reception of the blood in the syringe and its introduction into the patient's vein.

Various forms of apparatus have been devised for the operation of indirect transfusion of blood, and of these one

Fig. 106.

Apparatus for the indirect transfusion of blood.

of the best is that devised by Dr. J. G. Allen and modified by the late Dr. C. T. Hunter. (Fig. 106.)

Arterial Transfusion.

This procedure, which consists in injecting defibrinated venous blood into an artery, is occasionally practised. An artery, usually the radial at the wrist or the posterior tibial behind the inner malleolus, is exposed and secured by a ligature; it is then opened on the distal side of the ligature and the point of a canula or the nozzle of a syringe is introduced, directed toward the distal extremity of the limb, and blood, which has been previously defibrinated, is slowly injected. When a sufficient quantity has been introduced the canula is removed, and the division of the artery is completed and its extremities are secured by ligatures, and the wound is closed and dressed.

Auto-transfusion.

This procedure is recommended in cases of excessive hemorrhage to support a moribund patient until other means of resuscitation can be adopted. It consists in the application of rubber bandages or of muslin bandages to the extremities for the purpose of forcing the blood toward the vascular and nervous centres.

Intra-venous Injection of Saline Solution.

It has been proved by experiments and by clinical experience that human blood is not more efficacious in supplying volume to and restoring a rapidly failing circulation than normal salt solution, and as the latter can be obtained with much more ease than blood, its use has largely superseded the latter. The saline solution which is found most satisfactory to employ for this purpose is known as normal saline solution (0.7 per cent.).

The solution should be prepared with water which has been boiled to sterilize it, and should be of a temperature of about 100° when used.

A vein of the patient, at the elbow, should be exposed and should have placed under it, about one-half inch apart, two catgut ligatures; the distal ligature is then tied and an opening is made into the vein between the ligatures; a canula is next inserted into the opening into the vein, and is secured in position by tying the proximal ligature. The canula is first filled with the saline solution, and is then connected with a funnel by means of a rubber tube (Fig. 107), which is filled with saline solution to displace the air, and upon raising the funnel above the part the solution enters the vein; care should be taken to see that the funnel is kept well supplied with solution until a sufficient quantity has been introduced. The quantity introduced is regulated by the condition of the patient's pulse.

Saline solution may also be introduced by means of a syringe when the apparatus described cannot be obtained.

Fig. 107.

Funnel and tube for intra-venous injection.

Intra-venous Injection of Milk.

The intra-venous injection of cow's or goat's milk has also been employed as a substitute for transfusion of blood in patients who have suffered from excessive hemorrhage or from diseases which greatly deteriorate the quality of the blood, such as pernicious anæmia, typhoid fever, and in carbolic acid poisoning. In making one of these injections the milk should be fresh and should be warmed and strained through a fine wire or linen strainer, and it should be introduced by means of a canula inserted into a vein and secured in position by a ligature; to this canula is attached the rubber tubing and funnel, such as is employed in the intra-venous injection of saline solutions. The funnel and tube are next filled with milk prepared as above described, and it is made to enter the vein of the patient by turning the stop-cock and raising the funnel above the patient's body. This injection has been employed in the class of cases mentioned above with apparently beneficial results.

Artificial Respiration.

This procedure is resorted to in cases of threatened death from apnœa consequent upon drowning, profound anæsthetization or the inhalation of irrespirable gases, or any cause which checks or interferes with the function of breathing. Before resorting to artificial respiration care should be taken to see that nothing is present in the mouth or air-passages which will obstruct the entrance of air into the lungs, such as mucus, foreign bodies or liquids, and also that all tight clothing should be removed from the chest which will interfere with the free expansion of the chest walls.

When artificial respiration is resorted to the operator should persevere with it for some time, even when no apparent spontaneous respiratory movements are excited; for resuscitation has been accomplished in seemingly hopeless cases by patient perseverance with the manipulations.

When the first natural respiratory movement is detected the operator should not cease making artificial respiration, but should continue these movements in such a way as to coincide with the spontaneous inspiratory and expiratory movements until the breathing has assumed its regular character.

The temperature of the body should also be restored by frictions to the surface of the body by the hands or by rough towels and hot-water bottles, and warm coverings should be applied for the same object.

Mouth-to-Mouth Inflation.

This method of artificial respiration has been resorted to in cases of great emergency, especially in very young children. The operator draws the tongue forward, closes the nostrils, and applies his mouth directly to the mouth of the patient, and by a deep expiratory effort endeavors to force air into the chest; when this is accomplished the air can be expelled from the lungs by pressure upon the walls of the chest, and the procedure should be repeated about sixteen

times in a minute. The same object may be accomplished by passing a flexible catheter into the trachea through the mouth, or by passing an intubation-tube, to the upper part of which a rubber tube is attached, into the larynx; this can be passed with the fingers without difficulty, and the lungs can then be inflated by the operator blowing into the catheter or tube, or by attaching to it a pair of bellows.

Inflation of the lungs through the nostrils has been employed by Dr. Richardson, of London, who has devised a pocket-bellows for this object. The apparatus consists of two elastic bulbs, to which two rubber tubes are attached, which terminate in a single tube. In using this bellows the terminal tube is introduced into one nostril, the other nostril

FIG. 108.



Richardson's bellows for artificial respiration.

and mouth being closed; air is forced into the lungs by compressing one bulb, and withdrawn by compressing the other bulb.

This bellows may also be attached to a catheter or intubation-tube passed into the larynx, which would prevent the possibility of air escaping into the œsophagus, which is a complication which is liable to occur in mouth-to-mouth inflation or inflation through the nose.

Direct Method of Artificial Respiration (Howard's).

This method of artificial respiration is at the present time considered the most efficacious, and is the one adopted by the United States Life Saving Service, and although the

rules given are for the resuscitation of cases of apparent drowning, the same procedures may be adopted in cases of apnœa arising from other causes.

The rules laid down by Dr. Howard are as follows:

Rule I.—"To expel water from the stomach and lungs, strip the patient to the waist, and if the jaws are clenched separate them and keep them apart by placing between the teeth a cork or a small piece of wood. Place the patient face downward, the pit of the stomach being raised above the level of the mouth by a large roll of clothing placed beneath it. (Fig. 109.) Throw your weight forcibly two or three times upon the patient's back over the roll of

Fig. 109.

First manipulation in Howard's method.

clothing so as to press all fluids in the stomach out of the mouth."

The first rule applies only to cases of drowning, and in using Howard's method in apnœa from other causes it is to be omitted.

Rule II.—"To perform artificial respiration, quickly turn the patient upon his back, placing the roll of clothing beneath it so as to make the breast-bone the highest point

of the body. Kneel beside or astride of the patient's hips. Grasp the front part of the chest on either side of the pit of the stomach, resting the fingers along the spaces between the short ribs. Brace your elbows against your sides, and steadily grasping and pressing forward and upward throw your whole weight upon the chest, gradually increasing the pressure while you count one—two—three. Then suddenly let go with a final push which springs you back to your first position. (Fig. 110.) Rest erect upon your knees while you count one—two; then make pressure again as before, repeating the entire motions at first about four or five times a minute, gradually increasing to about ten or

FIG. 110.

Direct method of artificial respiration.

twelve times. Use the same regularity as in blowing bellows and as seen in the natural breathing which you are imitating. If another person is present let him with one hand, by means of a dry piece of linen, hold the tip of the tongue out of one corner of the mouth, and with the other hand grasp both wrists and pin them to the ground above the patient's head."

This method may be employed in cases of still-birth or in young children, the operator holding the chest of the child in his left hand and compressing it with the right hand.

Sylvester's Method of Artificial Respiration.

In employing this method of artificial respiration the patient should be placed on his back upon a firm flat surface; a cushion of clothing is placed under the shoulders,

FIG. 111.

Sylvester's method—Inspiration. (ESMARCH.)

and the head should be dropped lower than the body by tilting the surface on which he is laid. The mouth being cleared of mucus or foreign substances, the tongue is drawn forward and secured to the chin by a piece of tape tied around it and the lower jaw, or may be pulled out of the mouth and held by an assistant. The operator, standing at the patient's head, grasps the arms at the elbows and carries them first outward and then upward until the hands are brought together above the head; they should be kept in this position for two seconds, after which time they are brought slowly back to the sides of the thorax and pressed against it for two seconds. These movements are repeated

fifteen times in a minute until the breathing is restored, or until it is evident that the case is a hopeless one.

Fig. 112.

Sylvester's method—Expiration. (Esmarch.)

Marshall Hall's Ready Method of Artificial Respiration.

In this method the mouth should first be freed from mucus or foreign bodies, and the patient is turned upon his face with one wrist under his forehead, and a roll of clothing is placed beneath his chest. By turning the body briskly on the side and a little beyond, and then on the face, alternately, respiration is imitated. As the body is brought in the prone position, compression is to be made upon the posterior aspect of the chest. These manipulations should be made fifteen times in a minute.

In using any of these methods of artificial respiration the operator should persevere with them for an hour at least before abandoning the case as a hopeless one.

In cases where the apnœa is due to the presence of a foreign body in the larynx or trachea, it is evident that no efforts at respiration can be successful until the air-passages

are freed from the occluding body, and in such cases tracheotomy should be performed before artificial respiration is attempted; the tracheal wound should be held open by retractors, which in a case of emergency can be made from bent hairpins, or by a tracheotomy-tube if one be at hand.

Forced Respiration.

This is that method of artificial respiration by which air is forcibly passed into the lungs. This procedure is strongly advocated by Dr. George E. Fell, who has devised an apparatus by which it may be satisfactorily accomplished. Prof. H. C. Wood has also made use of forced respiration in the resuscitation of animals with an apparatus somewhat similar to that devised by Dr. Fell with good results, but has never applied it practically in the case of the human subject. Wood's apparatus consists of a pair of bellows, a few feet of rubber tubing and a face mask of rubber, and one or two intubation-tubes; the mask or intubation-tube is attached to one end of the rubber tube and the bellows to the other end of the tube. The mask is applied over the mouth, or if this is not used the intubation-tube is introduced into the larynx, and air is forced into the lungs by working the bellows. He also advises that in the tubing a double metal tube be introduced, with openings so placed that their size can be so regulated by turning the outer tube, that the operator can allow any excess of air thrown by the bellows to escape.

The apparatus of Fell, which he has used in a number of cases with good results, consists of a tracheotomy-tube, a tube connected with the air-control valve, which is attached to an air-warming apparatus, which in turn is connected with a bellows by another tube. In this apparatus air is forced into the lungs and allowed to escape when the lungs have been expanded by the elasticity of the lung tissue and the chest walls.

Forced respiration will prove of value in cases of narcotic poisoning and other accidents in which death is produced by paralysis of the respiratory centres. Dr. Fell has re-

ported a number of cases of narcotic poisoning in which he has used his apparatus with the most satisfactory results.

Aspiration.

This procedure is adopted to remove fluid from a closed cavity without the admission of air, and the instrument which is employed to accomplish this object is known as an aspirator. The two forms of aspirator most generally employed are those of Dieulafoy and Potain. (Fig. 113.)

Fig. 113.

Aspirator.

Potain's aspirator consists of a glass bottle, into the stopper of which is introduced a metallic tube, which is connected with two rubber tubes, one of which is connected with an exhausting pump, and the other with a delicate canula carrying a fine trocar; the apparatus is provided with stop-cocks to prevent the admission of air. In using this aspirator the bottle is exhausted of air by using the air-pump; the canula enclosing the trocar is next pushed through the tissues into the cavity containing the fluid to be removed; the trocar is next removed, and upon opening the stop-cock

LANE LIBRARY. STANFORD UNIVERSITY

the fluid is forced out of the cavity by atmospheric pressure and passes into the bottle or receiver. If the fluid contains masses of lymph or clots which block the canula, interrupting the flow of fluid, a stylet is passed through the canula to free it of the obstruction.

To diminish the pain produced in introducing the trocar and canula, the skin at the point to be punctured may be rendered less sensitive by holding in contact with it for a few minutes a piece of ice wrapped in a towel, or a towel containing broken ice and salt. Care should also be taken to see that the trocar and canula are perfectly clean; to accomplish this it should be carefully washed and placed in boiling water or a 5 per cent. carbolic solution before being used.

In introducing the trocar and canula the operator should be careful to avoid injuring any important veins, arteries, or nerves.

After removing the canula the small puncture should be dressed with a compress of antiseptic or iodoform gauze held in place by a bandage or adhesive straps.

The aspirator is frequently employed in cases of hydrothorax, empyema and ascites, to evacuate the contents of cold abscesses in diseases of the hip and spine, and to remove the contents of a distended bladder until a more radical operation can be performed. It is also a valuable instrument for diagnostic purposes, being frequently used to ascertain the character of the contents of deep-seated tumors containing fluid.

The Stomach-tube.

This consists of a tube about twenty-eight inches in length and three-eighths of an inch in diameter, which is introduced while the patient is in the sitting posture, the head being thrown backward so as to bring the mouth and gullet as nearly as possible in the same line. The tube being warmed and oiled, the surgeon standing in front of the patient passes it directly back to the pharynx, at the same

time introducing the index finger of the left hand to guide its point over the epiglottis; it is then passed gently downward into the stomach. If any obstruction is met with in its passage it should be withdrawn a little way and then pushed gently downward; all manipulations should be made without much force to prevent perforation of the wall of the œsophagus.

The introduction of the stomach-tube may be required for the evacuation of poisons from the stomach, or to wash out the cavity of this viscus, and it may also be used to introduce liquid nourishment into the stomachs of patients who are unable or unwilling to swallow food. In the recently introduced method of treating disorders of the stomach and intestines by washing them out, *lavage*, the introduction of a stomach-tube is required; the tube here employed is from twenty-four to thirty inches in length (Fig. 114), and the fluid is introduced by means of a funnel

Fig. 114.



The Stomach-tube.

attached to its free extremity, or it may be attached to a stomach-pump. In introducing liquid nourishment a syringe or funnel is fitted to the exposed end of the tube which has been passed into the stomach; the syringe or funnel having been filled with milk or beef-tea or broth, the contents are injected gently or allowed to run into the stomach.

In cases of poisoning, where it is desirable to withdraw the contents of the stomach and to wash out the organ, a stomach-tube and syringe may be employed; several syringefuls of warm water are first thrown into the stomach and then withdrawn by suction, but in such cases the use of the stomach-pump will be found more satisfactory.

The Stomach-pump.

This consists of a brass syringe, the nozzle of which is connected with two tubes, one at the end, the other at the side. The passage through the nozzle is regulated by a valve controlled by a lever. The nozzle of the pump is attached to the stomach-tube, and the end of the lateral tube is placed in a pan of warm water. By raising the piston and opening the valve, water may be drawn from the basin, and by closing the valve and depressing the piston it is passed through the stomach-tube into the stomach; when a sufficient quantity has been injected in this manner, by reversing the action of the valve the fluid is drawn out of the stomach and discharged through the lateral tube into a basin. This manipulation is continued until the water returns clear and the stomach has been completely washed out. A less complicated instrument will often serve as well as that just described (Fig. 115).

Fig. 115.

Stomach-pump.

Œsophageal Bougie.—This instrument—which may be passed through the œsophagus into the stomach for the purposes of diagnosis, or for the purpose of dilating strictures of the œsophagus—is passed in exactly the same

manner as the stomach-tube, and, as in the case of the latter instrument, it should be introduced without the use of much force, as perforations of the œsophagus have followed the forcible introduction of such bougies.

Vaccination.

This is a minor surgical procedure which every physician is called upon to perform. The surface may be prepared for the reception of the lymph by abrading the surface of the skin at one or two points with a dull lancet, or by making several superficial incisions with a knife, or by scratching the surface of the skin with the ivory-point charged with lymph, in lines with crossing lines, *cross-scratch*, until a little serum exudes. It is not advisable to draw blood, which washes away the lymph, and for this reason I prefer the abraded surface made by the dull knife or the ivory-point.

The lymph used may be the *humanized* or the *bovine*. The *humanized lymph* may be the viscid fluid taken from the vaccine vesicle on the eighth or ninth day, or the dried scab which separates when the wound is about healed; if the latter is used, a small portion of it is rubbed up with water until it forms a mixture of creamy consistence; this is rubbed into the abraded surface or the punctures. In using humanized lymph care should be taken to see that it is procured from a healthy subject.

Bovine lymph or *virus*, which is now most generally employed, is taken from the vaccine vesicles upon the udders and teats of heifers; ivory-points or quills are dipped into this lymph and allowed to dry, and in using them they are dipped in water for a moment, to moisten the lymph, before being applied to the abraded surface. The ivory-point is one of the most convenient means of vaccinating, as the surface may be abraded with it before the lymph is applied.

It has recently been advised that antiseptic precautions be exercised in performing vaccination, and although all of the details cannot be carried out, I have found that the

exercise of care as regards cleanliness of the surface has been followed by much fewer inflammatory complications in vaccination wounds. In an institution in which I vaccinate yearly a large number of cases, since I have adopted the following precautions I have had fewer bad arms.

The surface to be abraded, usually the left arm below the deltoid, is first washed with soap and water, and then with a 1 : 2000 bichloride solution. Two points of this surface, an inch apart, are then abraded by using a knife which has been washed or dipped in boiling water, or by using the ivory-point which has been dipped in water which has been boiled and cooled down. When the surface has been prepared in the manner described, the moistened virus is rubbed upon it and allowed to dry. Vaccination upon the leg, which is practised by some physicians to prevent the scar from showing, I think is not to be recommended, and I never practise it in this situation, as it is more difficult to keep this part at rest, and I have seen some very severe cases of cellulitis and phlebitis follow leg vaccination.

Hypodermic Injections.

The syringe used to make hypodermic injections is provided with a perforated needle, which is passed into the cellular tissue. (Fig. 116.) Care should be taken to see that the instrument and needle are perfectly clean before being used; if a metallic syringe is employed it should be rendered aseptic by soaking it for a few minutes in boiling water, or in a five per cent. carbolic solution. Hypodermic injections are generally made into parts in which the cellular tissue is abundant, and great care should be observed to avoid introducing the needle into a large vein or artery, as by neglect of this precaution serious symptoms have resulted, from the drug being thrown rapidly into the circulation instead of being slowly absorbed from the subcutaneous cellular tissue; the injury of superficial nerves should also be avoided. Care should also be taken to see that the solutions employed are sterilized if possible, and freshly made solutions should be preferred.

An unclean syringe or a solution which has not been sterilized may give rise to a troublesome abscess at the site of the injection.

FIG. 116.



Hypodermic syringe and needles.

To avoid using solutions for hypodermic use which undergo change from being kept, it will be found convenient to use the compressed pellets which are prepared by the manufacturing chemists, the alkaloids being compressed with a little sulphate of sodium, which increases their solubility, the solution being prepared with boiled water just before being used.

The portions of the body usually selected for hypodermic injection are the outer surface of the thighs or arms and

FIG. 117.



Method of giving a hypodermic injection.

the anterior surface of the forearm. In making a hypodermic injection the syringe is charged and the needle is fastened to the nozzle of the syringe; the skin is next pinched up and the needle is quickly thrust through this

into the cellular tissue; the syringe is then emptied by pressing down the piston, and when the cylinder is empty the needle is withdrawn; the small puncture in the skin resulting seldom bleeds and usually heals without difficulty. (Fig. 117.) In patients who have suffered from profuse hemorrhage, where transfusion of blood is not considered advisable, large injections of normal salt solution may be introduced into the cellular tissue by means of hypodermic injections, or the needle may be introduced into the cellular tissue and connected by a piece of rubber tubing, with an irrigator containing normal salt solution held above the part, and the solution gradually finds its way into the subcutaneous cellular tissue. A large quantity of fluid may be introduced in this way.

Exploring Needle.

This consists of a fine-grooved needle fitted into a handle (Fig. 118), which is introduced into tumors or swellings to

Fig. 118.



Exploring needle.

ascertain the nature of their contents, and its use is often of service for purposes of diagnosis. The exploring trocar (Fig. 119) is employed for the same purpose, or the needle

Fig. 119.

Exploring trocar.

of the hypodermic syringe or a fine needle attached to an aspirator may be used for a like purpose. When either the exploring needle or trocar is employed care should be taken

to see that it is rendered perfectly aseptic before being used; otherwise its employment is not without danger, for I have seen the introduction of an exploring needle into an effusion in a joint for diagnostic purposes followed by suppuration and destruction of the joint, which subsequently necessitated its excision.

SKIN-GRAFTING.

This is a minor surgical procedure which may be employed to hasten cicatrization where large granulating surfaces are exposed, such as result from extensive operations and from burns.

The operation consists in applying shavings of the epidermis or of the epidermis and cutis together, to the granulating surface and holding them in contact with it for a few days; the grafts often seem to disappear, but at the end of a few days, if the part is closely inspected, bluish-white points will be seen to occupy the positions at which the grafts were applied, which become converted into isolated cicatrices from which the healing process rapidly extends. To have a successful result following the use of skin-grafts the surface of the ulcer should be healthy, and the grafts should be applied at a number of points.

FIG. 120.



Scissors for skin-grafting.

The skin is removed by scissors made for this purpose (Fig. 120), or by raising the epidermis with a needle or with forceps and cutting out a small portion of it with a sharp

scalpel. The graft is next applied to the granulating surface with its raw surface in contact with the granulations; after a sufficient number of grafts have been applied, a piece of sterilized protective is laid over them and held in place by means of a few strips of isinglass plaster. An ordinary antiseptic gauze dressing is next applied, and the dressing is not disturbed for a week or ten days, at which time, if the grafts have taken, isolated cicatrices at the points where the grafts were applied will be found to exist.

Electrolysis.

Electrolysis, or the chemical decomposition induced by electricity, is employed in surgery to destroy morbid products, tumors, or exudations. For this procedure a galvanic or continuous-current battery is required, which is provided with electrodes and needles of suitable shapes. In applying electrolysis to a tumor, for instance, the needle connected with one of the poles of the battery is inserted into the tumor and the other rheophore is applied to the surface of the body, or two fine needles, carefully insulated nearly to their extremities, are connected with both poles of the battery by conducting cords; these are introduced into the tumor and a weak current is allowed to pass, and its strength is gradually increased as the operation advances; the current is passed for fifteen or twenty minutes, and the procedure is repeated at intervals of several days, until some decided change occurs in the tumor.

Electrolysis has been applied with success in the treatment of aneurism inaccessible to other operative procedures, in malignant growths, in nævi, goitres, cysts, hydatids, and is at the present time the most satisfactory method of removing superfluous hairs from those portions of the body in which their presence causes disfigurement.

Galvano-cautery.

Galvano-cautery batteries are constructed with plates of large size, placed closely together, so that the internal

resistance is reduced and a current is quickly obtained which will keep a metallic electrode at a white heat. The advantage in the use of this form of cautery is that the electrode can be introduced into the various cavities of the body while cold and quickly heated to the desired temperature. The electrodes are made of various shapes and sizes, according to the object desired (Fig. 121). Galvano-cautery is applied

FIG. 121.

Electrodes for galvano-cautery.

for the same purposes as the actual cautery, but, as previously stated, its use is more convenient in the various cavities of the body, its action can be more easily localized, and by its use hemorrhage is avoided. It is frequently employed to destroy morbid growths in the nasal passages, the throat, vagina, or uterus, and also may be employed in the treatment of superficial external growths; in using it for the removal of growths from the mucous membrane, its application may be rendered practically painless by previously thoroughly cocainizing the parts.

FARADIZATION.

The application of electricity in this form is often employed in surgical affections; in cases of wasting of the muscles following fractures or sprains, in some forms of club-foot, and in lateral curvature of the spine the judicious use of the faradic current will often be found to be followed by the most satisfactory results. The current is applied in such a manner as to bring about contraction of the affected or wasted muscles, and thus improve their nutrition.

Massage.

Massage consists in a variety of manipulations, such as pinching up the integuments and muscles, and rolling them between the thumb and fingers, in stroking or rubbing the surface with the palm of the hand from the periphery toward the centre, to empty the distended veins and lymphatics; rubbing the parts circularly with the extremities of the fingers and thumb or the palm of the hand, or kneading of the parts is another method of practising massage. Massage may also be practised by tapping the surface of the affected part with more or less force with the tips of the fingers held in a row, or with the ulnar border of the hand or with the palm of the hand. Before applying massage to an affected part, if there be a heavy growth of hair, it should be carefully shaved off; otherwise the manipulation may give the patient pain, and irritation of the hair follicles resulting in abscesses will be apt to occur. The part should also be rubbed over with olive oil, vaseline, or cocoa-butter before and during the manipulations.

Massage is often employed with advantage in the treatment of sprains and strains in their subacute and chronic states, and it will be found of great service in the later treatment of fractures involving the joints or their vicinity, in regaining the motion of the parts as well as in improving the nutrition of the muscles which have become wasted from disuse.

Passive Motion.

This manipulation consists in alternately flexing and extending or rotating the limb to imitate the normal joint movements. The motions should be carefully practised, and in cases of fracture they should not be undertaken until there is quite firm union at the seat of fracture, or if for any reason passive motion is made use of before this time the fragments should be firmly supported while it is being employed. Other forms of massage, such as stroking and kneading, may be employed in conjunction with passive motion in the treat-

ment of the troublesome stiffness of the joints resulting from fractures, dislocations, and sprains; passive motion applied in this manner will often restore the function of a stiff joint more satisfactorily and with less pain to the patient than the forcible manipulations of the joint which are practised under an anæsthetic.

The Clinical Thermometer.

For clinical observations two thermometer scales are in general use, the Centigrade and Fahrenheit; the latter is the one commonly employed in America and England. This scale has a limited range above and below the normal bodily temperature, which is $98\frac{2}{5}°$ Fahrenheit or 36° Centigrade. Thermometers are now made with a convex surface,

Fig. 122.



Clinical thermometer.

which serves to magnify the column of mercury, and thus enables the observer without difficulty to note the position of the index. (Fig. 122.)

The temperature of the body may be taken in the mouth, axilla, vagina, or rectum; the two former positions are those generally employed. When taken in the axilla care should be exercised to see that no clothing is interposed between the skin and the instrument, and when the mouth is used for thermometric observations the patient should be instructed to keep his lips tightly closed and breathe through

Fig. 123.



Surface thermometer.

his nose. The thermometer should be kept in place for from three to five minutes.

Surface thermometers are sometimes employed, the in-

struments for this purpose having bulbs of a discoid shape, or are drawn out in the form of a spiral or coil. (Fig. 123.) In using this form of thermometer to determine the amount of variation of the surface temperature, the temperature of corresponding parts of the body on the opposite side and the general temperature of the body should be taken at the same time.

The Rectal Tube.

The introduction of the rectal tube is best accomplished by placing the patient upon his left side, and the surgeon should introduce his index finger well oiled into the rectum and guide the tube upon this through the anus, and by gentle pressure it is gradually passed into the rectum; if a stricture exists in the rectum within reach of the finger, the latter should be used to guide the tube through the opening in this; if the tube becomes caught in a transverse fold of the mucous membrane, and becomes doubled upon itself, it should be withdrawn and a fresh attempt should be made to pass it; in passing a rectal tube all manipulations should be made with extreme gentleness, as it has been shown that its passage is not without danger, perforations of the intestine having followed its use in some cases. In cases of stricture of the rectum high up, the operator has to depend upon the sense of resistance experienced in passing the tube, and in such cases the manipulations should be most carefully made. When the rectal tube is employed to introduce fluids into the large intestine, the fluids may be introduced by means of a syringe, or by pouring them into a funnel attached to the free end of the tube, or by attaching the tube to a fountain syringe, thus allowing the liquid to pass slowly into the intestine.

The rectal tube is often employed with good results in relieving the intestine of excessive flatus, and in introducing water or oil into the intestine in cases of intestinal obstruction, and in those cases where the obstruction results from intussusception or fecal accumulations its use will often prove most satisfactory.

Rectal Bougies.

These instruments are made of the same material as the English flexible catheter, and are of various sizes. They should first be oiled, and carefully introduced in the same manner as the rectal tube. They are generally employed in cases of stricture of the rectum, and they should be used with great care to avoid perforating the wall of the rectum. A very satisfactory substitute for a rectal bougie is a tallow candle, one end of which is melted or rubbed down to a conical shape.

Enemata.

These may be administered by means of the ordinary syringe, or by means of a gravity or fountain syringe; the precautions which should be observed are to introduce the nozzle of the syringe gently and in the right direction, as perforation of the lower portion of the rectum has taken place from the careless and forcible introduction of the nozzle of the enema-syringe; the fluid should also be injected slowly, as by so doing there is less resistance and less tendency for the patient to pass the fluid before the desired quantity has been introduced.

The enema most commonly employed to empty the lower bowel is made by adding a tablespoonful of sweet oil and two teaspoonfuls of spirits of turpentine to one or two pints of warm water in which a little Castile soap has been dissolved; warm water and sweet oil are also frequently used for the same purpose.

Glycerin Enema.—One or two teaspoonfuls of glycerin injected into the rectum, or a suppository made of glycerin, will often be found an efficient substitute for the larger enemata of water.

Nutritious Enemata.

When it is found necessary to resort to feeding by the rectum, the substances employed should be injected into the

rectum by means of a syringe, and care should be taken to see that the quantity is not too large, and that it is of such a nature as not to cause any irritation of the walls of the rectum, or it will not be retained; two ounces in the case of an adult is generally a sufficient quantity to inject at one time.

Peptonized milk or beef juice, or the yolk of an egg beaten up with milk, are often employed, and any unirritating drugs may be mixed with the enema and administered at the same time.

Anæsthetics.

The substances which are employed at the present time to produce either local or general anæsthesia are ice, cocaine, rhigolene, nitrous oxide, chloroform, and ether.

Local Anæsthesia.

Cold.

Local anæsthesia may be produced by the application of cold, either by a piece of ice or a mixture of ice and salt

Fig. 124.



Application of rhigolene spray.

held in contact with the part for one or two minutes, or by directing a spray of rhigolene upon the surface of the part whose sensibility is to be obtunded. (Fig. 124) This form of local anæsthesia is made use of in minor surgical pro-

cedures, such as aspiration, the opening of abscesses, and the removal of superficial tumors.

Rapid Respiration.

Rapidly repeated deep inspirations kept up for a few minutes will produce insensibility to pain, but sensibility to contact is not obliterated. This form of anæsthesia may be made use of in slight operations, such as the opening of abscesses.

Cocaine.

Local anæsthesia produced by the employment of an aqueous solution of the hydrochlorate of cocaine, in strength from 2 to 12 per cent., is often made use of in minor surgical procedures, where the mucous membrane is to be operated upon or growths removed from it. Analgesia is produced by brushing the surface over with the solution of cocaine, or by applying a compress of absorbent cotton saturated with the solution to the part for a few minutes; in mucous cavities the latter method of application will be found most convenient. In using a solution of cocaine to produce anæsthesia in operations upon the eye a 2 or 4 per cent. solution is dropped into the eye, and the application is repeated until the analgesia is complete.

In applying cocaine to the urethra a 4 to 10 per cent. solution is injected into the urethra, and is allowed to remain for two or three minutes; more than one or two grains should not be injected at one time.

When it is desired to produce local anæsthesia of the skin or deeper tissues the application of the solution of cocaine to the surface is not satisfactory, and it should in such cases be injected hypodermically into the deeper layers of the skin and into the cellular tissue of the parts to be operated upon; to avoid mutiple puncture the needle is not completely withdrawn from the wound, but its direction is changed and the solution is thrown into different portions of the tissues. It is well, in situations where it can be accomplished, to cut off the circulation from the part to be

operated upon by placing around it a rubber strap or tube, which prevents its rapid absorption into the general blood current. It is well not to inject more than one grain of the drug in this way, for fatal results have followed the injection of larger quantities; this is especially the case in using cocaine in the urethra and rectum, and in these situations great caution should be exercised in its use.

Some persons also have an idiosyncrasy for cocaine, and children seem more susceptible to its constitutional effects than adults. I have seen several cases in children in which marked symptoms of cocaine poisoning resulted from the application of a 4 per cent. solution to the nasal mucous membrane.

In minor surgical operations, such as amputations of the finger, circumcision, opening of abscesses, and removal of superficial tumors, cocaine-anæsthesia may be employed with advantage, but its utility is most marked in operations upon the eye and those upon the mucous membranes of the nose, throat, rectum, vagina, and urethra. Applied to the surface of an ulcer for a few minutes, which is to be cauterized, it will render the operation almost painless to the patient.

Nitrous Oxide Gas.

This gas is administered for the purpose of producing anæsthesia, and the apparatus best suited for its administration consists of a cylinder of metal in which the gas is compressed; this is attached to a rubber bag which has a mouthpiece fastened to it; this is provided with a double valve, which prevents the expired air from passing back into the bag. The mouthpiece is adjusted over the mouth, and after removing any false teeth, or foreign bodies, from the mouth, the patient is instructed to take deep, full breaths, and in from one-half to one minute the face becomes congested and dusky, and the breathing becomes stertorous, indicating that the patient is fully under the influence of the gas. The anæsthesia from nitrous oxide cannot be prolonged for more than a few minutes, so that it can only be

employed in operations which take a short time for their performance, such as the extraction of teeth, opening of abscesses, and reduction of dislocations or fractures. In England nitrous oxide is frequently used to produce anæsthesia, and when this result is accomplished the anæsthesia is kept up by the administration of ether by the employment of a special apparatus devised for this purpose. Nitrous oxide gas is most commonly employed in dental surgery to produce anæsthesia for the removal of teeth, but is also occasionally employed in minor surgical operations; but from the fact that the apparatus for its administration is a bulky one, its use is not as convenient as ether or chloroform, and in this country it is not much employed in general surgery.

Ether.

Sulphuric ether is one of the most widely employed substances in surgery to produce anæsthesia; it is probably the safest of all anæsthetics, and for this reason should be given the preference over all others.

A patient should be prepared for the administration of ether by not allowing him to have any solid food for at least six hours before its inhalation; he should be in the recumbent posture, and any garments about the chest or neck should be loosened so that the respiratory movements are not interfered with. The surgeon should also see that any false teeth or foreign bodies which may be present in the mouth are removed before the administration of the drug is begun. As the vapor of ether often causes irritation of the mucus membrane of the lips and nasal passages, it is well to anoint these parts with a little vaseline or cold-cream before administering the ether.

It should also be borne in mind that the vapor of ether is very inflammable, and that it is heavier than the air, so that lights brought near the patient while being etherized should be held at a higher level than the ether can or inhaler.

In administering ether a towel folded into a cone or one of the various ether inhalers may be employed. The best

of these is Allis's inhaler (Fig. 125), which consists of a metallic framework covered with leather, which carries a number of folds of a roller bandage, giving a large surface for the rapid evaporation of the drug.

Fig. 125.

Allis's ether inhaler.

If a towel folded into a cone is used, a few layers of stiff paper interposed between the outer layers of the towel will keep the cone in shape and will prevent the evaporation of the ether from its external surface.

In debilitated patients or those who are weak from the loss of blood the administration of half an ounce to an ounce of whiskey from fifteen to thirty minutes before the anæsthetic is given is often advisable.

Half an ounce of ether is next poured over the inner surface of the towel or inhaler and it is brought near the mouth of the patient, and he is requested to take deep breaths or to blow the ether away, and as soon as he has become accustomed to the irritating qualities of the ether vapor, the cone or inhaler is held firmly over the mouth and nose, and the vapor is administered in as concentrated a form as possible; if the respiration and circulation are good there is no disadvantage in pushing the ether. When the conjunctiva is insensible to the touch of the finger, and the muscular relaxation is complete and the breathing tends to

become stertorous, the stage of complete anæthesia has been reached, and the ether should be withdrawn for a time or should be given only in such quantities as suffice to keep the patient in this condition.

The first effect from the inhalation of ether is to produce acceleration of the pulse and respiration; the mucous membrane of the air-passages is irritated and coughing often occurs; there is also in this stage a disposition to muscular movements, and it is frequently necessary to restrain the patient; the brain is also excited and the patient is apt to cry out. These symptoms call for a continuance of the administration of the ether and not for its withdrawal. Succeeding this stage, if the ether is pushed, profound anæsthesia takes place, as is evidenced by loss of consciousness, relaxation of the muscular system, moistened skin, loss of the special senses, contracted pupils, and slow and deep respiration tending to become stertorous.

Under the name of *first insensibility from ether* there exists early in the course of the administration of ether a primary anæsthesia, which lasts for a minute or so, and which may be taken advantage of to perform such minor surgical operations as the opening of an abscess or the reduction of a dislocation or the drawing of a tooth. The recovery from this condition is usually very prompt, and it is not followed by nausea and the after-effects which attend the prolonged administration of ether. During the administration of ether, particularly in the early stage, the patient may stop breathing, the face at the same time becoming cyanosed; this condition calls for the withdrawal of the ether, and if a deep inspiration does not quickly follow, pressure should be made upon the front of the chest, and when this is relaxed a deep inspiration usually takes place and no further difficulty is experienced.

If the patient has eaten solid food shortly before the etherization, vomiting is apt to occur; when this takes place the ether inhaler should be removed and the head should be turned to one side, or the patient should be rolled upon his side, the mouth being kept open to facilitate the escape of the vomited matters. The breathing also sometimes becomes

obstructed by the accumulation of mucus in the fauces; this should be removed by small sponges securely fastened to sponge-holders.

When the anæsthesia is profound it sometimes happens that the muscular relaxation is so complete that the tongue falls backward and the glottis is closed, the face becomes cyanosed and the pulse frequent and irregular, and death is threatened from asphyxia; in this event the head should be extended and the lower jaw should be pressed forward by the fingers placed beneath the ramus of the submaxillary bone. (Fig. 126.) This manipulation is usually sufficient

FIG. 126.

Pushing the lower jaw forward. (ESMARCH.)

to reëstablish the respiratory movements, but if so fortunate a result does not take place artificial respiration should be practised—Sylvester's or Howard's methods being given the preference—the patient's head being placed upon a lower level than the body, the tongue brought forward, and the fauces being cleared of mucus. The respiratory action should also be stimulated by the use of electricity—one sponge-electrode being placed over the sternum, the other being applied to the epigastrium during an inspiratory effort,

If artificial respiration is not satisfactorily applied in this way, forced respiration applied by means of a mask with tube and bellows attached (Fell's apparatus, see page 176), or an intubation-tube with a rubber tube attached, which is connected with a bellows, may be slipped into the larynx, and air may thus be directly forced into the lungs, or the trachea should be opened. This is especially to be recommended if the asphyxia has resulted from blood or vomited matters having entered the larynx. After opening the trachea and introducing a tracheal canula, a rubber tube and bellows is connected with this and respiratory movements are simulated by forcing air directly into the trachea.

The hypodermic injection of strychnia, atropia, or digitalis is also recommended, and the intravenous injection of ammonia is said to have been followed by good results.

Efforts at resuscitation in these cases should be persevered in for at least half an hour, as apparently hopeless cases have been saved by persistent use of these means.

The person intrusted with the administration of the anæsthetic should watch the patient closely and should not have his attention diverted by the operation; he should carefully watch the pulse, respiration, and the color of the patient's face, and be ready to withdraw the ether upon the development of any symptom of danger, and to meet such symptoms, should they arise, by the use of some of the means previously mentioned.

The administration of ether vapor by the rectum was a few years ago employed in many cases, and although anæsthesia was quickly produced, dangerous symptoms sometimes followed its employment, so that this method of administration has been abandoned.

An anæsthetic should never be given to a woman without the presence of a third person, as in some cases these agents give rise to erotic dreams, and it may be difficult to disabuse the patient's mind of the idea that an assault has been committed unless the evidence of eye-witnesses at the time of the anæsthetization can be brought forward to prove that such was not the case.

Chloroform.

A patient is prepared for the administration of chloroform as in the case of ether, the same precautions being taken as regards the removal of false teeth or foreign bodies from the mouth, and to see that the clothing about the chest and neck does not restrict the circulation or respiratory movements. Chloroform is certainly a much more dangerous anæsthetic than ether, and although it is widely used in the British Islands and upon the Continent, it is not used in this country except in certain districts—as in the southern and southwestern districts of the United States, and here its use is followed by fewer fatalities than in the northern districts, so that it is possible that its use is safer in warm climates. Clinical experience has demonstrated the fact that chloroform can be used in aged and very young subjects and in puerperal patients with comparative safety; it is also to be preferred to ether in patients suffering from emphysema of the lungs, bronchitis, and vascular degeneration of the kidneys. It is also employed instead of ether in operations upon the mouth when the actual cautery is employed, on account of its less inflammable character.

Chloroform is administered by pouring a drachm of the drug upon a folded towel, which is first held a few inches from the mouth and nose and gradually brought nearer, but is not allowed to come in contact with the face, as from its local irritating action it will blister the surface; the anæsthetizer should remember that one of the dangers in the administration of chloroform is the risk of too great concentration of its vapor, so that he should see that a sufficient admixture of atmospheric air takes place. Profound anæsthesia is evidenced by insensibility of the conjunctiva to the touch, by complete muscular relaxation, and by the absence of reflexes; the pupils in chloroform-anæsthesia are usually contracted. Various inhalers have been devised to regulate the amount of chloroform administered and to secure the proper admixture of atmospheric air, and the best of these is probably Mr. Clover's apparatus. (Fig. 127.) This consists of a bag holding 8000 cubic inches of air connected

with a face-piece by a flexible tube. The bag is charged by means of a bellows (Fig. 127, 1) measuring 100 cubic inches; and the air is passed through a box warmed with hot water, into which is introduced at each filling of the

Fig. 127.

Clover's chloroform apparatus. (Erichsen.)

bellows as much chloroform as is required for 1000 cubic inches of air. This is done with a syringe (Fig. 127, 2); the amount of chloroform required is usually from 30 to 40 minims. When the bag is full the tube is removed from the evaporator and the mouthpiece (Fig. 127, 3) is fitted to it. Additional air may be admitted by regulating the size of the opening in the mouthpiece; the patient, however, cannot receive a larger proportion of chloroform than the air in the bag is charged with.

Death from the administration of chloroform results from cardiac syncope or from respiratory arrest, and the dangerous symptoms develop so rapidly that the greatest promptness is required to meet them. The person administering chloroform should be constantly on the watch, and should not for a moment have his attention diverted from the patient; great vigilance is here, if possible, more important than during the administration of ether.

When dangerous symptoms arise they are to be treated by lowering the patient's head, and if respiratory arrest has occurred the same means to bring about respiratory action should be employed as for a similar condition during ether narcosis. Cardiac syncope is treated by the use of electricity, the electrodes with a rapidly interrupted current being swept over the chest; hypodermics of digitalis and strychnine and atropia may be employed to stimulate the heart and respiration, and as in ether narcosis the efforts should not be desisted from for some time, as by the persistent employment of these means apparently hopeless cases have been resuscitated.

The A. C. E. Mixture.

This mixture, which consists of 3 parts of chloroform, 1 part of ether, and 1 part of alcohol, has been employed by some surgeons in the place of ether or chloroform, with the idea that the dangers of chloroform are diminished by the combination of it with ether and alcohol. Clinical experience, however, has not proved this view to be correct, and I see no advantage in the use of this combination over that of ether or chloroform. If administered with as much care as chloroform, its administration is accompanied with the same safety. It should be administered upon a towel in the same manner as chloroform, and the patient should be watched as carefully during its inhalation as during the administration of the latter drug, and complications occurring should be treated in the same manner as those arising during the use of chloroform.

Trusses.

A truss for the palliative treatment of hernia is a mechanical contrivance with one or more pads and a strap; these are held in position by a spring to which they are attached, which holds the pad in contact with the skin over the hernial ring.

They are applied in all cases of reducible hernia, and are used in the treatment of hernia at all ages; in infants and young children the continued use of a properly fitting truss is often followed by a radical cure of the hernia.

Trusses are made with steel or rubber springs and with pads of wood, rubber, celluloid, or horsehair, covered with chamois, and their shape and the pressure which they exert varies with the kind of hernia for which they are applied.

A firm compress applied over the inguinal canal or crural ring, secured in position by a firmly applied spica-of-the-groin bandage, forms a very satisfactory temporary means of controlling either of these varieties of hernia.

A properly fitting truss should be worn without discomfort to the patient—that is, should not make too much pressure upon the skin at the points where the pads are applied—and should absolutely prevent the descent of the hernia. In testing the adequacy of a truss, after application, to prevent the escape of the hernia, the patient should be instructed to separate his legs, bend forward over the back of a chair, and cough or strain deeply; if this does not bring the hernia down, its control of the rupture may be considered satisfactory.

Trusses should be applied after the complete reduction of the hernia, while the patient is in the recumbent posture. When first applied the truss should be worn both during the night and day, and if the skin becomes tender at the points of pressure it should be sponged with alcohol and alum, then dried and dusted with powdered starch or lycopodium. Patients at first sometimes complain of discomfort in wearing a truss, but they soon become accustomed to its presence. After a truss has been worn for some time its use at night while the patient is in bed may be dispensed with, but the patient should not remove it until he is in bed in the recumbent posture, and he should reapply it before he rises in the morning. In children it is better to have the truss worn continuously, and if it is removed for bathing the nurse should be instructed to place her finger over the ring to prevent the descent of the hernia until the truss

is reapplied. In applying trusses to male children care should be taken not to make pressure upon an undescended testicle.

Trusses for Inguinal Hernia.

In measuring a patient for this form of truss the circumference of the body midway between the crest of the ilium and the great trochanter should be taken, and the distance from the symphysis pubis to the anterior superior spinous process of the ilium may also be given, as half of this distance corresponds to the position of the internal abdominal ring. In reducible inguinal hernia the truss-pressure should be exerted upon the inguinal canal and directly backward.

Fig. 128.

Truss for inguinal hernia.

To control this variety of hernia a single-spring truss (Fig. 128) may be employed, or the use of a truss having a double spring with flat pads on each side of the spine attached to the springs, and a smaller pad over the inguinal canal on the unaffected side, with a full pad on the side of

Fig. 129.

Hood's truss.

the hernia, will often be found most satisfactory. This, which is known as a Hood's truss, is one which will be found

a very satisfactory instrument both in inguinal and femoral hernia. (Fig. 129.)

Trusses for Femoral Hernia.

In measuring a patient for this variety of truss, the circumference of the body midway between the crest of the ilium and the great trochanter should be taken; the distance of the saphenous opening from the symphysis pubis, as well as from the anterior iliac spine, should also be taken. In reducible femoral hernia the truss-pressure should be directed backward against the femoral canal, and the pad should be large enough to make pressure upon the adjacent tissues through which the hernia passes, as well as upon the relaxed tissues covering the femoral canal. As in inguinal hernia, either a single or a double spring truss may be employed (Fig. 130).

In applying a truss for femoral hernia, care should be taken to see that the pad does not rest upon the pubis, and thus remove the pressure from the crural ring and adjacent tissues and prevent the proper control of the hernia.

Trusses for Umbilical Hernia.

In measuring a patient for this variety of truss, the circumference of the body over the umbilicus should be taken.

Fig. 130.

Hood's truss for femoral hernia.

Fig. 131.

Truss for umbilical hernia.

In reducible umbilical hernia the truss-pressure should be directed backward, and the pad should bear rather on the tendinous margins of the ring than on the hernial opening. A truss for this variety of hernia should have a flat or

slightly convex pad, which is held in position over the umbilical ring by means of springs having counter-pads on either side of the spine attached to their extremities; these are fastened together by a strap (Fig. 131).

A simple and satisfactory truss for umbilical hernia in infants consists of a penny covered by adhesive plaster, held over the umbilical ring by one or two strips of adhesive plaster about two inches in width, which should be applied so as to cover in about the anterior two-thirds of the body. A penny, or a small, flat compress of linen, will be found much more satisfactory than the conical rubber or cork pad which is often recommended.

Trusses for Irreducible Hernia.

The application of a truss to this variety of hernia secures the hernia from injury and prevents the further protrusion of the hernia; such trusses are secured in the same way as those for reducible hernia, but the pads are made concave or cup-shaped, or may have an air-cushion attached to the pad.

Use of Catheters and Bougies.

Catheters are hollow tubes, made either of metal, India-rubber, or other flexible substances.

Metallic catheters are made of silver, or, if constructed of other metals, they should be plated with silver or nickel, to give them a smooth, bright surface which can easily be kept perfectly clean; and their shape should conform to that of the normal urethra (Fig. 132). The shape of the metallic catheter is sometimes changed to meet certain indications; for instance the metallic catheter for use in cases of enlarged prostate is longer and has a longer curve than the ordinary instrument (Fig. 133). The metallic female catheter is shorter and has a much smaller curve than the instrument used for the male urethra.

Flexible Catheters.—The most commonly used variety of flexible catheter is that known as the English catheter, which is made of linen and shellac, and is provided with a stylet; it can be moulded into any shape desired by dipping

Fig. 132. Fig. 133.



Metallic catheter. Prostatic catheters.

it into hot water, which renders it very flexible, and, after moulding it to the proper curve, this can be fixed by immersing it in cold water, which hardens it again.

The French flexible catheters are made of India-rubber, or a combination of this material with other substances. These instruments are conical toward their extremities, and

terminate in an olive-shaped point; they are provided with one or two smoothly finished eyes near their vesical extremities (Fig. 134).

Fig. 134.



French flexible bougie and catheter.

Another form of flexible catheter, known as the elbow-catheter or Mercier's catheter (Fig. 135), has an angle or

Fig. 135.



Mercier's elbowed catheter.

elbow near its vesical extremity; this is often found a satisfactory instrument to use in cases of enlarged prostate. A

Fig. 136.

Soft rubber catheter.

variety of flexible catheter made of soft India-rubber is also sometimes employed. (Fig. 136.)

Catheters and bougies are made according to a certain scale. The English scale runs from 1 to 12; the American from 1 to 20; and the French from 1 to 30.

Bougies and Sounds.

Bougies are flexible instruments which correspond in size and shape to the English and French catheters, and besides these is the acorn-pointed bougie (Fig. 137) and the filiform bougie, which is made of whalebone or of the same material

Fig. 137.



Bulbous or acorn-pointed bougies.

as the ordinary French bougie and catheter. These instruments are of very small size and can often be passed through strictures which will admit no other form of instrument. (Fig. 138.)

Sounds.—These are solid instruments usually made of steel with a smooth surface and plated with nickel; they correspond in size and have the same curve as the metallic catheter; the handle is flattened to allow the operator to grasp them firmly; they are employed in the treatment of strictures by dilatation. (Fig. 139.) The sound used in dilating strictures of the meatus is straight and is shorter than the sound employed in the treatment of urethral strictures. (Fig. 140.)

Introduction of a Catheter.

The passing of a catheter is a minor surgical procedure which every practitioner is at times called upon to employ, and its passage through a healthy urethra is a matter of little difficulty. For the introduction of a catheter the patient may be in the standing, sitting, or recumbent pos-

ture, and the latter is the best in most cases; he should rest squarely on his back and have the thighs a little flexed and separated.

FIG. 138.

Filiform bougies.

FIG. 139.

Steel sound.

FIG 140.

Sound for dilating meatus.

Before passing a metallic catheter the surgeon should see that it is perfectly clean, and after warming and oiling it he stands upon the left side of the patient and grasps the penis with the left hand, and turns it over the pubis and introduces the beak of the catheter into the meatus, and

gently passes it along the urethra until its point passes beneath the symphysis pubis; at this point the handle is elevated and gently depressed between the thighs, and the beak will pass into the bladder. (Fig. 141.)

FIG. 141.



Introduction of catheter. (VOILLEMIER.)

When the prostatic region is reached difficulty is sometimes experienced in passing the catheter; this may be overcome by introducing the finger into the rectum and guiding the catheter through this, or if the prostate is found much enlarged the catheter should be withdrawn, and a prostatic catheter (Fig. 133) should be substituted for it.

The same manipulation is made use of in passing metallic sounds.

Flexible catheters and bougies are passed by grasping the penis and holding it in such a position that it is at a right angle to the axis of the body, and the catheter or bougie is passed into the meatus and carried through the urethra into the bladder by gently pushing the instrument downward.

In this variety of catheter, which has no curve, the surgeon has no means of guiding the point of the instrument, and if an obstruction is met he should withdraw the instrument slightly and make another attempt; all manipulations should be extremely gentle.

The same manipulations are employed in passing bougies through the urethra.

Passing the Female Catheter.

This should be introduced without exposure of the patient, she being in bed with the thighs slightly flexed and separated from each other. The surgeon introduces the fore-

Fig. 142.

Method of holding female catheter.

finger of the left hand between the nymphæ, bringing it from behind forward until he touches the space between the entrance of the vagina and the orifice of the urethra; the catheter is then introduced with the right hand held as shown in Fig. 142, and guided by the left forefinger is passed through the orifice of the urethra into the bladder.

Tying Male Catheter in the Bladder.

When it is desirable to retain a catheter for some time in the male bladder, it is necessary to secure it to prevent its slipping out. Either a metallic or flexible catheter may be employed, but, as a rule, the flexible instrument is to be preferred; there are several methods of securing it in the bladder.

By one method two narrow strips of tape, or two or three strong silk ligatures are attached to the rings at the end of a metallic catheter, or are securely fastened around the end of the flexible instrument; these are next brought backward, one on each side of the penis, and the skin is drawn forward and a strap of adhesive plaster half an inch in width is passed over the strings or tapes and carried three or four times around the body of the penis just behind the position of the glans penis. If the skin has been brought well forward before the straps have been applied, the ligatures are tightened as it slips back, and the catheter has not too much play (Fig. 143).

Fig. 143.

Tying in catheter. (Bryant.)

Another method consists in fastening a strong silk ligature around the catheter just in advance of the meatus; the two ends are next brought backward and tied in a knot behind the corona glandis; the ends are then carried around behind the corona and tied on one side of the frænum; the foreskin is slipped forward and covers the ligatures.

A simpler method of securing the catheter is to perforate the free end with a needle armed with a double ligature of silk or hemp; the needle being removed, two loops are made of the proper length, and these are passed through the ends of a T-bandage, which is secured around the waist, the tails being brought up on either side of the scrotum and

secured to the body of the bandage passing around the waist.

In the female bladder, when it is desirable to keep the bladder empty, the self-retaining catheter is usually employed, which consists of a catheter with a bulb at its vesical extremity, or an ordinary catheter with silk loops and a T-bandage may be employed in the same manner as in securing a male catheter.

The Endoscope.

This instrument is employed to explore the internal cavities of the body. When used to obtain a view of the

Fig. 144.



Endoscope of Desormeaux.

urethra, it consists of a straight conical metallic tube for the urethra, and for the bladder one which is somewhat curved like a vesical sound; there is also an eye-piece, an illuminating apparatus, and an arrangement of mirrors by which a strong light can be thrown upon whatever touches the end of the tube (Fig. 144). By the use of this instrument the urethra and inner surface of the bladder can be examined by the eye.

A view of the urethra may also be obtained by the use of the *urethroscope*, which consists of a straight, hard rubber tube provided with a rounded obturator which projects somewhat beyond the end of the tube. The instrument is introduced into the urethra until the bladder is reached, when it is slightly withdrawn and the obturator is removed and a strong light is thrown into the tube from a head mirror or from an electric lamp, and as the tube is withdrawn various portions of the urethra are exposed to the view of the surgeon.

Washing out the Bladder.

This procedure may be required in the treatment of cystitis, and it is accomplished by passing a flexible catheter with a large eye into the bladder, or a double catheter may be employed. A syringe, or better a rubber bulb holding about a pint, having a nozzle and stopcock (Fig. 145), is

Fig. 145.

Rubber bag with stopcock, for washing out the bladder.

filled with warm water, or with any medicated solution which is desired, and it is then attached to the free end of the catheter and the contents are gently injected into the bladder; care should be taken that the bladder is not too much distended. When the desired amount of fluid has

been injected, it is allowed to run out of the catheter, and the procedure may be repeated until the solution comes away perfectly clear.

Care should be taken to see that the bladder is perfectly emptied of the solution, and in cases of paralysis of the bladder gentle pressure should be made upon the abdomen over the pubis to accomplish this object. Solutions of boric acid, permanganate of potassium, and weak solutions of carbolic acid and of nitrate of silver are often employed in washing out the bladder in cases of chronic cystitis.

Urethral Injections.

In the treatment of urethral inflammations the injection of medicated solutions is generally made use of, and as these injections are usually made by the patient himself, he should be shown or instructed how to employ them. A rubber syringe having a conical nozzle and holding about two or three drachms is the best instrument to employ for this purpose. (Fig. 146.) The syringe having been filled with the solution, and the patient sitting upon the edge of a hard chair, with the thighs separated, grasps the syringe between the thumb and middle finger of the right hand, the tip of the index finger resting upon the end of the piston, and inserts its conical end from a quarter to half an inch within the meatus, which is held open by the thumb and finger of the left hand, and after its introduction it should be drawn tightly around it, the pressure being made laterally so as to narrow the aperture instead of broadening it, as is the case when the compression is in an antero-posterior direction. After the fluid has been thrown into the urethra in this manner the syringe is removed, and the patient is instructed to hold the lips of the meatus together for one or two minutes to prevent the escape of the fluid.

Fig. 146.

Shape of nozzle of urethral syringe.

Sutures.

A variety of materials are employed for sutures, such as silk, catgut, silver or iron wire, silkworm-gut, and horse-hair; the material most frequently employed at the present time is either catgut, silk, or silkworm-gut, although some surgeons still prefer silver wire. Catgut is practically the only substance employed as a suture which is absorbable; the other varieties of suture require removal after their application, although some sutures, such as the silk, when employed in subcutaneous wounds may be cut short, as they are apt to become encysted and produce no trouble. It matters little what variety of material be employed for suture if the surgeon is careful to see that it is rendered thoroughly sterile before being brought in contact with the wound.

Sutures of Relaxation are those which are entered and brought out at some distance from the edges of the wound, and are employed to prevent dangerous tension upon the sutures which close skin wounds. This form of suture is employed by the use of the quilled, button, or plate suture.

Sutures of Coaptation.—These are superficial sutures applied closely together and include only the skin; they are employed to secure accurate apposition of the cutaneous surface of wounds.

Sutures of Approximation are those which are applied deeply into the tissues to secure approximation of the deep portions of a wound; this object is accomplished by the use of the quilled, button, or plate suture.

Secondary Sutures.—These sutures are applied when the surfaces of the wound are covered by granulations, when the primary sutures have failed to secure apposition of the edges of the wound, or in cases of secondary hemorrhage where the opening of the wound has been necessitated to turn out the blood-clot and secure the bleeding vessel, or in plastic operations where the primary sutures have failed to secure adhesions of the edges of the flaps. They are also employed with advantage in cases in which it is necessary to pack a wound with antiseptic gauze, or to allow hæmostatic forceps

to remain clamped upon bleeding tissues in a wound at the time of operation. The sutures should in such a case be introduced and loosely tied at this time, and when the packing or forceps are removed at the end of two or three days the sutures are tightened so as to secure apposition of the edges of the wound.

Surgical Needles.

Needles for surgical use are of different sizes and shapes (Fig. 147); straight needles are the ones most commonly

Fig. 147.



Surgical needles.

employed, but curved needles will be found most convenient for the introduction of sutures in wounds of certain

Fig. 148.

Mounted needle.

locations. Tubular needles are often employed in introducing sutures in wounds in which the use of an ordinary

needle is difficult: for instance, in the operation for cleft palate, and for the introduction of sutures in deep wounds, a mounted needle will often be found very useful (Fig. 148). Needles should be sharp and clean and should be rendered thoroughly aseptic before being used. A needle-holder is often required for the satisfactory introduction of needles in wounds of certain localities (Fig. 149); if

FIG. 149.

Needle-holder.

this is not at hand the needle may be held by a pair of dressing forceps or a pair of hæmostatic forceps.

METHOD OF SECURING SUTURES AND LIGATURES.

Metallic sutures are usually secured by twisting the ends together or by passing the ends through a perforated shot and clamping the shot with a shot-compressor, which securely fixes them.

Sutures and ligatures of catgut, silk, silkworm-gut or horsehair are secured by tying, and several different knots are employed to secure them.

Reef or Flat Knot.

This is one of the best forms of knot to use in securing sutures or ligatures, and it is made by passing one end of the thread over and around the other end, and the knot thus formed is tightened; the ends of the thread are next carried toward each other and the same end is again carried over and around the other, and when the loop is drawn tight we have formed the reef or flat knot (Fig. 150).

FIG. 150.

Reef or flat knot.

Surgeon's Knot.

This knot is formed by carrying one end of the thread twice around the other end (Fig. 151); and after tighten-

FIG. 151.



Surgeon's knot.

ing this loop the same end is carried over and around the other end as in the case of the final knot of the reef or flat

FIG. 152.



Surgeon's knot and reef knot combined.

knot. The surgeon's knot and reef knot combined is one of the best methods of securing sutures or ligatures of

catgut or silk, as the first knot is not apt to relax before the second knot is applied. (Fig. 152.)

Granny Knot.

This method of tying the ligature or suture should not be employed, as the resulting knot is not as secure as the reef knot and is apt to relax; it differs from the latter in the fact that one end of the thread having been carried

FIG. 153.



Granny knot.

across and around the other end, the knot is completed by carrying the same end under and around the other end of the thread (Fig. 153).

The *Staffordshire* knot, which is much used to secure the pedicle in the removal of the ovaries or ovarian tumors, is applied as follows: A handled-needle armed with a stout silk ligature is passed through the pedicle, and then withdrawn so as to leave a loop on the distal side; this loop is drawn over the ovary or tumor and one of the free ends is passed through it so that one end is above while the other end is below the retracted loop. (Fig. 154.) The ends are then seized and drawn through the pedicle; at the same time the thumb and forefinger are pressed against it, until sufficient constriction is made, and the ends are finally secured by tying as in the securing of an ordinary ligature.

FIG. 154.



Staffordshire knot.

Varieties of Suture.

The Interrupted Suture.

This variety of suture is the one most usually employed in the apposition of wounds, consisting of a number of single stitches, each of which is entirely independent of those on either side. In applying this suture the surgeon holds the edge of the wound with the fingers or forceps and thrusts the needle, previously threaded, through the skin three or four lines from the edges of the wound. He then passes the

Fig. 155.



The interrupted suture.

needle from within outward through the tissues of the opposite flap at the same distance from the edge of the wound. (Fig. 155.) Each stitch is secured as soon as it is passed—by tying if a silk, catgut or silkworm-gut suture be used, or by twisting if a silver-wire suture is employed. A suture may be used with a needle threaded on each end, and in this case both needles are passed from within outward. The sutures may be secured as soon as applied or they may be left unsecured until a sufficient number have been introduced and then they may be secured by tying or twisting. Care should be taken to see that they make no tension on the

edges of the wound and that they are so introduced as to make the best possible apposition of the parts.

In extensive and deep wounds it may be found necesary to introduce both deep and superficial sutures, the former bringing about apposition of the muscles and deep fascia, the superficial layer bringing together the superficial fascia and skin.

The *deep* or *buried* sutures are often employed to unite fascia, muscles or tendons, and the best material for this variety of suture is either catgut or silk.

Continued or Glover's Suture.

This variety of suture is applied in the same manner as the interrupted suture, but the stitches are not cut apart

FIG. 156.



Continued or glover's suture: method of securing.

and tied; it is made with silk or catgut, and is secured by drawing it double through the last stitch and using the free end to make a knot with the double portion attached to the

needle. (Fig. 156.) This suture is generally employed in intestinal sutures, but may also be employed in bringing about apposition of the edges of wounds in tissues of loose structure.

The Twisted or Hare-lip Suture.

This is a very useful form of suture where great accuracy and firmness of apposition of the edges of the wound are desired. It is applied by thrusting pins or needles through both lips of the wound, the edges being kept in contact over the wound by figure-of-eight turns with silk or wire. (Fig. 157.) The ends of the pins should be cut off by pin-cutters after the sutures are applied, or should be protected by pieces of cork or plaster to prevent them from injuring the skin of the patient and causing him pain.

The India-rubber Suture.

This is applied by first passing the pins or needles through the edges of the flaps, and instead of the twisted figure-of-

Fig. 157.

Twisted or hare-lip suture.

Fig. 158.



India-rubber suture.

eight suture of silk, delicate rings of India-rubber are employed. (Fig. 158.)

The twisted or hare-lip suture is frequently employed in plastic operations about the face and in other parts of the body, where accurate apposition of the flaps is desired.

The Quilt Suture.

This variety of suture is made with silk or catgut, and is employed in wounds to effect very close approximation of

the parts and to prevent bagging; it is often employed in connection with the continued suture, and is applied as shown in Fig. 159.

Fig. 159.



The quilt suture.

The Quilled Suture.

In making use of this suture a needle armed with a double thread of wire or silk is passed through the tissues as in applying the interrupted suture, but at a greater distance from the edges of the wound. Into the loops on one side of the wound is inserted a quill or piece of a flexible catheter or bougie, and on the opposite side the free ends of the sutures are tied around a similar object after being tightened. (Fig. 160.) This form of suture makes deep and equable pressure along the whole line of the wound. In applying this suture it may be found well in some cases to introduce a few superficial interrupted sutures along the line of the wound to secure accurate approximation of the skin.

This form of suture was formerly much employed in cases of deep wounds to secure accurate apposition of the deep

FIG. 160.



The quilled suture. (SMITH.)

portions of the wound, but recently the introduction of buried catgut sutures has supplanted the use of this variety of suture.

Button or Plate Suture.

This suture is applied by passing a needle armed with a double thread as in the case of the quilled suture, the ends of the suture being passed through the eyes of a button or through perforations in a lead plate before being threaded in the eye of the needle. After the suture prepared in this way has been passed through both sides of the wound, the needle is removed and the free ends of the suture are passed through the eyes of a button or the perforations in a lead plate on the opposite side of the wound, and are tightened and secured. (Fig. 161.) This form of suture may be employed in deep wounds to accomplish the same purpose as the quilled suture, and allows the cutaneous margins of the wound to remain free from compression, and here, as in the case of the quilled suture, a few interrupted

FIG. 161.



Button suture. (SMITH.)

sutures may be introduced between the button or plate sutures to secure accurate apposition of the skin surfaces if desired.

Tongue-and-groove Suture.

This variety of suture, devised by the late Dr. Joseph Pancoast, consists in slipping the margin of the flap which has been bevelled into a groove, made by dissecting up the margin of the skin surrounding the raw surface which is to be covered. In applying this suture the wire or thread used has a needle applied on each end, and after passing the sutures so as to secure the flaps the free ends are secured over a pad of adhesive plaster or a disc of lead or a button. (Fig. 162.)

Fig. 162.



Tongue-and-groove suture.

Shotted Sutures.

This suture receives its name not from any special method of application, but solely from the way in which it is secured; any of the previously mentioned varieties of sutures may be employed. The material used in applying this suture may be catgut, silver wire, silkworm-gut, silk, or horsehair, and after the suture has been passed the needle is removed, and the ends are passed through a perforated shot; the ends are then drawn upon to bring the edges of the wound in contact, and the shot is pressed down to the skin and clamped by means of a shot-compressor. The suture is then cut off flush with the surface of the shot.

This method of securing sutures is especially useful in closing wounds in the mucous cavities, such as the vagina, rectum, and mouth, where the knot or twist of the wire might cause irritation of the surface or pain to the patient; it is also a useful method of securing sutures in plastic operations; it also facilitates the removal of the sutures, as the shot is not apt to be obscured by the swollen

tissues and is easily seized by forceps when the loop is divided.

Removal of Sutures.

Where sutures are buried in the tissues or used to approximate parts in cavities which are subsequently closed, such material should be used for sutures as will be absorbed in a few days, or will become encysted and remain harmless in the tissues—such as catgut, silkworm-gut, or silk—and it is needless to state that sutures used with this end in view should be rendered perfectly aseptic before being employed.

Catgut sutures, when well prepared and used for sutures in external wounds, usually undergo absorption in from ten to fifteen days; the loop buried in the tissues is absorbed and the knot may be removed from the surface with forceps or comes off with the dressings.

The other substances, such as silk, silkworm-gut, silver wire, and horsehair, are removed by cutting one side of the loop and making traction upon the knot of the suture with forceps, or in the case of the wire suture, after dividing the loop and straightening out one end of it, the wire should be withdrawn in a curved direction.

Sutures which are not causing any irritation should be allowed to remain in position until the wound is solidly healed. The time usually required for their retention in cases of aseptic wounds is from eight to twelve days.

Lembert's Suture.

Lembert's suture is used in wounds of the viscera covered by the peritoneum, with the object of bringing in contact the peritoneal surfaces. This form of suture is usually employed in closing wounds of the intestine or stomach. (Fig. 163.)

A needle armed with a fine catgut or silk thread is passed, and it is better to employ a round needle, such as the ordinary sewing-needle, in preference to the bayonet-pointed needle, as there results by its use less bleeding from the

punctures. The needle is first carried through the peritoneal and muscular coats of the intestine a short distance from the wound, and it is then carried across the wound

FIG. 163.



Lembert's suture. (BRYANT.)

FIG. 164.



Lembert's suture. *a*, serous; *b*, muscular; and *c*, mucous coat. (SMITH.)

and passed through the same portions of the intestine a short distance from the edge of the wound on the opposite side, and when the suture is tightened the peritoneal surfaces of the intestine are inverted and brought into contact with each other (Fig. 164); the interrupted or continued suture may be employed in making this form of suture.

Gély's Suture.

In applying this form of suture in intestinal wounds a ligature armed with a fine needle at each end is employed,



out at the lower angle of the external wound; a thread is also tied under the head of each pin and brought out at the upper angle of the wound, and at the end of three or four days the pins are removed by means of the threads attached to them, and at the same time the sutures, having been freed by the removal of the pins, are withdrawn.

Czerny Suture.

This suture is applied in intestinal wounds by passing the needle armed with a catgut or silk thread through the serous membrane on one side of the wound of the intestine and out at the wound surface so as not to include the mucous membrane; the needle is then passed through the wound surface on the opposite side, avoiding the mucous membrane, and brought out through the serous membrane a short distance from the edge of the wound. By this suture the lips of the wound are approximated. For additional security in preventing the escape of the contents of the intestine and to secure approximation of the serous surfaces a few Lembert sutures should be introduced.

Jobert's Suture.

This suture which was employed in transverse wounds of the intestine which completely or incompletely divided the

FIG. 167.



Jobert's suture.

gut, is introduced after turning the lower end of the bowel in upon itself. When the division of the gut was incom-

plete he employed only one suture, when complete two; the ends of the sutures were brought out of the external wound. (Fig. 167). By this method of suture the two serous surfaces are brought into contact.

Sutures Employed in Intestinal Anastomosis.

When it is desired to form a permanent orifice between two portions of the gut, the ends of the gut are closed and an opening is made in each portion of the gut and the walls

Fig. 168.



Method of applying Senn's decalcified bone plates. (Greig Smith.)

of the gut surrounding the openings are held in contact with each other by sutures attached to perforated plates of decalcified bone; this is the method devised by Senn. The manner of using the bone plates and sutures is shown in Figs. 168 and 169. To accomplish the same purpose

rubber rings or perforated plates of rubber have been employed, also rings made from catgut, to which the sutures are attached, are applied in the same manner as Senn's plates. In using the rubber rings or plates it is well to divide them at one or two points and unite them by catgut sutures which will soften and be dissolved in a few days and allow the ring or plate to change its shape and facilitate its passage through the bowels; if catgut rings are employed these will be softened and dissolved in a short time so as to be passed without difficulty. Intestinal anastomosis may be

FIG. 169.



Diagram showing position of bone plates in intestinal anastomosis after resection of the bowel. (ROBERTS.)

employed instead of Jobert's suture or the circular suture in wounds completely dividing the intestine and after resection of the gut for the removal of growths or for stricture.

Sutures Employed in Gastrostomy.

In this operation, when the wall of the stomach has been exposed, two hare-lip pins should be thrust through the integument and tissues near the edge of the wound and then through the peritoneal and muscular coats of the stomach, to bring the surface of the stomach in contact with the peritoneum covering the inner surface of the abdominal walls in the region of the wound; a few sutures of silk may also be

introduced to secure the wall of the stomach to the edges of the wound. The opening of the stomach is postponed for four or five days if possible, until the adhesion between its walls and the abdominal parietes is secure, and at this time the sutures and the pins are removed.

When immediate opening of the stomach is required for any reason, after the wall of the stomach has been exposed two silver wire sutures are passed through the peritoneal and muscular coats of the stomach by means of a needle;

FIG. 170.



Sutures for immediate gastrostomy. (ROBERTS.)

these sutures should be placed transversely to the external abdominal wound and serve to draw the wall of the stomach in contact with the inner margins of the abdominal incision.

A long silk suture is next passed through the outer coats of the stomach so that the loops project upon the external surface of the organ (*a*). A needle, having a hook near its extremity (*c*), is passed through the abdominal wall and engages the loop and draws it to the surface of the abdomen

near the edge of the abdominal wound; the same manipulation is repeated until all of the loops have been brought to the surface. (Fig. 170.)

A piece of rubber tube is carried around the external wound and slipped through the loops which project upon the surface of the abdomen (*c*), and by drawing the loops tight over the rubber tube and tying the ends of the suture the stomach wall is secured in contact with the inner margins of the abdominal wound, and it may be safely opened after being thus fixed.

In the operation of *gastrotomy*, where the stomach has been exposed and opened and the foreign body removed, or its cavity has been explored, or its orifices dilated as the case may be, the wound in the wall of the stomach is closed with Lembert's sutures, silk or catgut being the material employed for sutures. The abdominal wound is next closed with deep sutures which include the parietal peritoneum.

Ligatures Used in the Treatment of Vascular Growths.

Various forms of ligature are used for the strangulation of vascular growths; the material used for ligatures is usually strong silk or hemp thread, catgut, or silver wire.

The Single Ligature with Pin.

This is applied by first inserting a hare-lip pin through the skin near the edge of the growth, passing it under the growth and bringing its point out through the skin at a point opposite the point of entry; a strong silk or hemp ligature, which has been well waxed, is passed under the ends of the pin surrounding the base of the tumor and is drawn tight enough to strangulate the growth, and is secured by two knots (Fig. 171). If the growth is of considerable size it is better before applying this ligature to introduce a second pin at right angles to the first one, and then secure the ligature under the pins. In applying these forms of

ligatures to healthy skin, the patient is saved much pain, and the separation of the mass is hastened, by cutting a groove in the skin with a sharp knife at the point where

FIG. 171.



Vascular tumor strangulated with pin and ligature. (ROBERTS.)

the ligature is to be applied; the ligature when tied is buried in the groove thus made.

Double Ligature in Vascular Growths.

This ligature is applied by passing a needle or a needle with a handle, armed with a double ligature, through the skin near the growth, and then passing it under the tumor and bringing it out through the skin at a point directly

FIG. 172.



Method of applying double ligature. (ROBERTS.)

opposite the point of insertion; the ligature is then divided and the needle removed and the tumor is strangulated by tying firmly the corresponding ends of the ligature on each side of the tumor, each ligature strangulating one-half of the growth (Fig. 172).

The double ligature may also be applied by first passing a pin under the growth and then passing a needle armed

with a double thread under the tumor at right angles to the pin, and after removing the needle the ends of the liga-

FIG. 173.



Method of applying double ligature and pin. (BRYANT.)

ture are tied and the tumor is strangulated in two sections (Fig. 173).

Quadruple Ligature.

In applying this ligature two needles carrying a double ligature are passed under the growth at right angles to each other, or if the handled needles be used they may be first passed in this manner, and then threaded with double ligatures, which are carried under the growth as they are withdrawn. The needles being removed, the surgeon ties two ends of the ligature together and repeats this procedure until the growth has been strangulated in four sections. (Fig. 174.)

Subcutaneous Ligature.

This is applied by introducing a needle armed with a ligature through the skin near the growth, and carrying it through the subcutaneous tissues around the growth for a

short distance, then bringing it out through the skin. The needle is again introduced through the same puncture and is again brought out through the skin at some distance from

Fig. 174.

Method of applying quadruple ligature. (Liston.)

the first point of exit, and is next introduced through this puncture and brought out at a more distant point. In this way the growth is completely encircled by a subcutaneous ligature, which finally is brought out at the point of entrance; the tumor is strangulated by firmly tying together the ends of the ligature. (Fig. 175.)

If a needle armed with a double ligature is first passed under the growth the ligature is divided, and by passing each end of the divided ligature subcutaneously around the

growth it may be strangulated subcutaneously in two sections.

FIG 175.



Method of applying subcutaneous ligature. (HOLMES.)

Erichsen's Ligature.

This ligature is employed to strangulate tumors of irregular shape in a number of sections. A strong silk or hemp ligature three yards in length, one-half of which is stained

FIG. 176.



Method of applying Erichsen's ligature. (ERICHSEN.)

black, is carried by a needle as a double ligature under the growth at various points so as to leave a series of loops about nine inches long on each side of the tumor (Fig. 176); the black loops being cut on one side, the white on the

Fig. 177.



Erichsen's ligature applied.

other, the ends are then firmly tied so as to strangulate the growth in sections. (Fig. 177.)

Elastic Ligatures.

Ligatures made of India-rubber varying from half a line to several lines in thickness are often made use of in surgery. They may be employed to strangulate growths such as moles or nævi, or in the treatment of fistulæ, and are especially useful in the treatment of those cases of fistula-in-ano in which the internal opening into the bowel is situated high up, as the division of such fistulæ by this means is accomplished without hemorrhage and with less risk than by the employment of the knife. In applying elastic ligatures in such cases the ligature, after being passed through the fistula by means of a probe, is carried out through the internal opening; the anus is next well stretched, and the elastic ligature is then firmly tied with two or three knots; the greater the tension made before the ligature is tied the more rapidly will it cut its way out. The smaller sizes of rubber drainage-tubes may be substituted for the solid rubber ligatures.

Treatment of Hemorrhage.

The surgeon may be called upon to treat the following varieties of hemorrhage: *arterial*, *venous*, or *capillary;* and these again are classified according to the time of their occurrence, as—*primary*, that is, bleeding which occurs at the time the wound is inflicted; *intermediary* or *consecutive*, that which occurs within twenty-four or forty-eight hours after the reception of the injury, which generally takes place during the period of reaction; and *secondary*, which takes place after forty-eight hours, and may occur at any time subsequent to this period until the wound is healed. The treatment of hemorrhage is either *constitutional* or *local*.

The *constitutional* treatment of hemorrhage consists in keeping the patient in the recumbent posture and avoiding any sudden elevation of the head or arms which might induce fatal syncope. Opium is a valuable remedy and should be freely used. Ergot, gallic acid, acetate of lead and tincture of iron may also be employed, and stimulants and food should be carefully administered, and in extreme cases autotransfusion or transfusion of blood or normal salt solution, as described on page 168, may be resorted to.

In the *local* treatment of hemorrhage various measures may be adopted which may be either temporary or permanent in their action.

Temporary Control of Arterial Hemorrhage.

This may be effected by pressure applied directly to the bleeding vessel in the wound or by pressure applied indirectly to the main artery between the point of its injury and the centre of the circulation, and this pressure may be made by the fingers, *digital compression*, by compresses, or by means of tourniquets.

Digital Compression.

This constitutes one of the most valuable means employed in the temporary control of hemorrhage; the finger is pressed

directly upon the bleeding vessel in the wound or is used to make pressure upon the artery from which the bleeding arises at some point between the wound and the centre of the circulation. (Fig. 178.) Control of hemorrhage by digital presssure can only be maintained for a few minutes,

Fig. 178.

Digital compression of the femoral artery.

for the fingers of the surgeon or assistant soon become tired, so that it is only employed until means are adopted for the permanent control of the bleeding. Digital compression of the radial and ulnar arteries is frequently resorted to for the control of hemorrhage during amputations of the fingers, also of the axillary and femoral arteries in amputations at the shoulder- and hip-joints.

It is also used to control hemorrhage from wounds either the result of accident, or those made by the knife of the surgeon, in which case the finger is placed directly upon the divided vessel, or employed to hold a sponge or compress firmly in the wound.

Compresses.

By the use of compresses placed directly in the wound or applied to the vessel between the wound and the centre of the circulation, the temporary control of hemorrhage may be very satisfactorily accomplished. Where it is possible, the compress which is applied in the wound should be made of

antiseptic gauze, thereby diminishing the chances of wound-infection.

The compress should be held in position by a bandage firmly applied and is generally employed only as a temporary expedient until a more permanent means of controlling the bleeding is adopted.

Tourniquets.

These instruments, which are employed for the temporary control of hemorrhage from wounds, are of many different kinds.

Fig. 179.



Petit's tourniquet.

Petit's tourniquet, which is the best for ordinary use, consists of two metal plates connected by a strong linen or silk strap, with a buckle—the distance between the plates being regulated by a screw. (Fig. 179.) In applying this tourniquet a compress or roller bandage is placed directly over the artery to be compressed and may be held in position by a few turns of the roller bandage. The lower plate

of the tourniquet is placed directly over this pad and the strap is tightly secured around the limb to keep the instrument in place. The screw is then turned so as to separate the plates and tighten the strap, thus forcing the compress or pad upon the artery controlling its circulation. This instrument is very generally employed for the control of hemorrhage in wounds of the extremities and is especially useful in amputation of these parts, being placed over the main artery some distance above the seat of operation.

The Spanish Windlass.

An improvised tourniquet, known as the Spanish windlass, may be employed in cases of emergency; it is prepared by folding a handkerchief or piece of muslin into a cravat and placing a compress or smooth pebble on the body of the cravat; this is placed over the artery to be controlled, and the ends of the handkerchief are tied loosely around the limb; a short stick is passed through this loop, and by twisting the stick the loop is tightened and the compress is forced down upon the artery (Fig. 180).

FIG. 180.

The Spanish windlass.

Many other forms of tourniquet have been devised which have the pad and counter-pad arranged as to make pressure upon the vessel desired, such as Lister's aorta compressor (Fig. 181), which is employed in the treatment of aneurism of the iliac vessels, and for the control of hemorrhage in amputation at the hip-joint. Hoey's clamp (Fig. 182) and Signorini's tourniquet (Fig. 183) are constructed upon the same principle, and are frequently employed to control the

FIG. 181.



Lister's aorta compressor.

circulation in the femoral artery in cases of operations on the thigh and leg, and in the treatment of femoral or popliteal aneurism.

FIG. 182.



Hoey's clamp.

FIG. 183.



Signorini's tourniquet.

The *elastic tube*, or *strap of Esmarch's apparatus* (Fig. 184) may also be employed for the temporary control of arterial hemorrhage, being applied above the wound, and if this is not at hand, any strong rubber cord, or a piece of large-sized drainage-tube may be used as a substitute. In hemorrhage from wounds of the hands and feet, especially in children, and in controlling hemorrhage from wounds of the penis, a piece of drainage-tube, firmly applied above the wound, may be employed with advantage. This tube or

FIG. 184.

Elastic strap of Esmarch's apparatus.

strap, although generally employed to control hemorrhage from vessels of the extremities, may be used to control the femoral artery as it crosses the brim of the pelvis, by placing a compress over the artery in this position, and then applying the elastic band to secure it with a figure-of-eight turn, passing it under the thigh, crossing over the pad, and then carrying the ends around the pelvis, and securing them.

To make pressure on the axillary artery, a compress should be placed in the axilla, and the middle of the tube is placed over this to hold it in position; the ends of the tube are then carried over the shoulder and crossed, and then carried to the opposite axilla and secured.

In amputation at the shoulder-joint, to make pressure upon the subclavian artery, which is difficult to compress by an ordinary tourniquet, the handle of a large key well padded may be used; it is firmly pressed against the vessel above the clavicle, and held by an assistant, and

will prove a very satisfactory means of controlling the circulation in this vessel.

Hæmostatic Forceps.

The temporary control of arterial hemorrhage by the use of hæmostatic forceps is now very generally employed in surgical operations, and their use has done much to diminish the shock following operations from the loss of blood. The hæmostatic forceps in general use is self-retaining; it is clamped upon the bleeding vessel, and is allowed to remain until the operation is completed, when the vessel is secured permanently by the application of a ligature, and the forceps is removed. The use of these forceps will be found very satisfactory in controlling hemorrhage during the removal of tumors and in cases of amputation, and for the temporary control of bleeding during the operation of tracheotomy they will be found most efficient, as also in abdominal operations, in which their utility was first demonstrated. (Fig. 185.)

Fig. 185.



Hæmostatic forceps.

Esmarch's Bandage and Tube.

This apparatus, which is applied to the limbs to render them bloodless during operations, consists of a rubber bandage two and a half inches in width and three or four yards in length, and a rubber tube two yards in length, to one end

of which is attached a chain and to the other a hook, or better a rubber strap, one inch in width and one and a half yards in length with a hook and chain. The bandage is applied to the extremity of the limb and is carried up the limb to a point some distance above the seat of proposed operation; the bandage is applied firmly, each turn overlapping one-fourth of the preceding one, and when the last turn has been made the rubber tube or strap is wound firmly around the limb and secured by fastening the hook into one of the links of the chain. (Fig. 186.) After securing the tube or strap the rubber bandage is removed from the limb

FIG. 186.



Esmarch's bandage and tube applied.

and if the tube has been firmly enough applied the limb will be found to be blanched, and should be free from blood during the operation. Care should be taken not to apply the tube or strap too tightly in poorly developed limbs, or on parts of the limb where large nerve trunks approach the surface, as they may be subjected to an amount of pressure which will interfere with their functions subsequently. I have knowledge of one case of this nature in which permanent paralysis of the limb followed the use of Esmarch's apparatus; the tube should be applied with just enough firmness to control the circulation.

As the strap, when firmly applied, completely cuts off the circulation of parts below, it should be applied for as short a time as possible, as gangrene has resulted from its prolonged use.

After the removal of the tube there is generally quite free capillary hemorrhage, due to paralysis of vasomotor nerves from pressure, but this in a short time stops. This apparatus is of the greatest service in controlling hemorrhage at the time of operation, and in amputations and removal of vascular tumors from the limbs will be found most satisfactory. In operations upon bone, either osteotomy or sequestrotomy, it is especially useful, as it allows the surgeon to have a view of the parts unobscured by hemorrhage. I have found its use most satisfactory in operations for the removal of foreign bodies, such as needles imbedded in the hands or feet or extremities.

Permanent Control of Arterial Hemorrhage.

To secure this end the surgeon may resort to the use of position, cold, heat, styptics, pressure, cauterization, ligation, torsion, or acupressure.

Position.

In arterial hemorrhage from wounds of the extremities elevation of the part will be found to materially diminish the amount of hemorrhage; in hemorrhage from wounds of the arteries of the hand, forearm, foot, or leg, forcible flexion of the forearm on the arm or of the leg on the thigh will be found useful in diminishing the force of the blood-current.

Cold.

The application of cold by means of a stream of cold water or an ice-bag or pieces of ice will often be found an efficient means of controlling hemorrhage from vessels of small calibre; it is especially applicable to hemorrhage from wounds of the vessels of the mouth, nostrils, vagina, or rectum.

Hot Water.

Hot water will be found a very efficient means of controlling hemorrhage from small vessels, and it may be used

in the form of a hot antiseptic solution. It is of especial value in capillary or parenchymatous hemorrhage and is employed in the form of a douche or by means of sponges dipped in the hot solution and packed into the wound. Injection of hot water is a most satisfactory method of controlling uterine hemorrhage.

Styptics.

These agents are sometimes employed to control capillary bleeding or hemorrhage from small vessels, and although their use is often satisfactory as regards the control of the bleeding, they have the disadvantage of interfering with the primary union in wounds, and since the value of asepsis in wound treatment has been demonstrated they are now very seldom employed. The most valuable styptics which are used are alcohol, alum, oil of turpentine, perchloride of iron, and persulphate of iron or Monsel's solution, acetic acid, and vinegar.

Pressure.

For the permanent control of arterial hemorrhage pressure may be applied directly to the bleeding-point or surface by means of a compress of antiseptic gauze or by strips of gauze packed firmly into the cavity from whose surface the bleeding arises.

Compresses are used with the best results where the proximity of a bone gives a firm substance upon which the vessel may be compressed, as is the case in the vessels of the scalp. Pressure applied by means of packing with strips of gauze will be found most efficient in controlling hemorrhage from cavities such as the nose, vagina, or rectum, and in the cavities resulting from the removal of necrosed or carious bone. Pressure may be indirectly applied by flexing the proximal joint over a compress or by firm bandaging of the limb.

In controlling bleeding from a divided artery in a bony cavity, such as the inferior dental, a piece of catgut ligature may be forced into the canal, and will control the bleeding in a most satisfactory manner.

The troublesome hemorrhages sometimes occurring after the removal of a tooth may be controlled by packing the alveolar cavity with a strip of gauze, or by introducing a wedge-shaped piece of cork and holding it in place by fastening the jaws together by means of a bandage.

Cauterization.

The use of cauterization by means of a hot iron is a satisfactory method of arresting hemorrhage. Care should be taken to have the iron only of a dull-red or black heat, as the result desired is not the destruction of the tissues, but the coagulating effect of heat upon them. The form of cautery iron employed will depend upon the position of the vessel. Paquelin's cautery is also a satisfactory apparatus to use for the control of hemorrhage.

Control of arterial bleeding by cauterization is often resorted to in operations upon the jaws and in the removal of tumors from the mouth or pharynx or of the tonsils; it is also frequently employed to control hemorrhage in operations upon the uterus and the rectum, and also that resulting from the removal of abdominal tumors, where the application of a ligature is difficult and often impossible.

Torsion.

This method of controlling arterial hemorrhage consists in seizing the end of the artery, drawing it slightly out

FIG. 187.



Hewson's torsion forceps.

of its sheath and twisting it; it may be accomplished with a single pair of forceps or by two pairs of forceps. In the latter method the vessel is held by one pair of forceps and is twisted by the second pair.

Torsion of arteries in accidental wounds is quite common, and in many cases controls the hemorrhage until surgical aid is rendered. I have seen the femoral artery in Scarpa's triangle completely controlled in this manner in the case of an avulsion of the thigh from railway injury.

In vessels of moderate size it may be practised with one pair of forceps, and the ordinary double-spring artery forceps (Fig. 188) will be found satisfactory for such cases. In larger arteries two forceps should be employed, or some of the numerous forms of torsion forceps which have been devised for this purpose. (Fig. 187.)

Fig. 188.

Double-spring artery forceps.

Ligation.

The use of the ligature is by far the most generally employed method of controlling arterial hemorrhage. The materials used for ligature are silk, hemp thread, catgut, horsehair, iron or silver wire. Catgut or silk is the material generally employed. The vessel is seized with a pair of artery forceps or a tenaculum (Fig. 189) and drawn out of its sheath,

Fig. 189.



Tenaculum.

and a ligature of prepared catgut is thrown around it and secured by a surgeon's knot, or by a reef knot and surgeon's knot combined, and when firmly tied the ends are cut short

in the wound. Silk ligatures which have been rendered aseptic are applied in the same manner and the ends may be cut short in the wound.

When ligatures are applied to vessels in their continuity they may be threaded into an eyed probe or aneurism needle (Fig. 190) and carried around the vessel and secured.

FIG. 190.

Aneurism needle armed with ligature.

A convenient method of applying a ligature to a bleeding-point in a deep wound or to a vessel in tissues which are of such a nature as not to permit of the isolation of the vessel, is to use a curved needle threaded with a catgut ligature,

FIG. 191.

Artery occluded by suture. (ESMARCH.)

which is passed deeply into the tissues near the vessel and brought out on the opposite side; the ligature thus placed is then firmly tied, and the ends are cut short in the wound. (Fig. 191.)

Acupressure.

In this method of controlling arterial hemorrhage a needle or pin is used which is thrust through the tissues in such a way as to compress the artery. There are a number of methods of using the needle or pin and a few of these will be described.

First Method of Acupressure.

In this method the surgeon places a finger of his left hand upon the mouth of the bleeding vessel and with his right hand introduces the needle from the cutaneous surface and passes it through the thickness of the flap till its point projects for a couple of lines or so from the surface of the

FIG. 192.

Acupressure—first method; raw surface. (ERICHSEN.)

FIG. 193.

Acupressure—first method; cutaneous surface. (ERICHSEN.)

wound a little to the right side of the tube of the vessel. By forcibly inclining the head of the needle toward his right he brings the projecting portion of its point firmly down on the site of the vessel, and after seeing that it occludes the artery he makes it reënter the flesh as near as possible to the left side of the wound and pushes the needle through the flesh till its point comes out again at the cutaneous surface. (Figs. 192 and 193.)

Second Method of Acupressure.

A straight needle threaded with a short piece of iron or silver wire, for the purpose of afterward retracting and

removing it, is passed down through the soft parts a little to one side of the vessel; its point is then raised up and passed over the artery and is then turned down again and thrust into the soft tissues on the other side of the vessel. (Fig. 194.)

Third Method of Acupressure.

In this method the point of the needle is passed into the tissues a few lines to one side of the vessel, then passed under it and afterward pushed on, so that the point again emerges a few lines beyond the vessel. A loop of wire is next passed over the point of the needle and then after

FIG. 194.



Acupressure—second method. (ERICHSEN.)

FIG. 195.



Acupressure—third method. (ERICHSEN.)

being carried over the vessel and passed around the opposite end of the needle it is drawn sufficiently tight to close the vessel, and the ends of the wire are secured by making a twist around the stem of the needle. (Fig. 195.)

Fourth Method of Acupressure.

This method is identical with the third, except that a long pin is used in place of the needle, the head of the pin remaining outside the wound.

Fifth Method of Acupressure.

This method consists in passing a pin or needle through the soft tissues close to the artery, and by giving the pin a quarter or half rotation twisting the vessel upon itself,

fixing the pin by thrusting its point deeply into the tissues beyond. (Fig. 196.)

Fig. 196.



Acupressure—fifth method. (Erichsen.)

Sixth Method of Acupressure.

This method consists in applying the pin as in the fourth method, but differs from it in crossing the ends of the wire behind the pin so as to embrace the mouth of the vessel between them.

Seventh Method of Acupressure.

This method consists in passing a long needle or pin through the cutaneous surface deeply into the soft parts at some distance from the vessel, making it emerge near the vessel, bridging over the vessel and then thrusting it down into the soft parts on the other side of the vessel and making its point emerge again from the integument.

Treatment of Venous Hemorrhage.

Bleeding from small veins often stops spontaneously unless there is some pressure upon the wounded veins upon the cardiac side of the wound. It is, however, very satisfactorily controlled by position or by the application of a compress and bandage, or by the use of a ligature; if the divided vein be a large one it is well to secure both ends of the vein by ligatures. The free bleeding arising from ruptured varicose veins of the leg is easily controlled by the application of a compress and bandage, while hemorrhage

from the larger veins, such as the jugular, should be controlled by the application of ligatures as in the case of wounded arteries. The application of the *lateral* ligature to small wounds of veins of large size, such as the femoral, has been recommended, and it consists in pinching up the wall of the vein so as to include the orifice of the wound and throwing a delicate ligature around it.

The use of the actual cautery may also be required for the control of venous hemorrhage in positions in which its arrest by pressure or the ligature is not feasible.

Treatment of Capillary Hemorrhage.

Capillary or parenchymatous hemorrhage is usually arrested spontaneously by the exposure of the injured surface of the wound to the air, but it is often so profuse that its arrest becomes a matter of importance. To control this form of bleeding, pressure may be applied to the bleeding surface for a short time, and if this fails to arrest it, sponging the surface with dilute alcohol will sometimes prove satisfactory; but the best application to arrest hemorrhage of this nature is hot water, which may be used in the form of a hot bichloride solution. Acetic acid and vinegar are also sometimes employed for the same purpose. In cases where the means mentioned above fail to control the bleeding, it may be necessary to pack the wound with strips of antiseptic gauze; this dressing is most serviceable when the hemorrhage comes from cavities such as result from the removal of tumors or excisions of joints, and for the control of bleeding following the removal of necrosed or carious bone, packing the cavity resulting is the method very generally employed. To control hemorrhage from the mucous cavities, such as the nose, rectum and vagina, this method of treatment is frequently resorted to.

Treatment of Secondary Hemorrhage.

Secondary hemorrhage following the use of the ligature or other means of controlling bleeding is, since the adop-

tion of the antiseptic method of wound-treatment, a much less frequent complication of wounds. The treatment of this complication is both constitutional and local; the constitutional treatment consists in the use of those remedies which were mentioned as serviceable in primary hemorrhage, and the drugs upon which the most reliance is to be placed are opium and ergot.

The local treatment of this form of hemorrhage consists in the use of the various means of controlling hemorrhage which have been mentioned before, such as the ligature, hot water, pressure, or the actual cautery. If possible, it is well to secure the vessel from which the bleeding arises in the wound; if for any reason this cannot be done, the main artery should be ligated above the wound if the hemorrhage be arterial.

Rules for Ligating Wounded Arteries.

The following rules for the application of ligatures to wounded arteries are laid down by Ashhurst:

1. In cases of primary hemorrhage, no operation should be performed upon an artery, unless it is at the moment actually bleeding. The exception to this rule is in the cases where the vessel is seen to pulsate in the wound or where the wound involves the region of a large artery and the patient has to be transported or may be in a position not to receive surgical aid subsequently if needed; under these circumstances, the vessel should be tied or the wound should be explored to ascertain the fact that no important vessel has been injured.

2. In applying a ligature to a wounded artery, the surgeon should cut down directly upon it at the point from which it bleeds and secure it in the wound.

This rule holds good for both primary and secondary hemorrhage.

3. Two ligatures should be applied, one to each end of the artery if it be completely divided, and one on each side of the wound if the latter has not completely severed the coats of the artery. This procedure is adopted for the

reason that the arterial anastomosis is so free that the proximal ligature will not always, even temporarily, arrest the bleeding; and if it does accomplish this object at the time, after the collateral circulation is established, bleeding is apt to occur from the distal extremity of the divided vessel. If the coats of the artery are not completely severed their division should be completed, either before or after the application of the proximal and distal ligatures, thereby favoring the contraction and retraction of the ends of the divided vessel.

Control of Hemorrhage from Special Parts.

Epistaxis or hemorrhage from the nose may be so profuse as to require surgical interference. To control this form of hemorrhage the application of iced compresses to the surface of the nose may first be made use of, and if this fails to control the bleeding, the surgeon or the patient should grasp the cartilaginous portion of the nose with his thumb and forefinger in such a manner as to keep the nostrils tightly closed, which will prevent the passage of air through the nose and thus permit clots to form, arresting the flow of blood. If these simple means fail to arrest the bleeding the nasal cavity or cavities may be packed with strips of antiseptic gauze introduced into the anterior nares, and pushed backward by a director or probe; this will often be found a perfectly satisfactory means of arresting the bleeding. This method may be supplemented by a plug of antiseptic cotton introduced into the posterior nares with the fingers. The use of a rubber tampon, consisting of a rubber bag introduced into the nares in an empty state and afterward inflated, has also been recommended for the control of this variety of hemorrhage.

Another method of controlling hemorrhage from the nose consists in introducing a small piece of sponge, tied to a strong silk ligature, into the anterior nares and pushing it back along the floor of the nose to the posterior nares; a small piece of sponge about the size of a marble with a

hole in the centre is threaded on the ligature and pushed back until it comes in contact with the first piece of sponge introduced, and thus by introducing a number of pieces of sponge in this way the nasal cavity may be completely filled up and the bleeding is arrested. Care should be taken to see that the sponge has been rendered aseptic before being introduced, and the nasal cavity should also be washed out with an antiseptic solution before its introduction. The sponges may be allowed to remain in place for twenty-four to forty-eight hours. (Fig. 197.)

Fig. 197.

Plugging the nares from the front. (Roberts.)

Plugging the nares by means of Bellocq's canula is also employed to arrest hemorrhage from the nasal cavities; the canula, armed with a strong ligature, is passed along the floor of the nose until it reaches the pharynx, when the spring being protruded, the ligature is seized and brought

out of the mouth and secured to a plug of lint or antiseptic gauze of the required size, and upon withdrawing the instrument the plug is brought into position in the posterior nares and the end of the ligature is allowed to protrude from the mouth to facilitate its removal. (Fig. 198.) An

FIG. 198.

Plugging the nares with Bellocq's canula. (FERGUSSON.)

ordinary flexible catheter may be employed in place of Bellocq's canula for the introduction of the ligature.

Hemorrhage from the Urethra.

In hemorrhage from the urethra, if profuse, the blood will trickle from the meatus, or if efforts at micturition are made the first gush of urine will contain blood, but afterward will be clear, and the last urine will contain a few drops of pure blood.

This variety of bleeding, if it proceeds from the anterior portion of the urethra, may be controlled by the introduction of a catheter and the application of a bandage around the penis, carefully applied so as to make only moderate pressure.

If the bleeding comes from the posterior portion of the urethra, it will often be controlled by the application of cold or pressure to the perineum, or by the introduction of a cold steel bougie, or by the injection of a solution of tannic acid.

Hemorrhage from the Bladder.

In this variety of hemorrhage the first portion of the urine may be blood-stained and the last portion will contain more blood and clots as the organ contracts, which distinguishes it from hemorrhage from the kidneys, in which the admixture of blood with the urine renders it of a smoky color or dark-red if the bleeding is profuse.

To control bleeding from the bladder a catheter should be introduced and the urine and clots withdrawn; the bladder should next be washed out with a warm or cold boric acid solution, or in severe cases weak astringent solutions, such as tannic acid or alum, may be employed. The application of ice to the perineum and supra-pubic regions may also be employed with advantage.

Hemorrhage from the Rectum.

This variety of bleeding may be controlled by the injection of cold or astringent enemata. If the bleeding be profuse a speculum should be introduced, and when the source of the bleeding has been discovered the actual cautery or a ligature should be applied. If this is not feasible the rectum may be plugged with strips of antiseptic gauze, or a piece of a rubber catheter of large calibre may be wrapped with gauze and introduced into the rectum, the end of the catheter being allowed to protrude; by using this tube flatus can escape, and if the bleeding is not controlled blood will escape through the tube, preventing the risk of concealed hemorrhage. If the bleeding arises from hemorrhoids or polypus of the rectum, the operative treatment of these conditions should be undertaken to permanently control the bleeding.

Opening and Dressing of Abscesses.

Acute abscesses, as a rule, should be opened by incision, and this is best done with a straight, narrow, sharp-pointed bistoury; the incision should be deep enough to freely expose the cavity of the abscess, and should be so planned as to be parallel with and not across important structures, and it should also be made at as dependent a portion as possible. Abscesses of the limbs are opened by a longitudinal incision, and those in the region of the anus and breast by an incision radiating from the anus or nipple.

In deep-seated abscesses in the region of important structures the method of opening suggested by Mr. Hilton may be employed with advantage; it consists in making a small incision through the skin and cellular tissue; a director is next pushed through the tissues into the abscess cavity, which will be shown to have been reached by the escape of a little pus along the director; a dressing forceps with the blades closed is now pushed along the director into the abscess cavity, and when this has been accomplished the director is withdrawn and the forceps are removed with the blades expanded so as to dilate the wound and allow the pus to escape.

The cavity of the abscess being emptied of pus, it should be irrigated with a stream of carbolic acid solution 1 : 40, or bichloride solution, and if the cavity is not very large or deep no drainage-tube need be introduced, and a small piece of protective may be placed between the lips of the wound to prevent their adhesion; but if, on the other hand, the cavity is extensive and deeply situated, a rubber drainage-tube should be introduced to the bottom of the cavity to secure free drainage and fixed at the surface of the skin by a safety-pin. A piece of protective which has been dipped in bichloride solution is next placed over the wound, and over this is laid a gauze dressing, consisting of a number of layers, which has been moistened in carbolic or bichloride solution; this is covered by a number of layers of dry gauze which is in turn covered by a piece of rubber tissue, and

over this is placed a few layers of bichloride cotton and the dressing is finally secured by a roller bandage. The dressing is removed at the end of two or three days, the cavity being washed out with one of the antiseptic solutions previously mentioned; the drainage-tube may be shortened or removed and the dressings are reapplied as at the primary dressing. Under this method of treatment acute abscesses usually heal more promptly and with less suppuration than under the older methods of treatment in which poultices were applied.

Chronic or cold abscesses, which occur chiefly in connection with diseases of the bones or joints or of the lymphatic system, and are generally tubercular in their origin, may be opened in various ways, the time at which this should be done depending upon the size and situation of the abscesses and the amount of constitutional disturbance which the patients experience from their presence.

A cold abscess may be evacuated by means of the aspirator; the pus being withdrawn as far as possible, the puncture is sealed with a small piece of gauze covered with iodoform collodion. Reaccumulation of the pus often takes place and the aspiration has to be repeated a number of times. The greatest difficulty in the successful removal of the contents of cold abscesses by means of aspiration is the presence of masses of lymph in the pus which occlude the canula and often prevent the complete emptying of the cavity.

These abscesses may also be evacuated by making a puncture through the skin and overlying tissues with a narrow bistoury, the surface having been previously thoroughly washed with soap and water and with a carbolic or bichloride solution; a director is next pushed through this small wound into the cavity of the abscess and the pus is allowed to escape by stretching the wound by the director; when the cavity is emptied of pus it is washed out with a carbolic or bichloride solution introduced into it by pushing the nozzle of a syringe into the cavity, and this is allowed to escape in the same way as the pus previously did, and when the irrigating solution has all escaped the cavity may

be injected with an emulsion composed of iodoform one part, glycerin ten parts; after this has been introduced the small wound is closed by a compress of antiseptic gauze held in place by a compress of bichloride cotton and a bandage or by strips of adhesive plaster. The injection of the iodoform emulsion need not be repeated as long as iodoform continues to be excreted with the urine. Cold abscesses are also treated by making a free incision into the abscess cavity with full antiseptic precautions, and after the escape of the purulent matter the walls of the abscess should be thoroughly scraped with a curette, and after the cavity has been freely washed out with a carbolic or bichloride solution large drainage-tubes are introduced and an antiseptic dressing is applied to the wound. The dressings are removed as soon as they become soaked and the drainage-tubes are shortened or removed as the discharge diminishes and the cavity contracts.

In evacuating chronic abscesses by means of the aspirator or by a small puncture, there is absence of shock and the loss of blood is insignificant, so that these procedures should generally be first employed, and the more radical operation of incision and curetting of the cavity of the abscess, which is accompanied with a certain amount of shock and hemorrhage, should be reserved for those cases in which the less severe operations have failed to be followed by a satisfactory result.

Diffused suppuration is treated by numerous punctures or incisions, which allow the purulent matter to escape, and where sloughs are present free incisions may be required to give exit to the necrosed tissues; the introduction of drainage-tubes may also be required. The wounds and the cavities, as far as possible, should be washed out with a carbolic or bichloride solution and an antiseptic gauze dressing should be applied.

Sinuses resulting from abscesses, if superficial, should be laid open freely and their surfaces should be scraped with a curette and they should then be lightly packed with strips of bichloride or iodoform gauze and should be covered by an antiseptic dressing. If they are too deep to be treated

by incision their healing may be facilitated by the injection of stimulating solutions introduced by means of a syringe; the employment of solutions of chloride of zinc, nitrate of silver, and sulphate of copper varying in strength from five to twenty grains to the ounce of water will often prove satisfactory.

Dressing of Wounds.

Incised wounds present the conditions favorable for prompt healing and they should first be carefully irrigated with an antiseptic solution to remove any blood-clots or foreign bodies, and after any hemorrhage which is present is controlled by the use of ligatures, if the wound be an extensive or deep one, provision should be made for drainage by introducing a drainage-tube or a few strands of prepared catgut to the bottom of the wound, allowing the extremity to project from the most dependent portion of the wound. In superficial incised wounds, after the hemorrhage has been controlled, it is not usually found necessary to make any provision for drainage. If the wound be a deep one, involving the muscles and deep fascia, buried sutures of catgut should be applied to approximate the muscles and fascia, and if important nerves or tendons have been divided their ends should be brought into apposition by sutures of catgut or sterilized silk; the superficial portions of the wound should next be brought together by the introduction of a number of interrupted sutures, catgut, silkworm-gut, silver wire or silk being employed for this purpose; the accurate apposition of the edges of wounds of this variety is secured by the introduction of a number of sutures placed closely together.

After a wound of this variety has been closed the subsequent dressing is accomplished by dusting the surface of the wound with iodoform or aristol, and a piece of protective a little larger than the wound, which has been dipped in a 1 : 40 carbolic solution, is placed over it; over this is placed a pad of antiseptic gauze, composed of ten or twelve layers, which has been soaked in a 1 : 40 carbolic solution or a

1 : 2000 bichloride solution, and over this is laid a pad of dry antiseptic gauze of the same thickness, overlapping the wet gauze by a few inches in all directions; a few layers of bichloride cotton are next applied over the gauze dressings and the whole dressing is secured in position by the application of an antiseptic gauze bandage. Under this form of dressing prompt healing of incised wounds is the rule, and the wound need not be re-dressed for a week or ten days unless some indications exist for the change of dressing at an earlier period. At the time of the first dressing the catgut drain or the drainage-tube is usually removed and if the adhesion of the edges of the wound is firm the sutures may also be removed. An antiseptic dressing is usually next applied and allowed to remain in position for a few days longer.

Lacerated wounds present edges which are torn and not sharply cut, and the vitality of the injured parts is often so seriously impaired that prompt union in this variety of wounds is not, as a rule, to be looked for. Wounds of this nature should first be irrigated with an antiseptic solution, as in the case of incised wounds, and blood-clots and foreign bodies should be removed. If the wounds be deep, drainage-tubes should be introduced; on the other hand if they be superficial or if the edges are not closely approximated, provision for drainage may be omitted. The torn or irregular edges of the wound should next be brought into apposition at a few points, by the introduction of a few catgut or silkworm-gut sutures, applied not very closely together; and if the edges are discolored and their vitality seems markedly impaired, it is better not to use sutures, but rest satisfied by bringing them as nearly as possible in contact by the use of a few strips of isinglass plaster moistened with a bichloride solution. If the edges of the wound are so much crushed as to have their vitality destroyed, they may be trimmed away with scissors until a surface possessing fair vitality is secured. The evil results arising from the introduction of sutures into this variety of wounds with the idea of closely approximating their edges are so common, that the surgeon who dispenses with the use of sutures en-

tirely errs upon the safe side. The use of many sutures in wounds of this nature often causes marked tension in the wound, which is frequently followed by impairment of the vitality of the injured tissues and sloughing results.

The wound should next be dressed antiseptically, and if it runs a favorable course it need not be re-dressed for a week or ten days; the time required for the repair of a wound of this nature is longer than that for an incised wound, and a larger number of dressings may be required.

In lacerated wounds of the extremities continuous irrigation of the wound by a warm bichloride or carbolic solution, applied as described (page 144), is often followed by the most satisfactory results; wounds produced by machinery and railway accidents, in which the vitality of the tissues is much impaired, are particularly favorable cases for this method of treatment, and here the same caution should be exercised as regards the introduction of sutures.

Contused Wounds.—This variety of wounds possesses many characteristics in common with lacerated wounds; the edges are bruised and the injury of the subcutaneous tissue is often more extensive than the size of the external wound would lead one to suspect. They are dressed in the same manner as lacerated wounds, and the same objection here exists to the use of sutures as in the latter class of injuries.

Punctured Wounds.—These wounds are inflicted by sharp-pointed instruments, and it often happens that a portion of the vulnerating body remains in the wound, as is frequently the case in wounds produced by needles; and another complication in this variety of wound is the injury of vessels, giving rise to concealed hemorrhage, or of nerves resulting in neuritis. Simple punctured wounds should be carefully washed with an antiseptic solution and covered by an antiseptic gauze dressing, and if no complication exists their healing is usually very rapid.

When, however, a foreign body remains in the wound, as it often happens in punctured wounds produced by needles and pins, the punctured wound should be converted into an incised wound, and the body should be searched for and

removed if possible, and in doing this in the case of wounds of the extremities the operation is much facilitated by the employment of Esmarch's bandage and strap. After the removal of the foreign body the wound is treated as an incised wound, and an antiseptic dressing should be applied. When concealed hemorrhage occurs after a punctured wound, the wound should be laid open and the bleeding vessel searched for and ligatured if possible, and the wound should afterward be dressed as an incised wound.

Poisoned Wounds.—These wounds are caused by the absorption, by means of a cut or abrasion in the skin, of fluids from a dead body in making dissections or post-mortem examinations or in operating upon living subjects, and often result in serious consequences. Such wounds, as soon as possible after their reception, should be carefully washed out with a solution of bichloride of mercury, 1 : 2000, or a 30-grain solution of chloride of zinc, and then dressed with an antiseptic dressing. If, however, this precaution is not taken or the wound has escaped notice, and in a few hours becomes inflamed and painful, and evidences of lymphatic involvement show themselves, the wound should be opened and its surface should be thoroughly washed out with a 30-grain solution of chloride of zinc, and finally with a 1 : 2000 bichloride solution, and it should then be dressed with an antiseptic gauze dressing. Under this method of dressing the poisoned wound is often converted into a healthy one, even after the lymphatic involvement is well marked, and it usually heals promptly without further constitutional disturbance.

Gunshot Wounds.—These wounds are produced by small shot, balls, or fragments of shells, and are of the nature of contused and lacerated wounds, and the vulnerating body as well as portions of the clothing are often imbedded in the tissues.

In dressing these wounds any foreign bodies, if they can be located, should be removed, and in the search for and removal of balls from the extremities the application of the Esmarch bandage and strap will be found most useful. The wound should next be thoroughly washed out with a 1 : 2000

bichloride solution, and an antiseptic dressing applied as in the case of other contused and lacerated wounds.

Powder burns resulting from the explosion of powder, in addition to the burning and laceration of the tissues, are accompanied by the introduction of grains of unburnt powder into the skin, which, if not removed, leave permanent points of pigmentation. These wounds should first be washed with an antiseptic solution, and upon the face, to avoid unsightly pigmentation of the skin, care should be taken to pick out the small masses of powder with a needle or the sharp point of a tenotomy knife. The surface should then be dressed with lint spread with an ointment of boric acid or an ointment of aristol, consisting of half a drachm or a drachm of aristol to an ounce of vaseline, this dressing being covered by a few layers of bichloride or borated cotton, held in place by a roller bandage.

Contusions or *bruises* differ from contused wounds in the fact that the skin is not broken, though in spite of this fact there may exist very extensive laceration of the subcutaneous tissues, accompanied by more or less extravasation of blood from the injured vessels. When not severe enough to require operative treatment, they should be dressed by applying over them several layers of lint saturated with lead-water and laudanum, and over this dressing is placed a layer of waxed paper or rubber tissue, and the dressing is secured by a roller bandage.

A solution which I find most satisfactory in the dressing of contusions is as follows:

Ammonii chloridi	grs. xx.
Tr. opii }	
Alcoholis }	āā f℥j.
Aquæ	q. s. ad f℥j.

Several layers of lint saturated with this solution are laid over the contused tissues, and are covered with waxed paper, oiled silk, or rubber tissue.

Extensive collections of blood following contusions often remain in the tissues for some time, but usually are absorbed. If this result does not follow, or an abscess forms,

the blood or pus should be removed by aspiration or by incision with full antiseptic precautions.

Burns and Scalds.

The dressings employed in the treatment of burns and scalds are similar, as the injury to the tissues is practically the same in both classes of injuries. Superficial burns or scalds, in which the effect of the heat has only extended to the superficial layer of the skin, may be treated by the application of lint saturated with a solution of carbonate of sodium, a drachm to an ounce of water; this dressing rapidly relieves the pain, and is a satisfactory application in this variety of burns and scalds. In cases in which the effects of heat have extended to the deeper tissues, the affected surface may be dressed with *carron oil*, which is prepared by rubbing together lime-water and linseed oil until a thick creamy paste results; lint is saturated with this mixture and laid over the surface of the burn or scald. The dressing is a comfortable one to the patient, but soon becomes offensive, and for this reason requires frequent renewals.

The disadvantage met with in the antiseptic method of dressing burns and scalds is the fact that the raw surface presented offers the most favorable conditions for the absorption of the antiseptic substances employed in the dressings, and for this reason the use of bichloride of mercury, carbolic acid, and iodoform is not to be recommended in burns or scalds involving a large extent of surface, on account of the toxic symptoms which may result from their employment.

A recent burn or scald, by reason of the heat employed in its production, is practically an aseptic wound, and it may be dressed by covering it with boric acid ointment, and placing over this a number of layers of borated or salicylated cotton, and holding the dressings in position by a roller bandage.

Aristol, as a substitute for iodoform, may be employed in the dressing of burns or scalds, being either dusted over the

surface or used in the form of an ointment, and over this application should be placed a few layers of borated or salicylated cotton.

When blebs are present upon the surface of the burn or scald, they should be opened to allow the serum to escape. If suppuration occurs or the tissues become necrosed by reason of the severity of the injury, the surface of the burn may be washed with a 1 : 60 carbolic solution or 1 : 4000 bichloride solution and the same dressing should then be applied.

The ulcers resulting from the separation of the dead tissues should be touched with a solution of nitrate of silver, four grains to the ounce of water, and dressed with lint spread with ointment of boric acid or aristol. In the dressing of extensive burns or scalds of the neck, face, and region of the joints, the possibility of serious deformity from contraction of the tissues in healing should not be lost sight of, and position, splints and bandages, should be employed to prevent, as far as possible, this complication.

Bedsores.

These sores usually occur over the sacrum or hips in patients who are confined to bed for a considerable time, as the result of a long-continued pressure, or in those cases where the vital powers are depressed by adynamic diseases, and are also a frequent and troublesome complication in spinal injuries. Their formation may be prevented in many cases by the use of air-cushions or of a water mattress, and by keeping the parts exposed to pressure scrupulously clean and frequently bathing them with stimulating lotions, such as alcohol, olive oil and alcohol equal parts, or soap liniment. The parts should also be protected from pressure by the application of adhesive plaster, or, still better, soap plaster spread upon chamois. When the bedsore has actually formed, and in many cases its formation is very rapid and the slough will be found to involve a large surface of the skin over the sacrum, and to extend down to the bone,

we have present a very serious complication, and one which requires most careful treatment.

The dressing of a bedsore before the separation of the slough consists in relieving the part from pressure by the use of an air-cushion placed under the buttocks, and the application of a fermenting poultice until the slough has separated. When the slough has become detached the ulcer remaining should be well washed with a carbolic or bichloride solution and the granulations should be touched with a 5-grain solution of nitrate of silver; and resin cerate, iodoform, aristol, or boric acid ointment, spread upon lint, should be applied to the surface of the ulcer, and a piece of soap plaster a little larger than the ulcer should be placed over this dressing and held in place by broad strips of adhesive plaster. This dressing should be renewed every day or every other day, and means should be adopted to protect the parts from further pressure, and the constitutional condition of the patient should be improved by the administration of a nutritious diet, tonics, and stimulants. The application of the galvanic current has been employed with good results to promote the healing of the ulcer in obstinate cases.

Sprains.

Sprains of joints from twists or other external violence resulting in the stretching or laceration of the ligaments are injuries which require careful dressing.

Sprains may be first treated by the application of cold- or hot-water dressings for a few hours, or by the application of lead-water and laudanum, the joint being kept at rest by the use of a splint or by confining the patient in the recumbent posture in the case of sprains of the joints of the lower extremities.

After a few days' use of the lead-water and laudanum dressing the swelling usually subsides and the joint may be fixed by the application of a moulded soap-plaster splint or felt splint held in place by a firmly applied roller bandage, which should be worn for a week or ten days; in ordi-

nary cases after this time the splint may be removed and the patient should be encouraged to use the joint. In cases of severe sprain, on the other hand, the pain and swelling persist for some time, and here the fixation of the joint by a soap plaster, or better by a plaster-of-Paris bandage, will be found useful for a few weeks. If upon the removal of this dressing the parts are still painful and swollen, the swollen tissues should be painted with tincture of iodine; or the method of applying tincture of iodine recommended by Mr. Jordan, that is, the application of the iodine in a broad band around and not over the swollen tissues, may be employed. The joint should next be surrounded by a piece of lint spread with an ointment composed of equal parts of ointment of mercury and ointment of belladonna, and a moulded soap-plaster splint being fitted to the joint, it is held in place by a firmly applied bandage. This will be found a most satisfactory dressing in the treatment of sprains after they have passed their acute stage. The dressing is removed at intervals of three or four days, the joint is sponged off with alcohol, and a similar dressing is reapplied; and this method of dressing may have to be continued for some weeks, but the results obtained by its continuous use are often most satisfactory. An ointment of icthyol one part to lanolin three parts may also be used in the same manner as the ointment of belladonna and mercury with good results in the treatment of these injuries. The employment of pressure in the treatment of sprains, by means of strapping, is also sometimes advantageous.

In the chronic stage of a sprain, after all dressings have been removed, the methodical use of massage is often most beneficial; and after the parts have been thoroughly manipulated a flannel bandage should be applied which, by its elasticity, gives a certain amount of support to the parts.

Sprain-fracture.—Under this name Mr. Callender has described an injury which consists in the separation of a ligament or tendon from its point of insertion, with the detachment of a thin shell of bone; this injury is apt to occur about the ankle-, knee-, elbow-, and wrist-joints, and

the treatment is the same as that of an ordinary fracture in the same locality. This injury is probably much more common than is generally supposed in connection with sprains of the joints, and is, I think, in many cases the cause of the tardy restoration of the function of sprained joints, this injury being overlooked and the injury simply being treated as a sprain, and the patient being encouraged to use the part before the union of the bone has been accomplished.

Strains of muscles and fascia varying in severity from simple stretching of the fibres to absolute rupture are treated by putting the parts at rest and by the application of pressure by means of adhesive straps or of a bandage; in strains of the muscles and fascia of the back the use of broad strips of adhesive plaster, applied as in cases of fracture of the ribs, will be found most satisfactory, and in the treatment of the later stages of the injury the employment of massage will often be followed by good results.

Tracheotomy.

This operation consists in dividing the tissues over the trachea in the median line of the neck, and after the trachea has been exposed it is opened by dividing two or three of the tracheal rings.

The operation of tracheotomy may be required to relieve the dyspnœa dependent upon membranous or diphtheritic laryngitis, growths in the larynx or trachea, growths external to these organs causing pressure upon them, œdema of the mucous membrane of the larynx or trachea from inflammation from burns or scalds, or from the inhalation of irritating gases or the swallowing of corrosive liquids. The operation may also be required for the removal of foreign bodies from the larynx, trachea, or from the bronchi, as well as for the relief of the dyspnœa due to their presence, and it is also required in cases of fracture or laceration of the larynx or trachea, and occasionally in cases of spasm of the glottis, and in cases of glossitis to overcome the

mechanical obstruction which prevents the entrance of air into the air-passages.

The ease with which the operation is performed varies much in different cases; it is, as a rule, a much simpler operation in adults than in children. In the latter subjects the shortness of the neck, the relatively greater size of the thyroid gland and the possible presence of the thymus body, the great vascularity of the parts, and the abundance of adipose tissue, render the trachea difficult to expose and open.

Under certain circumstances the operation may be performed with very few instruments; but if the surgeon has the choice he will find it convenient to have at hand two small scalpels, one short grooved director, a tenaculum, two aneurism needles which may be used as retractors, one pair of artery forceps, hæmostatic forceps, two pairs of dissecting forceps, a pair of scissors, a sharp-pointed tenotome, a pair of tracheal forceps, a tracheal dilator, tracheotomy tubes, tapes, ligatures, sponges, a flexible catheter, and feathers. The *director* should be short: the ordinary grooved director is too long to use with satisfaction in operating upon the short necks of children: so that I have had made a shorter and somewhat broader one, which has a bevelled extremity which allows it to be passed with ease between the different layers of the tissue. (Fig. 199.)

Fig. 199.



Author's tracheotomy director.

Hæmostatic forceps are also of great use in controlling hemorrhage during the operation in case of the division of vessels which bleed freely, when the operator from the urgency of the case does not think it justifiable to ligature them at the time of their division. They may also be employed under similar circumstances to clamp the isthmus of

the thyroid gland on either side of the trachea when it becomes necessary to divide it to expose the trachea.

A sharp-pointed tenotome is the instrument I prefer to employ in opening the trachea as its sharp point enables it to be easily thrust into the trachea, and its short cutting surface and the narrowness of the blade obscure as little as possible the line of incision and thus enable the operator to see exactly where he is cutting.

Tracheal dilators of various kinds are employed, but the most satisfactory tracheal dilator which I have employed is that of Golding-Bird (Fig. 200), which is a self-retaining instrument; the blades are slipped through the tracheal incision and are then expanded by turning the screw to which they are attached.

Trousseau's tracheal dilator, the blades of which are introduced through the incision in the trachea and are expanded by bringing together the handles, is also a satisfactory instrument (Fig. 201), but is not as useful as the tracheal

FIG 200.

Golding-Bird's tracheal dilator.

FIG. 201.

Trousseau's tracheal dilator.

dilator previously mentioned, as it has to be retained in position by the hand. Tracheal dilators may be improvised from bent hairpins or pieces of wire, which will often serve a useful purpose where ordinary dilators cannot be obtained.

It is also well to have at hand a number of pliable feathers to be used in cleaning the trachea or larynx of mucus or membrane after it has been opened, and by their use this object can be accomplished with little risk of injury to the mucous membrane.

Tracheal forceps, which are constructed with a double

spring and curved blades are also useful in removing membrane or foreign bodies from the larynx above the wound or from the trachea below the tracheal incision. (Fig. 202.)

Fig. 202.

Tracheal forceps.

Tracheotomy-tubes of various shapes are made of silver, aluminium, hard and soft rubber, but the tube which I think is the most satisfactory for general use is a silver quarter-circle tube with a movable collar (Fig. 203), and provided with a fenestrated guide. (Fig. 204.) A good tracheotomy-tube is

Fig. 203.

Silver tracheotomy-tube.

Fig. 204.

Silver tracheotomy-tube with fenestrated guide.

one which inflicts the least possible injury upon the mucous membrane of the trachea, and to insure this object the part of the tube within the trachea should lie exactly in its axis and its free extremity should be capable of as little move-

ment as possible. The tracheotomy-tube is held in position after being introduced by means of tapes attached to the shield of the tube and tied around the neck.

Position of Patient for Tracheotomy.

The best position in which to place the patient for this operation is that which brings the neck into the greatest prominence, and this can best be obtained by laying the patient upon his back upon a firm table and placing under the shoulders a round cushion; or an empty wine-bottle, or a roller-pin wrapped in towels will answer the same purpose. If an anæsthetic is not used the arms should be held by an assistant, which is better than securing them by a binder fastened around the chest, which restricts respiratory movements.

Use of an Anæsthetic in Tracheotomy.

As a rule, I think it is better not to administer an anæsthetic in performing this operation, as little pain is experienced, in cases in which the dyspnœa is well marked, after the incision in the skin has been made, and I have seen the dyspnœa which was well marked before the use of the anæsthetic suddenly become so alarming that the trachea had to be opened before it was thoroughly exposed, which is a procedure always attended with risk. So strong is my conviction that the risks of the operation are much increased by the employment of an anæsthetic that in later years I have abandoned its use.

Operation of Tracheotomy.

The trachea may be opened above the isthmus of the thyroid gland or below it, and these operations constitute respectively the *high* and *low* operations.

The *high* operation is generally selected, because at this point the trachea is more superficial and is more easily exposed, whereas in the *low* operation the trachea is more difficult to expose by reason of its relatively greater depth,

the large size and number of the veins, and its proximity to the large arterial trunks.

The patient being placed in position, the operator stands at the head of the patient; this position I prefer, as it is easier from this point to keep the incisions exactly in the median line of the neck. The operator next makes himself familiar with the landmarks of the neck; locating the position of the *cricoid cartilage*, he makes an incision through the skin in the median line of the neck from one and a half to two inches in length, the position of the cricoid cartilage being the middle point. There is no disadvantage in making a longer incision if a freer exposure of the parts is required. Having divided the skin, the operator will often see a large vein lying in the superficial fascia—the *superficial anterior jugular;* this should be displaced, and the fascia divided upon the director.

The surgeon should keep his incisions strictly in the median line of the neck, for this is the line of safety; and he should be careful, as the wound increases in depth, not to make the incisions too short, so that it becomes funnel-shaped.

When the deep fascia is exposed it should be picked up and divided upon the director, and any large veins in the line of the wound should be carefully displaced, or, if this is impossible, they should be ligatured on each side and then divided between the ligatures.

The operator now looks for the intermuscular space between the *sterno-hyoid* and the *sterno-thyroid muscles*, which can generally be found without difficulty, and the muscles are now separated in this line with the handle of the knife or with the director, and the *isthmus of the thyroid gland* will be exposed. The muscles should now be held aside by retractors placed on either side. A caution here as to the use of retractors may not be out of place: the operator should place them himself and allow the assistants to hold them. I once almost lost a case in which I had the trachea exposed, and while I turned aside to pick up a knife with which to open it, my assistant, in replacing a retractor which had slipped, included the movable trachea in the grasp of the

retractor, pulling it to one side and completely shutting off respiration; when I attempted to find the trachea to open it I could only feel the anterior surface of the vertebræ at the bottom of the wound, and it was only when I appreciated what had occurred, and lifted the retractor, allowing the trachea to spring back into its normal position, that I was able to open it. Mr. Durham and Mr. Marsh mention somewhat similar cases in which the trachea and vessels were held aside with retractors by assistants until the surgeon had exposed the cervical vertebræ.

The operator should carefully explore the wound with the finger, to locate exactly the position of the trachea, and to ascertain, if possible, the presence of any anomalous arteries.

The isthmus of the thyroid gland is exposed, which generally occupies a position over the first three tracheal rings; this is usually surrounded by a plexus of veins which should be displaced with the director, or, if this is impossible, they should be ligatured on each side and divided between the ligatures. The thyroid isthmus is next displaced upward or downward, according as the surgeon desires to open the trachea below or above this body. This is often done without difficulty, especially its upward displacement; but when there is difficulty in displacing it downward, a procedure recommended by Bose may be employed, which consists in making a transverse incision across the cricoid cartilage to divide the layer of fascia by which the isthmus is bound down; a director is then passed into this incision, and the isthmus is gently depressed without difficulty.

Having displaced the isthmus of the thyroid gland upward or downward, the trachea, yellowish-white in appearance, covered by the tracheal fascia, should be exposed; this fascia should next be thoroughly broken up with the director or handle of the knife so as to bare the trachea, and in doing this the operator can feel it crepitate under the finger from the suction of air drawn in with inspiration. Having arrived at this stage of the operation the operator should examine the wound to see that it is free from hemorrhage and he should also replace the retractors so as to expose as

large a portion as possible of the trachea, for, be the case ever so urgent, he now feels assured that he can open the trachea in a moment if the breathing should cease. The trachea is now fixed with a tenaculum introduced into it a little to one side of the median line; an incision is made into it with a narrow knife from below upward, from one-half to three-fourths of an inch in length (Fig. 205), care

FIG. 205.

Opening the trachea. (LISTON.)

being taken to see that this incision is in the median line, for if the trachea be opened by a lateral incision the wound does not heal so promptly and the tracheotomy-tube does not fit well, and its lower extremity may cause injury to the mucous membrane of the trachea. If the wound be a deep one, after fixing the trachea with the tenaculum the operator may lift it slightly from its bed, thereby bringing it more prominently into view and making it more superficial in the wound, thus facilitating its opening. As soon as the incision is made into the trachea there is a gush of air from the wound in the trachea mixed with blood or membrane; this should be wiped away with a sponge and a tracheal dilator should next be introduced and the trachea should be cleared of membrane, if it is present in the region of the wound, with a feather or with forceps. The tracheotomy-tube is

next introduced and is secured in position by tapes tied around the neck.

If respiration has ceased artificial respiration should be resorted to or the use of a tube attached to a bellows, or Fell's apparatus, and these efforts should be continued for at least fifteen minutes, for I have seen resuscitation take place in patients who were apparently dead by a persistent employment of artificial respiration.

The *care of the tube* is a matter of some importance after its introduction; the inner tube should be removed at short intervals, washed and replaced, and if the operation has been done for an inflammatory condition of the larynx or trachea a moistened feather should occasionally be passed through the tube into the trachea to withdraw any mucous or membrane which is present. In cases of croup after tracheotomy the use of a spray of steam or of a spray composed of

Carbonate of soda	℥j to ℥ijss.
Glycerin	f℥ij.
Water	f℥vj.

applied by means of a steam atomizer, the spray being directed over the opening of the tube, will be found most satisfactory in softening the discharges and thus facilitating their expulsion through the tube.

The tracheotomy-tube is usually allowed to remain in the trachea from five to ten days; its permanent removal is indicated as soon as the patient is able to breathe through the larynx with the wound in the trachea closed; its use may be required for a longer time, but as soon as the indication for its presence has disappeared the sooner it is removed the better, for its presence sometimes sets up a troublesome tracheitis. After its removal the wound rapidly diminishes in size, the healing taking place by granulation and contraction. Difficulty is occasionally met with in the permanent removal of tracheotomy-tubes; for the causes and treatment of this complication the reader is referred to special works upon tracheotomy.

Where the operation of tracheotomy is done for the re-

moval of *foreign bodies* from the air-passages, the steps of the operation are the same, but after the removal of the foreign body the treatment of the wound is somewhat different. If the foreign body has remained in the trachea only for a short time the wound in the soft parts may be closed by means of sutures or may be allowed to remain open, being covered by a piece of moistened gauze, and the use of the steam spray is here also beneficial for a few days. If, however, the body has remained in the larynx, trachea, or one of the bronchi for some time and has set up a certain amount of inflammatory trouble, it is better to introduce a tracheotomy-tube and allow it to remain for a few days. If it is found impossible to locate or remove the foreign body at the time of operation, a tracheotomy-tube should be introduced and allowed to remain until the foreign body is expelled through the tube or removed subsequently by means of forceps.

Laryngotomy.

In this operation an opening is made into the air-passages through the *crico-thyroid membrane*. It is a simple operation, and one which is practically free from risk, and can therefore be performed much more rapidly and safely in urgent cases than tracheotomy.

In this operation the same objection exists to the use of an anæsthetic as in tracheotomy, and therefore it should be dispensed with. The patient being placed in the recumbent posture, with the shoulders slightly elevated and the head thrown back to make the neck as prominent as possible, the surgeon feels for the prominence of the *thyroid cartilage*, and steadying the larynx between the finger and thumb of the left hand, he makes an incision in the median line over the centre of the thyroid cartilage and extending downward for an inch or an inch and a half. The skin and superficial fascia being divided, the fascia between the *sterno-hyoid muscles* and the areolar tissue is exposed and divided, and the *crico-thyroid membrane* is exposed. The knife is then passed transversely through the membrane into the larynx,

care being taken that both that membrane and the mucous membrane which covers its inner surface are divided at the same time. As soon as the knife enters the cavity of the larynx blood and mucus will be forcibly expelled.

The wound should be carefully enlarged and a tube introduced, which differs from the ordinary tracheotomy-tube in being slightly flattened; this is secured in position by tapes tied around the neck as in the case of the ordinary tracheal tube. The only bleeding which is likely to occur is from the *crico-thyroid arteries* or *veins*, and if these cannot be avoided, and are divided in the operation, they should be temporarily secured by hæmostatic forceps or ligatured, and if the case is not extremely urgent, all bleeding should be arrested before the crico-thyroid membrane is incised.

The after-treatment of cases of laryngotomy is similar to that of cases of tracheotomy; the same attention is required in the care of the tube and in the general management of the patient.

LARYNGO-TRACHEOTOMY.

This operation consists in making an incision into the air-passages by dividing one or two of the upper rings of the trachea, the crico-tracheal membrane, the cricoid cartilage, and the crico-thyroid membrane. This operation is employed in cases where, from the age of the patient, the crico-thyroid space is too small to admit of a sufficient opening, or in those in which, for any reason, the surgeon does not deem it advisable to attempt to open the trachea lower down. The incision in the skin and superficial fascia of the neck is made in the same manner as in the operation of laryngotomy, but is carried a little further downward. It may be necessary to displace the isthmus of the thyroid gland downward to expose the upper portion of the trachea, and when the trachea is exposed the incision should be made through this and the cricoid cartilage from below upward.

This operation is more often performed in the high operation of tracheotomy than is generally supposed. A tracheotomy-tube is introduced through the wound and secured by

tapes tied around the neck, and the care of the tube should be similar to that in cases of tracheotomy.

Intubation of the Larynx.

This procedure, at the present time, is widely employed as a substitute for tracheotomy in the treatment of the dyspnœa due to inflammatory affections of the larynx or trachea, or stenosis of the larynx; it consists in the introduction of a metallic tube into the larynx, which is allowed to remain in place for a few days. The operation has been recently reintroduced to the profession by Dr. O'Dwyer, of New York, who has devised a set of ingenious instruments for the purpose of laryngeal intubation.

Fig. 206.

Mouth-gag.

The instruments required are a mouth-gag (Fig. 206), with which the jaws are separated and held open; an instrument for the introduction of the tube, which is fastened

Fig. 207.



Intubation-tube and introductor.

to the obturator which fills the cavity of the tube (Fig. 207), and an instrument for extracting the tube after it has been placed in the larynx. (Fig. 208.) The tubes are of metal

and have a collar which rests upon the false cords and bulge slightly toward their middle and again taper toward their lower extremity; at the collar of the tube there is a perforation through which a strand of silk is passed which is made into a loop; this is used to allow the operator to remove the

FIG. 208.

Intubation-tube extractor.

tube if on its introduction it is found to have passed into the œsophagus instead of the larynx, and also is used to remove the tube if it becomes occluded with membrane while in the larynx. The intubation set now in common use is provided with six tubes ranging in size from such as are suited for a child of one year or less up to the age of twelve or fourteen years. (Fig. 209.)

FIG. 209.

Scale of intubation-tubes.

In performing the operation of intubation of the larynx the child is placed upon the lap of the nurse or assistant wrapped in a blanket and the arms are secured by the nurse holding the elbows so as not to interfere with the respiratory movements.

The patient's head is next secured by an assistant, and the position of the head, neck and body, should be that as if it hung from the top of the head, and this position should be firmly maintained during the insertion of the tube. The mouth-gag is next inserted upon the left side and the blades dilated so as to open the jaws widely, and as the gag is self-retaining this position is easily maintained. The jaws being thus held open, the operator,

sitting on a chair facing the patient, next introduces the index finger of the left hand into the mouth and passes it over the tongue until he feels the epiglottis; the introducing instrument to which the tube is attached is held in the right hand and this is now introduced into the mouth, first seeing that the silken loop is free, and it is swept over the tongue and passed down until it touches the epiglottis; this is hooked up by the index finger of the left hand and the tube is passed into the larynx; the index finger of the left hand is then transferred to the edge of the tube and by drawing upon the trigger of the instrument with the index finger of the right hand the obturator is detached, and the instrument is withdrawn, and before removing the finger it is well to place it upon the head of the tube and to sink it well into the larynx. As soon as the obturator is removed there is usually a violent expiratory effort which is accompanied by a gush of mucus, muco-purulent matter or membrane from the tube, and after this escapes the breathing is usually satisfactorily established. If the operator has passed the tube into the œsophagus and has detached it from the introducing instrument, and no improvement in the respiration takes place, it should be withdrawn by the silk loop and attached to the obturator and another attempt should be made to introduce it into the larynx.

The mistake which inexperienced operators make in introducing the tube is in not hugging the posterior surface of tongue closely, so that they pass the tube over the epiglottis into the œsophagus.

The silken loop may be brought out at one side of the mouth and fastened around the ear or fastened to the side of the face by strips of adhesive plaster for a few hours, so that by drawing upon it the nurse or attendant is able to withdraw the tube instantly if it should become obstructed with membrane; or, if it is coughed up, by this means it may be withdrawn from the œsophagus if it has not been expelled from the mouth. Some operators keep the loop attached to the tube during the time it is retained in the larynx, others prefer to remove it after several hours and remove the tube by means of the extracting instrument when

required. The tube is removed at the end of the second or third day and if the child is able to breathe comfortably for an hour or two it is not reintroduced; if, however, the dyspnœa returns it is reintroduced and allowed to remain one or two days longer; several attempts may have to be made before the tube is permanently removed, but it is usually dispensed with from the third to the eighth day.

The most serious complication which is apt to occur during the introduction of the intubation-tube is the detachment and pushing of a mass of membrane in front of the tube into the trachea; if this is too large to be expelled through the tube the breathing is suddenly arrested, and the tube should be removed, and if the mass of membrane does not escape upon the expiratory efforts of the patient the trachea should be rapidly opened as the only means of re-establishing the respiratory function. So much do I dread this accident, which has occurred in a few cases, that I never introduce an intubation-tube without having at hand the necessary instruments to do a tracheotomy if it should be suddenly required, and if possible obtain the consent of the parents or friends to perform tracheotomy if it should be indicated.

One of the greatest troubles after intubation of the larynx is the satisfactory feeding of the patient; liquids as a rule are not swallowed well, a portion of them escaping into the tube, causing coughing and difficulty in breathing. The diet I usually order is of semi-solids, such as corn-starch, soft-boiled eggs, and mush; and if these are not well swallowed it may be necessary to resort to nutritious enemata or the use of a stomach-tube to introduce food. Some patients swallow liquids and semi-solids quite well if the head is dropped a little lower than the body during the act of deglutition.

PART III.

FRACTURES.

In the following article the author has endeavored to confine himself simply to a description of the varieties of fracture and to their dressing and treatment, and he has tried as far as possible to avoid the multiplication of dressings, being satisfied to describe a few of the methods of dressing most frequently employed. He has also avoided the description of complicated splints and dressings, by the use of which in certain fractures most excellent results are obtained, but has preferred to recommend the employment of simple splints and dressings, which can be obtained by physicians practising in districts remote from large cities, where the services of an instrument-maker cannot be obtained to construct special apparatus for the treatment of these injuries.

Varieties of Fractures.

A complete fracture is one in which the line of separation completely traverses the bone, involving the entire thickness of the bone.

An incomplete fracture is one in which there is only a partial separation of the bone-fibres (Fig. 210), under which name are included *partial* or "*green-stick*" *fracture*, in which some of the bone-fibres have given way, while the remaining fibres have been bent by the force and have not been broken. (Fig. 211.) Fissured, punctured, indented,

and perforating fractures are also included in the class of incomplete fractures. (Fig. 212.)

A simple or closed fracture is a fracture in which there are but two fragments, and the seat of injury in the bone does

Fig. 210.

Incomplete fracture of femur.

Fig. 211.

Partial or green-stick fracture of radius.

Fig. 212.

Fissured fracture of humerus. (Gurlt.)

not communicate with the external air by a wound in the soft parts.

Compound or open fractures are fractures in which the seat of injury in the bones communicates with the external air by a wound in the soft parts.

Comminuted fractures are those in which there are more than two fragments, the lines of fracture intercommunicating with each other. (Fig. 213.)

A multiple fracture is one in which a bone is the seat of two or more distinct fractures at different points, the lines of fracture not necessarily communicating with each other.

Complicated fractures are such as are accompanied by some serious injury of the parts in the region of the fracture—as, for instance, the laceration of important blood-vessels or nerves, contusion or laceration of the muscles, or dislocation of a neighboring joint.

FIG. 213.

Comminuted fracture of patella.

FIG. 214.

Impacted fracture.

FIG. 215.

Transverse fracture of femur. (GURLT.)

Impacted fractures are those in which one fragment is driven into and fixed in the other, the impaction taking place at the time of fracture, or being caused by force subsequently applied. (Fig. 214.)

Direction of Fracture.

A transverse fracture is one in which the general line of division of the bone is at right angles with the long axis of the bone. (Fig. 215.) Transverse fractures of the long bones are rarely met with, the line of fracture usually being more or less oblique.

Fig. 216.

Oblique fracture of humerus. (Stimson.)

Fig. 217.

Longitudinal fracture of tibia. (Stimson.)

An oblique fracture is one in which the line of separation is oblique to the long axis of the bone. This is one of the most common directions of the line of fracture. (Fig. 216.)

A longitudinal fracture is one in which the line of separation runs in the general direction of the long axis of the

bone. (Fig. 217.) This form of fracture is rare, but is sometimes met with in the long bones as the result of gunshot injury.

Epiphyseal fracture or separation occurs before complete ossification has taken place between epiphysis and diaphysis, and is rarely seen after the twentieth year of life; the direction of the epiphyseal separation is transverse. (Fig. 218.)

Fig. 218.

Epiphyseal fracture of head of humerus. (Moore.)

The *deformity* or displacement in fractures is either angular, transverse, longitudinal, or rotary.

Repair of Fractures.

The process of repair in cases of fracture is concisely stated by Ashhurst as follows: "The traumatic irritation propagated from the broken bone causes swelling of the periosteum, active proliferation, and formation of a sheath of new bone around the seat of fracture; this is the *ensheathing* or *ring callus* of surgical writers. At the

same time, the medulla feels the effect of the irritation, becomes hardened, and partially ossified; this constitutes the *interior* or *pin callus*. Lastly, the osseous tissue itself undergoes cell-proliferation, and union of the fragments takes place—*mutatis mutandis*—precisely by the same process that we have already studied in considering wounds of the soft tissues. The new material which is thus developed between the fragments themselves, constitutes what Dupuytren called the *intermediate*, *permanent*, or *definitive callus*, in contradistinction to the *ensheathing* and *interior* forms of callus, which are *temporary* or *provisional*."

Examination of Cases of Fracture.

In examining a case of fracture to locate the nature and seat of fracture, the clothing should be removed from the part with as little disturbance as possible, and it is better, in most cases, to cut or rip the clothing, rather than to attempt to remove it in the ordinary manner. The surgeon should first inspect the injured part, and, where possible, compare it with its fellow, as in the case of injuries of the extremities; much valuable information is also derived from the patient or his friends as to the manner in which the injury was produced. The part should next be carefully examined by the surgeon; if it be one of the extremities which is injured, it should be gently lifted, firm extension being made at the same time, the surgeon by his touch and by gentle movements seeking to locate the seat of fracture; and he may, by this manipulation, at the same time develop crepitus.

All manipulations should be made with care, and with the greatest gentleness, not only to save the patient from pain, but also to prevent the soft parts in the region of the fracture from being injured by the rough or sharp fragments of the bone. Rough handling of fractures may increase the muscular spasm by the irritation caused by the sharp fragments of the bones, and may also result in the injury of important vessels and nerves, and indeed a simple fracture may be converted into a compound one by forcible and injudicious manipulations.

The sooner the examination is made after the fracture has occurred the better, for at this time there is less swelling in the region of the injury, and the surgeon can locate the bony prominences with much more ease, and can often discover the exact seat of the fracture with the least amount of manipulation of the parts. When a case of suspected fracture is not subjected to examination for several days after the reception of the injury, the parts in the region of the supposed fracture are often so much swollen that it is impossible to accurately locate its seat, and in such a case it is often necessary to wait until the swelling has subsided before the position of the fracture can be satisfactorily fixed, the case being treated in the meantime as one of fracture.

Anæsthetics may be employed to relieve the patient from pain and to obliterate muscular spasm in the examination of fractures, and their employment is often of the greatest service in the diagnosis of obscure or complicated fractures, especially those in the neighborhood of joints ; but the surgeon should remember that all manipulations should be made with the same gentleness as when the examination is conducted without anæsthesia, for there is the same risk of injury to the surrounding structures by the fragments; this precaution is often neglected when an anæsthetic has been given, the surgeon often being inclined to handle the parts more roughly than he otherwise would ; such practice cannot be too severely condemned.

Provisional Dressings in Cases of Fracture.

It generally happens that fractures occur at localities more or less distant from the point where the treatment of the fracture is to be conducted, and the transportation of the patient and the temporary dressing of the fracture are, therefore, matters of the first importance. In fractures of the upper extremities, if the fracture be simple, the clothing need not be removed, and the arm should be bound to the side by some article of clothing, or supported in a sling made from handkerchiefs or the clothing, and the patient can usually

walk or ride for a short distance without much injury to the parts in the region of the fracture or inconvenience to himself. When the bones of the lower extremities or the trunk are the parts involved, the transportation of the patient is a matter of more difficulty. When the bones of the trunk are involved, the part should be surrounded by a binder firmly pinned or tied, made from the clothing or from towels, or sheets or other strong materials which are at hand. When the bones of the lower extremity are involved, if the fracture be a simple one, the clothing need not be removed, and the motion of the fragments should be prevented by applying to the sides of the limb, extending above and below the seat of fracture, strips of wood, shingles, pasteboard, bundles of straw, strips of bark taken from trees, or bundles of twigs, these being held in place by handkerchiefs or strips torn from the clothing. Umbrellas or canes, or broomsticks (Fig. 219), applied in the same manner, may be employed, the object of any of these dressings being to secure temporary fixation of the fragments of bone during the transportation of the patient.

Fig. 219.



Provisional dressing for fracture of the leg. (Esmarch.)

If the fragments are not fixed in some way, but are allowed to move about during the transportation of the patient, much damage may result to the soft parts surrounding the fractured bones, and simple fractures may become compound ones by the bones being forced through the skin, the discomfort of the patient at the same time being much increased.

Having applied any dressing to bring about fixation of the fragments, the patient should next be placed upon a broad board or settee; if a mattress cannot be obtained, the fractured limb should be laid upon a mass of clothing, or upon some straw, and he should be placed in a wagon or carried to the point where the subsequent treatment of the fracture is to be conducted.

Reduction or Setting of Fractures.

This should be effected as soon as possible after the occurrence of the injury and as soon as the surgeon is prepared to apply the dressings to keep the parts in their proper position; reduction at an early period is less painful to the patient and is accomplished with more ease to the surgeon than at a later period when marked swelling and inflammation are present at the seat of fracture. It consists in bringing the fragments by manipulation as nearly as possible in their normal position, and it is accomplished by extension and manipulation with the hands, care being taken to use as little force as possible to attain the object. Very little force is often required if the surgeon places the part in such a position as to relax the muscles which produce the displacement; when this is accomplished the fragments can usually be pressed into position by the fingers without the application of any considerable force. When the reduction of a fracture has been accomplished the fragments are retained in position by the application of various splints or dressings which serve to prevent their displacement.

Materials and Appliances Used in the Dressing of Fractures.

Fracture Bed.

Many ingenious forms of beds have been devised for the use of patients suffering from fractures of the bones of the trunk and lower extremities, but a simple bedstead provided

with a firm hair mattress having a perforation near its centre, into which is fitted a firm pad, and provided with a pan which slides in a framework beneath a corresponding opening in the bedstead, will prove a useful appliance. The mattress is covered by a sheet perforated to correspond to the opening in the mattress, and when the pad is removed the evacuations of the patient are passed into the pan.

In fractures of the trunk or lower extremities it will be found more convenient in handling the patient to use a single bed not over thirty-two or thirty-six inches in width, and it is not essential that the mattress be perforated, as a bed-pan can usually be slipped under the patient; the mattress should be a firm one stuffed with hair. The use of an ordinary tin pie-plate covered with a piece of old muslin to receive the fecal evacuations may be substituted for the bed-pan and will be found in many cases more satisfactory, especially in the case of children suffering from fracture of the lower extremity.

Splints.

After the reduction or setting of the fragments in cases of fracture they are usually retained in position until union occurs by the use of splints held in position by means of bandages or strips of muslin. Splints may be made of wood, or of tin, lead, copper or wire which possess the requisite amount of firmness and permit of their being moulded to the part, which latter may be found useful in certain cases.

Wooden splints.—The simplest and best splints are made from wood—white pine, willow or poplar being the best material to employ for their construction, being sufficiently strong to give fixation to the parts and at the same time being light. Splints made from smooth white pine, willow or poplar boards from one-eighth to one-half an inch in thickness may be employed in the form of straight or angular splints, and their preparation is a matter of little difficulty.

Wooden splints before being applied to the part should be well padded with cotton, wool, oakum, or hair, and where

lateral wooden splints are employed in the treatment of fractures of the lower extremity it is usual to place bran-bags or junk-bags between the limb and the splint. The carved wooden splints which are sold by the instrument-makers are not to be recommended, as a rule, for unless the surgeon has a large number to select from it is rare that a splint can be obtained to accurately fit any individual case.

Binder's board or *pasteboard* is an excellent material from which to construct splints; it is first soaked in boiling water and when sufficiently soft is padded with cotton or a layer of lint and moulded to the part, and secured in position by a bandage; as it becomes dry it hardens and retains the shape into which it was moulded.

Undressed leather is also an excellent material from which to construct splints; it is applied by first soaking it in boiling water, and after padding it with cotton or lint it is moulded to the part and retained in position by a bandage.

Felt made from wool saturated with gum shellac, pressed into sheets, is also a good material from which to construct splints. This material is prepared for application to the surface by heating it before a fire until it becomes pliable, or by dipping it into boiling water.

Gutta-percha splints made from sheets of this material, in thickness from $\frac{1}{16}$ to $\frac{1}{5}$ of an inch, may often be employed with advantage; it is prepared for use by immersing it in hot water, when it becomes soft and can be moulded to the surface. Care should be taken that it is not allowed to become too soft by too long immersion to permit of its being conveniently handled.

Paper splints made from layers of manilla paper stiffened with starch constitute a very fair substitute for some of the varieties of splints previously mentioned.

Plaster-of-Paris, *starch*, *chalk and gum*, *silicate of potassium* or *sodium* may be employed for the construction of splints, either movable or immovable, in the treatment of fractures; their method of preparation and application is

described (p. 84 *et seq.*); the plaster-of-Paris dressing is the one which is most generally used at the present time.

Fracture-box.—This is a form of splint used in the treatment of fractures of the lower extremity, and consists of a piece of board eighteen to twenty inches in length, with a foot-board firmly secured at its lower extremity; the sides are secured by hinges which allow them to be raised or lowered (Fig. 220). A fracture-box of greater length is required for the treatment of fractures about the knee-joint.

FIG. 220.

Fracture-box with movable sides.

Bran, Sand, or Junk Bags.

These are constructed by taking a piece of unbleached muslin five feet in length and fourteen and a half inches in width, doubling it and securing the free margins except at the mouth by stitches so as to form a bag; the bag is then inverted so that the edges of the seams are brought in the inner surface of the bag. The bags are next filled with dry sand, bran, or hair, or with straw, and the mouth of the bag is closed by stitches or by being tied with a string. Bran bags with splints or sand bags are frequently employed in the treatment of fractures of the femur.

Bandages made of muslin are used to retain splints in the treatment of fractures, and are also sometimes applied directly to the injured part before the application of splints to control muscular spasm and limit the amount of swelling; when a bandage is so used it is known as a *primary roller*. The use of the *primary roller* is sometimes of the greatest service in the dressing of fractures; but its use in inexperienced hands has often been followed by such unfortunate results in the early treatment of fracture, or in cases which are not under constant observation, that I think it is a safe rule of practice to discard entirely the use of the primary roller.

Compresses made from a number of folds of lint, of cotton or oakum, are often employed to retain fragments in position or to make localized pressure upon certain points in the treatment of fractures. The compresses are held in position by strips of adhesive plaster, by a few turns of a roller bandage, or by the splints. Compresses are sometimes employed to protect bony prominences of the skeleton from the pressure of the splints; but this purpose is often better effected by the use of small pieces of soap plaster spread on chamois fitted over the prominent points.

FIG. 221.

Rack for supporting bed-clothes in fracture of the lower extremity.

A rack or cradle, made of wire or wooden hoops, is often employed to support the weight of the bed-clothes in the treatment of fracture of the lower extremity (Fig. 221).

Dressing of Special Fractures.

FRACTURE OF THE NASAL BONES.

Fractures of the nasal bones are often accompanied with fractures involving the septum, the nasal process of the maxillary bone, and the nasal spine of the frontal bone.

The treatment consists in replacing the fragments, if displacement exists, by manipulation with the fingers over the seat of fracture and by pressure made from within the nostrils by a probe or a steel director. When the displacement is once corrected it is not apt to recur, and in the majority of cases no dressing is required. Before resorting to any manipulation within the nasal cavities the mucous membrane should be thoroughly cocainized to render the

operation painless to the patient. When there is depression of the fragments or displacement of the septum, after correcting the deformity by raising the depressed fragment or bending the septum into place by a director, the parts may be held in position by packing the nasal cavity firmly with a strip of antiseptic gauze.

In lateral displacements of the nasal bones from fracture, after reducing the displacement, a small compress held over the fragment by strips of adhesive plaster will be the only dressing required.

Mason transfixes the nose, after reduction of the fragments, with a stout needle, and steadies the pieces with a

FIG. 222.

Mason's dressing for fractures of nasal bones.

strip of plaster crossing the bridge of the nose and fastened to the ends of the needle. The needle is kept in position for eight or ten days (Fig. 222). Roberts, in cases in which there is a displacement of the cartilaginous portion of the

nose, after reducing the deformity, holds the parts in position by transfixing them with steel pins.

Profuse hemorrhage sometimes occurs after fracture of the nasal bones and may require plugging of the nares to control it.

Fractures of the nasal bones are usually firmly united in from ten to twelve days, and dressings may be dispensed with after this time.

Fractures of the Malar Bone and Zygoma.

These fractures are usually the result of direct force; the displacement is upward or backward, and when the zygomatic arch is broken the fragments from pressure upon the masseter muscle or on the tendon of the temporal muscle may interfere with the movement of the lower jaw in mastication. This displacement is corrected by cutting down upon the fragment and elevating it or by passing a tenaculum into the fragment and raising it.

Outward displacements may be corrected by pressure and the application of a compress. The dressing of these fractures after the correction of the deformity, consists in the application of a compress of lint over the seat of fracture, held in position by strips of adhesive plaster or a bandage. There is little tendency to recurrence of the deformity after it has been corrected, and union at the seat of fracture is usually firm at the end of three weeks.

Fractures of the Upper Maxilla.

These fractures may involve the body, the nasal processes or the alveolar processes. The displacement should be corrected and if any teeth have been displaced they should be replaced; if there is comminution of the alveolus the teeth in the separate fragments may be fastened together by fine wire to fix the fragments and hold them in place; and the teeth of the lower jaw should be brought up in contact with those of the upper jaw, and the jaws should be secured together by the application of a Barton's or a Gibson's

bandage (Fig. 223). Inter-dental splints, made of cork with grooves to fit the teeth, or of gutta-percha, are also employed in the dressing of these fractures. The patient should not be allowed to move the jaw in mastication, and

FIG. 223.

Dressing for fracture of the upper jaw.

should be nourished by liquid and semi-solid food which can be taken without removing any teeth to give space for its introduction.

The bandage should be removed every second or third day, and after the face and neck have been sponged off with alcohol it should be reapplied.

These fractures are usually firmly united at the end of four or five weeks, and dressings may be dispensed with at this time.

FRACTURES OF THE LOWER MAXILLA.

The lower jaw may be broken at or near the symphysis, the most usual seat of fracture being near the mental foramen; it is often broken at two places at once and the fractures are in many cases rendered compound by laceration of the mucous membrane, or the injury may consist in a separation of a portion of the alveolar process of the bone. The

dressing of a fracture of the lower jaw, after reducing the displacement and replacing any loosened or detached teeth,

FIG. 224.

Dressing for fracture of the lower jaw.

FIG. 225.

Four-tailed bandage applied for fracture of the lower jaw.

consists in applying a pad of lint under the chin and bringing the jaw up against the upper jaw and holding the com-

press in place and securing the jaws firmly in contact by applying a Barton (Fig. 224), modified Barton or Gibson's bandage. The bandage should be removed and reapplied at the end of the second or third day, and at like intervals during the course of treatment. The patient should be fed upon a liquid or semi-solid diet, not being allowed to chew any solid food until the union at the seat of fracture has become firm.

A very satisfactory temporary dressing for fracture of the lower jaw consists in the application of a four-tailed sling. (Fig. 225.)

Some surgeons prefer to use an external splint moulded from pasteboard or gutta-percha fitted to the chin in the

FIG. 226.



Shape of splint before being fitted to chin.

dressing of this fracture, this being padded with cotton and held in place by a Barton or Gibson bandage. (Fig. 227.) Where there is much difficulty in keeping the fragments in position the wiring together of the teeth may be employed, or the fragments may be perforated with a drill and held in place by a strong silver-wire suture; inter-dental splints of metal or gutta-percha are also sometimes used for this purpose. During the course of the treatment in fracture of the jaws the mouth often becomes very offensive from the fermentation of the saliva and discharges, and it is well to use frequently a mouth-wash of chlorate of potash, tincture of myrrh, glycerin and water.

FIG. 227.

Splint moulded to fit chin.

The dressings for fracture of the lower jaw are usually applied for four or six weeks, the union usually being quite firm at the end of this time.

Fracture of the Hyoid Bone.

In fracture of the hyoid bone, if displacement exists, its reduction is facilitated by pressure made with the finger in the pharynx.

The treatment consists in enforced quiet and the use of opium if cough is a prominent symptom, and the inflammatory symptoms may require the employment of active local treatment. A dressing may sometimes be employed with advantage, consisting of a splint of pasteboard or leather moulded to the anterior portion of the neck.

Fractures of the Larynx or Trachea.

In fractures of the larynx or trachea where there is little displacement and dyspnœa is not marked, the parts should be supported by the application of compresses of lint held in place by strips of adhesive plaster. If, on the other hand, the respiration is embarrassed or there is free expectoration of blood, tracheotomy should be performed, and if the injury be seated in the larynx the displacement of the fragment may be overcome by manipulation with the finger or a director through the tracheal wound, or the larynx may be packed with a strip of antiseptic gauze to control hemorrhage or hold the fragments in position, the patient in the meantime breathing through a tracheotomy tube secured in the tracheal wound; the packing should be removed in a few days, the tracheotomy-tube being permanently removed as soon as the patient can breathe comfortably through the larynx with the tracheal wound closed. In fractures of the trachea the opening into the trachea should be below or at the seat of injury.

Fractures of the Trunk

Fractures of the Ribs.

Fractures of the ribs are more frequent than fractures of any other bones of the trunk; the ribs most commonly broken are those from the fourth to the tenth; the most common seat of fracture is near the junction of the costal cartilages or at the angle. The dressing of fractures of the ribs is best accomplished by enveloping the side of the chest on which the rib or ribs are broken with broad straps of adhesive plaster. The adhesive straps should be two and a half inches in width and long enough to extend from the spine to the middle of the sternum. The straps are warmed and the first strap is firmly applied a short distance below the seat of fracture, extending from the spine to the mid-sternal line; a number of ascending straps are applied in this way, each strap overlapping the preceding one by about one-third of its width, until half the chest is covered in. (Fig. 228.) This dressing usually gives the patient much comfort, and the straps need not be renewed until they become slightly loosened, usually at the end of a week or ten days; they should then be renewed in the same manner.

Fig. 228.



Adhesive plaster dressing for fracture of the ribs.

The dressings for fractures of the ribs are usually dispensed with at the end of three or four weeks, as repair of the fracture is generally well advanced by this time.

A satisfactory temporary dressing for fractures of the ribs consists in surrounding the chest by a broad binder of stout linen or muslin; indeed, some surgeons prefer to employ this dressing during the course of treatment, but as a rule I think it is not as good a dressing as the adhesive

plaster dressing, as the former confines the movements of both sides of the chest.

Fractures of the Costal Cartilages.

These fractures often take place at the junction of the cartilages with the ribs or in the body of the cartilages, and the union of the fracture usually takes place by the production of a mass of bone at the seat of fracture. The dressing for fractures of the costal cartilages consists in the application of strips of adhesive plaster applied in the same manner as for fracture of the ribs, and the dressing should be retained for about the same time.

Fractures of the Sternum.

Fractures of the sternum are rare injuries, but diastasis of the bones of the sternum is a more common accident. The dressing for either variety of injury is the same, and

Fig. 229.

Adhesive plaster dressing for fracture of the sternum.

consists in the application of a compress over the seat of fracture held in place by a broad bandage, or, better, by strips of adhesive plaster (Fig. 229), applied so as to cover and fix the anterior portion of the chest, covering the entire length of the sternum. This dressing should be retained

for at least four weeks, being renewed if it becomes loose at the end of a week or ten days.

Fractures of the Pelvis.

These fractures are often serious injuries from implication of the pelvic viscera. The reduction of the displacement should first be accomplished as far as possible by external manipulation, together with internal manipulation by the fingers introduced into the rectum, or into vagina in the female. The patient should be placed upon a firm bed on his back, with the knees slightly flexed over a pillow, and the parts should be kept at rest by surrounding the pelvis with broad straps of adhesive plaster or a stout muslin binder, or by a firmly applied padded pelvic belt. The hip-joints should be kept at rest by the application of pasteboard splints or by sand-bags. The dressings should be retained for a period of at least six weeks.

When these fractures are complicated by injury of the pelvic viscera various operative procedures may be required, which will compel the surgeon to modify the method of dressing.

Fractures of the Sacrum and Coccyx.

The dressing of these fractures, after effecting reduction of the fragments as far as possible by pressure within the rectum, and, when the sacrum is involved, the application of broad adhesive straps, or of a padded belt, should be employed, and the patient should be kept at rest in bed. When the *coccyx* only is fractured, after reduction of the displacement, the patient should be confined to bed and the bowels should be kept at rest by the use of opium by suppository. The patient should be kept at rest for three or four weeks, and, in case of fracture of the sacrum, the dressings should be retained for this time.

Fractures of the Vertebræ.

Fractures of the vertebræ are always most serious injuries, not only from the injury of the bones themselves, but also from the damage to the spinal cord, membranes, and nerves, which often accompanies them.

In transporting, or turning in bed, a patient suffering from fracture of the vertebræ, great care should be exercised, for rough or sudden motions might cause a displacement of the fragments which might, by injury of, or pressure upon, the spinal cord, rapidly prove fatal.

In the treatment of fractures of the spine, if the deformity is marked, efforts should be made to reduce it by extension and counter-extension, and the result may be successful, especially if the fracture be associated with a dislocation of the vertebræ. In some cases the use of permanent extension by means of weights attached to the legs, shoulders, and chest by adhesive plaster and bandages has been successful in reducing the deformity.

The patient should be placed upon his back upon a bed with a hair mattress, or better, if it can be obtained, a water-bed, which consists of a rubber mattress filled with water, which distributes the weight of the patient's body evenly over the surface. Whatever form of bed be used, the greatest care should be exercised to keep the patient absolutely clean, and the parts of the body or limbs which are exposed to pressure should be frequently bathed with alcohol or soap liniment, and to distribute the pressure, small pads should be placed under the parts and changed at intervals. These precautions are necessary to prevent, if possible, the formation of extensive bedsores, which are a frequent and troublesome complication of these injuries.

The bowels should be carefully watched, and, if constipation is present, it should be relieved by the use of enemata; and, as it is not desirable to lift the patient to slip a bed-pan under him, the discharges can be received in a flat tin plate pushed under the thighs and buttocks, or on pads of oakum or old muslin.

The care of the bladder is also a matter of the greatest importance; the retention which at first exists should be relieved by the use of a flexible catheter introduced with great gentleness, and when incontinence supervenes the catheter should also be used at intervals; the employment of a soft instrument, if used with care, is not apt to produce any injury to the urethra or bladder.

The employment of a plaster-of-Paris jacket has been followed, in some cases, by good results, and it may be applied early in the case, or it may be used after the patient has been kept in the recumbent posture for some weeks; by its use it is often possible to get the patient out of bed and allow him to sit in a chair.

In fractures involving the cervical vertebræ, care should be exercised in lifting or moving the head, and it is often of advantage in these cases to apply short sand-bags to the sides of the neck and head, to give additional fixation to the parts while the patient is in the recumbent posture, or, if he is allowed to get out of bed, to apply a moulded leather or pasteboard splint to the neck, shoulders, and back of the head for the same purpose.

Trephining of the spine in cases of fracture of the vertebræ, to remedy the displacement and relieve the cord from pressure, has been recommended and employed in some cases, and although the operation under strict antiseptic methods is not attended with much risk, the results obtained up to the present time scarcely seem to warrant its performance.

The course of treatment in cases of fractures of the vertebræ, if the patient does not succumb to the injury in a few days or weeks, often extends over many months, and recovery is often more or less incomplete as regards the function of the parts below the seat of fracture.

Fractures of the Skull.

The treatment of fractures of the skull, whether simple or compound, depends largely upon the nature of the injury and the condition of the cranial contents. In simple frac-

tures unaccompanied with cerebral symptoms no special dressing is required, but in compound fractures where loose fragments are present, these should be removed; and if there is no depression of the fragments, and if no cerebral symptoms are present, the wound should be drained and closed and dressed antiseptically, the dressings being held in place by a recurrent bandage of the head.

The patient should be put to bed and the use of an ice-cap to the head is often of service. The diet should be restricted and calomel and opium and bromide of potassium should be administered; it is well to keep the patient for a few weeks in a quiet and darkened room. Where cerebral symptoms are present, either in simple or compound fractures, and trephining is resorted to, the dressing of the wound is similar, and the same general treatment should be adopted. In all cases of fracture of the skull, whether subjected to operative treatment or not, it is well to keep the patient at rest in bed for three or four weeks, and he should be cautioned to avoid excesses and should not resume active work for some months.

Fractures of the Upper Extremity.

Fractures of the Clavicle.

Fractures of the clavicle may be complete or incomplete, and in the latter variety of injury, the deformity is not usually very marked. The indications for treatment in complete fractures of the clavicle are to relax the sterno-cleido-mastoid muscle, to prevent the weight of the arm on the injured side from dragging down the outer fragment of the clavicle, and by fixing the scapula, to carry the attached external fragment outward and forward. A large number of dressings have been devised and used to accomplish these objects. The treatment of fractures of the clavicle by *position* is accomplished by placing the patient in bed on his back upon a firm mattress with a low pillow under his head, and the arm on the side of injury should be fastened

to the side of the chest by a few circular turns of a bandage passing around the arm and chest; the deformity is usually very satisfactorily reduced upon the patient assuming this position, and after three weeks' rest in this position the union is generally sufficiently firm to allow the patient to get out of bed and be about with the arm bound to the side or carried in a sling or with a Velpeau bandage applied, without any recurrence of the deformity.

A satisfactory *temporary dressing* for fractures of the clavicle consists in the application of a four-tailed bandage; the bandage is made from a piece of muslin two yards in

FIG. 230.

Four-tailed bandage for fracture of clavicle.

length and fourteen inches in width; a hole is cut in its centre about four inches from its margin, to receive the point of the elbow; the bandage is then split into four tails in the line of the hole and to within six inches of it; the body of the bandage should be applied so that the point of the elbow rests in the hole, and a folded towel being placed in the axilla, the lower tails should be carried, one anteriorly, the other posteriorly, diagonally across the chest and back to the neck on the side opposite the seat of fracture and secured; the remaining tails are next carried around the

lower part of the chest and secured so as to fix the arm to the side of the body. (Fig. 230.)

In some cases the deformity is corrected by the application of a posterior figure-of-eight bandage, the forearm on the side of injury being carried in a sling. (Fig. 231.)

Fig. 231.



Posterior figure-of-eight dressing for fracture of the clavicle. (Hamilton.)

Sayre's dressing for fracture of the clavicle consists of two strips of adhesive plaster three and a half inches wide and two yards in length. The first strip is looped around the arm just below the axillary margin, and is pinned or sewed with the loop sufficiently open not to constrict the arm. The arm is then drawn downward and backward until the clavicular portion of the pectoralis major muscle is put sufficiently upon the stretch to overcome the action of the sterno-cleido-mastoid muscle, and in this way draws the sternal fragment of the clavicle down to its place. The strip of plaster is then carried completely around the body and pinned or stitched to itself on the back. (Fig. 232.) The second strip is next applied, commencing upon the front of the shoulder of the sound side; thence it is carried over the top of the shoulder diagonally across the back,

under the elbow, diagonally across the front of the chest to the point of starting, where it is secured by pinning or sewing. A slit is made in this strip to receive the point of the elbow. Before the elbow is secured by the plaster, it should be pressed well forward and inward. (Fig. 233.)

Velpeau's dressing may also be used in the treatment of fractures of the clavicle. (Fig. 234.) A compress may also

FIG. 232.

Sayre's dressing for fracture of the clavicle. First strip applied.

FIG. 233.

Sayre's dressing for fracture of the clavicle. Second strip applied.

be secured by the vertical turns of this bandage over the seat of fracture if needed. The application of the bandage is described (p. 55).

In any form of dressing in which the arm is held against the side of the chest, it is well to apply a folded towel or piece of lint between the arm and chest to prevent the surfaces from becoming excoriated.

A modified form of the Velpeau dressing for fracture of the

clavicle is applied as follows: A soft towel or piece of lint is placed against the side of the body and over the front of the chest, and held in position by a strip of adhesive plaster; the arm is next placed in the Velpeau position, a good-sized pad of lint is next applied over the scapula, and this is held in place by a broad strip of adhesive plaster two and a half inches in width and one and a half yards in length; this strip is continued downward and forward so as to pass over the point of the elbow, and is carried diagonally across the chest to the shoulder of the opposite side and is secured, a slit being cut in it to receive the point of the elbow; a compress of lint is next placed over the seat of fracture and held in place by a strip of adhesive plaster; an additional strip of plaster is next carried from the spine around the arm and chest and secured on the opposite side of the chest; circular turns of a roller bandage are then passed around the chest, including the arm, from below upward until the arm is securely fixed to the body, and the dressing is finished by making one or two turns of the third roller of Desault. (Fig. 235.)

Fig. 234.

Velpeau's dressing for fracture of the clavicle.

In the treatment of fractures of the clavicle in *children* the Velpeau or modified Velpeau dressing will be found to be the most satisfactory dressing to employ, and as these patients are particularly apt to disarrange their dressings it is well to render the dressing additionally secure by applying a few broad strips of adhesive plaster over the turns of the roller bandage, the strips following the turns of the bandage.

The removal of dressings and their reapplication will depend upon the comfort of the patient and the manner in which they keep their position. As a rule in fractures of

the clavicle the dressings are removed at the end of the second or third day, the parts are inspected, and the skin is sponged off with dilute alcohol or whiskey; the dressings are then reapplied, and if they are comfortable and the

FIG. 235.

Modified Velpeau dressing for fracture of the clavicle.

parts are in good position, the dressings are made at less frequent intervals until union is completed at the seat of fracture.

Union in cases of fracture of the clavicle is generally quite firm at the end of four or five weeks, and at this time the dressings may be removed, and the patient should carry the arm of the affected side in a sling for several weeks, and should not undertake any work requiring forcible movements of the arm until eight or ten weeks have elapsed from the receipt of the injury.

The time required for union in fractures of the clavicle in children is somewhat shorter; the dressings may be removed at the end of three weeks.

Fractures of the Scapula.

Fractures of the scapula may involve the body, neck, acromion or coracoid process of the bone. Fractures of this bone are quite rare.

Fracture of the Body of the Scapula.

In dressing this fracture, if deformity is present, it is reduced by manipulation, and compresses of lint are placed above and below the seat of fracture and held in place by adhesive strips; the arm is next fixed to the side of the body by spiral turns of a roller bandage passing around the arm and chest, and the forearm is supported in a sling.

Fracture of the Neck, Acromion or Coracoid Process of the Scapula.

These fractures may be dressed by placing a pad of lint or a folded towel in the axilla and binding the arm to the

Fig. 236.

Velpeau dressing for fracture of the scapula.

body by spiral turns of a roller bandage passing around the arm and chest and supporting the forearm in a sling.

Or these fractures of the scapula may be dressed by first placing a pad of lint or a folded towel in the axilla and then securing the arm in the Velpeau position by the application of a Velpeau's bandage. (Fig. 236.) In fractures of the acromion or coracoid processes the union is usually fibrous. In the treatment of fractures of the scapula the dressing should be retained for about four weeks.

Fractures of the Humerus.

Fractures of the humerus may involve the upper extremity, the shaft or the lower extremity of the bone.

Fractures of the Upper Extremity of the Humerus include fractures of the head and anatomical neck of the bone, fractures through the tuberosities, fractures through the surgical neck of the humerus, and epiphyseal fracture or disjunction of the upper epiphysis of the humerus.

The most satisfactory dressing for all fractures of the humerus above the upper third of the bone is applied as follows: A primary roller should be evenly applied from the tip of the fingers to the seat of the fracture, the arm being flexed at the elbow before the bandage is carried above this point, to prevent the dangerous constriction which might result if the bandage were applied with the arm in the straight position, and it were afterwards flexed at the elbow. A folded towel or a thin pad of lint should next be placed in the axilla and over the outer surface of the chest, to furnish a firm basis of support for the humerus and also to prevent excoriation from the contact of the skin surfaces. A splint of pasteboard, felt or leather (Fig. 237) is next moulded to the shoulder and arm; this should be long enough to extend some distance below the seat of fracture and wide enough to cover in about one-half of the circum-

Fig. 237.

Moulded splint for shoulder and arm.

ference of the arm, and is padded with cotton and fitted to the shoulder and arm. The splint and arm are next secured to the side of the body by spiral turns of a roller bandage including the arm and chest in its turns and applied from the elbow to the top of the shoulder. The forearm is carried in a narrow sling suspended from the neck (Fig. 238). This dressing should be removed at the end of twenty-four

Fig. 238.

Dressing for fracture of the upper extremity of the humerus.

or forty-eight hours, and after the parts have been inspected and sponged over with alcohol, the dressings should be reapplied in the same manner, and if the patient is comfortable they need not be disturbed again for three or four days, subsequent dressings being made at the same intervals. Union in fractures of the upper extremity of the humerus, except in *intra-capsular* fracture, in which bony union is the exception, is usually quite firm at the end of five or six weeks, and the dressings can be dispensed with at this time.

Fractures of the Shaft of the Humerus.

The dressing consists in the application of a primary roller from the tips of the fingers to the seat of fracture; a short

well-padded wooden splint extending from the axilla to a point a little above the internal condyle is next placed on the inner surface of the arm and against the chest; a moulded pasteboard or felt splint, fitted to the shoulder and outer side of the arm and extending a short distance below the seat of fracture, is padded with cotton and applied to the shoulder and arm. The splints are held in position and the arm is secured to the body by spiral turns of a roller bandage carried around the chest and arm, and the forearm is carried in a sling suspended from the neck. The dressing is much the same as that for fracture of the upper part of the humerus, with the addition of the short internal splint.

Fractures of the shaft of the humerus may also be dressed by first applying a primary roller and then placing the forearm and arm upon a well-padded internal angular splint. (Fig. 239.) Care should be taken to see that the end of

FIG. 239.

Internal angular splints.

the splint extends only to the axilla and does not press upon the brachial vein. A pasteboard or felt moulded splint is next applied to the shoulder and outer side of the arm. The splints are held in position by turns of a roller bandage beginning at the fingers and carried up to the shoulder. (Fig. 240.) The arm is supported by a sling applied at the wrist, and sometimes for additional security the arm is secured to the side of the body by spiral turns of a bandage carried around the arm and chest. The after-treatment of

these fractures as regards the removal and renewal of the dressings is the same as in cases of fracture of the upper portion of the humerus.

Fig. 240.

Dressing for fracture of the shaft of the humerus with internal angular splint.

In fractures of the shaft of the humerus the dressings should be retained for five or six weeks.

Fractures of the Lower Extremity of the Humerus.

These include fractures at the base of the condyles, splitting fractures between the condyles or those of the internal or external condyle, and epiphyseal fracture or disjunction of the lower epiphysis of the humerus.

In dressing fractures of the lower extremity of the humerus, if a primary roller is employed it should be carried up only to the elbow; the displacement is reduced by extension and manipulation, and before applying any splint it is well in many cases to apply over the region of the fracture several folds of lint saturated with lead-water

and laudanum, and to cover this dressing with waxed paper or rubber tissue to diminish as far as possible the swelling, which is very marked after these injuries. An anterior angular splint (Fig. 241) well padded with cotton or oakum

FIG. 241.

Anterior angular splint.

is next applied and held in position by the turns of a roller bandage applied from the fingers to the upper portion of the splint. (Fig. 242.) These fractures may also be dressed with a well-padded internal angular splint, this splint being

FIG. 242.

Dressing for fracture of the lower extremity of the humerus with anterior angular splint.

substituted by an anterior angular splint at the end of ten days or two weeks.

These fractures may also be dressed by placing the arm in a posterior angular trough (Fig. 243) made of pasteboard

or leather. Some surgeons prefer to dress fractures of the condyles of the humerus with the arm in the extended position upon a straight anterior splint, or with short, narrow pasteboard splints applied around the joint, as favoring more accurate coaptation of the fragments. If this position is employed a straight wooden splint is applied to the anterior surface of the arm and forearm, or moulded splints of pasteboard may be used, and after the union is moderately firm, at the end of two weeks, the elbow should be flexed and kept in this position during the remaining time of the treatment.

Fig. 243.

Posterior angular trough.

When fractures of the lower extremity of the humerus involve the elbow-joint a certain amount of impairment of joint-motion is apt to occur either from ankylosis or from displacement of the fragments which in many cases it is impossible to completely reduce, so that flexion and extension of the joint is restricted. Bearing these facts in mind, it is well to make passive motion in these cases as early as the second or third week. It is well to explain to the patient or his friends that impairment of joint-motion may result in these fractures in spite of the greatest skill and care in the treatment. In a case of fracture in the region of the condyles of the humerus the dressings should be removed in twenty-four hours and it should be re-dressed in the same manner, and if the swelling does not increase and the dressing is comfortable to the patient it should afterward be dressed at less frequent intervals; the union is generally quite firm at the end of four weeks and the splint may be removed at this time. Fractures of the condyles of the humerus are very common in children and epiphyseal disjunctions of the lower epiphyses are also met with; the dressing of these injuries in this class of patients is similar to that described for fractures of the condyles of the humerus.

Fractures of the Olecranon Process of the Ulna.

Fractures of the olecranon may consist in simply a separation of the cortical layer of bone over the summit of the process to which the triceps is principally attached, or the line of fracture may pass through the sigmoid fossa.

Fractures of the olecranon are dressed with the arm slightly flexed at the elbow, or with it completely extended;

Fig 244.

Adhesive strap applied to draw fragment downward.

the former position is possibly a little less irksome to the patient. The separation of the fragment by the action of

Fig. 245.

Fracture of olecranon dressed in the extended position.

the triceps muscle is usually not very marked; but, if the displacement is marked, it may in a measure be overcome by the use of a compress above the fragment, over which figure-of-eight strips of adhesive plaster are fastened to draw it

down into position (Fig. 244). A primary roller should then be carefully applied to the forearm and arm with figure-of-eight turns at the elbow to reinforce the action of the strips of plaster, and a well-padded straight wooden splint extending from the upper third of the arm to the ends of the fingers is next securely fastened to the arm by the turns of a roller bandage (Fig. 245).

This fracture may also be dressed by first applying a primary roller up to the elbow, and then placing the arm upon a well-padded anterior obtuse-angled splint, or a straight splint with a good-sized pad of lint or oakum fastened at a point corresponding to the position of the flexure of the elbow. When either of these splints is placed upon the arm a position of moderate flexion is obtained. A compress of lint is next placed above the fragment, if there is displacement, and one or two narrow strips of adhesive plaster are fastened to this and passed obliquely downward and attached to the splint on either side. The splint is then securely fastened to the arm by the turns of a roller bandage applied from the fingers to the upper end of the splint.

The dressings in a case of fracture of the olecranon should be removed at the end of twenty-four or thirty-six hours, or sooner if there is evidence of swelling of the tissues in the region of the fracture, and they should be reapplied in the same manner. If the dressing is comfortable to the patient, and there is no evidence of swelling, the subsequent dressings should be made at less frequent intervals ; the dressings are usually retained in this fracture for five or six weeks. Passive motion should not be made until this time, as flexion of the elbow tends to separate the fragments, unless union has taken place. The repair of a fracture of the olecranon is, in most cases, by fibrous union, but in a few instances bony union has been found to have taken place.

Fractures of the Coronoid Process of the Ulna.

Fractures of the coronoid process are rarely met with, and their dressing is accomplished by placing the arm in a flexed

position and applying a well-padded internal right-angled splint, or a posterior right-angled splint, and securing it to the arm by the turns of a roller bandage. A moulded pasteboard or leather gutter may be substituted for the angular splints. The dressings should be changed at intervals, and after their removal at the end of three or four weeks, passive motion should be practised.

Fractures of the Head and Neck of the Radius.

These fractures are also quite rare, and, when met with, should be dressed, after reducing the fragments by manipulation, by flexing the elbow and keeping it in this position by the application of a well-padded anterior right-angled splint, the splint being firmly secured in position by the turns of a roller bandage applied from the tips of the fingers to the upper end of the splint. The splint should be changed at intervals, and should not be permanently removed for four weeks, at which time passive motion, consisting in flexion and extension at the elbow and pronation and supination of the forearm, should be made. (Fig. 242.)

An internal angular splint applied to the inner surface of the forearm and arm may also be used in the treatment of these fractures. (Fig. 240.)

Fractures of Both Bones of the Forearm.

These fractures are often met with as the result of direct or indirect violence, and after reducing the displacement, which is always marked when both bones are broken, and is not so marked when one bone only is broken, by making extension from the hand and by manipulation; the forearm is placed in the supine position or in a position between pronation and supination. The supine position is, as a rule, to be preferred in any fracture of the radius, as the upper fragment is supinated by the action of the biceps and supinator brevis muscles, and, therefore, unless the lower

fragment be placed in the supine position union with rotary deformity will almost inevitably ensue.

Two straight wooden splints, well padded, a little wider than the forearm, are employed. The anterior splint should be long enough to extend from the elbow to the tips of the fingers, and the posterior splint should extend from the elbow to the wrist. A primary roller should never be ap-

FIG. 246.

Dressing for fracture of both bones of the forearm.

plied to the forearm in dressing these fractures, as its application diminishes the interosseous space and its use has been followed by gangrene of the hand and forearm. In applying the anterior splint to the palmar surface of the forearm and hand care should be taken to see that the upper end of the splint does not press upon the brachial artery and vein at the elbow when the forearm is flexed; the posterior splint is next applied from the elbow to the wrist and the splints are held in position by the turns of a bandage carried from the fingers to the elbow. (Fig. 246.)

In fracture either of the shaft of the radius or of the ulna alone, the deformity is usually not so marked as when both bones are broken at the same time, the unbroken bone acting as a splint; the dressing for these fractures is the same as for fracture of both bones of the forearm.

The dressing should be removed in twenty-four or thirty-six hours, and after inspecting the parts and sponging them

with dilute alcohol the splints should be replaced in the same manner and secured, and the dressings should be removed and renewed at intervals of two or three days for two weeks at least, and after this time the dressings should be made at less frequent intervals. The time required for union in these fractures is usually five or six weeks, and the splints should be retained for this time.

Fractures of the forearm should be seen by the surgeon frequently for the first two weeks of the treatment, for it is in these fractures that the most unfortunate results have occurred from neglect of this precaution.

In children *incomplete* or *green-stick* fractures of the bones of the forearm are very common; their dressing, after reducing the deformity, which consists in bending the bones back into place, which often converts the incomplete fracture into a complete one, is accomplished in the same manner as described above. In these patients there is a great tendency to displace the splints or rather to draw the forearm out of the splints, and to prevent this I often employ an anterior angular splint, in place of the straight anterior splint, the upper portion of which, being fastened to the arm, prevents the child from dragging the arm out of the dressings.

Fractures of the Lower End of the Radius.

The most common fracture of the radius is one situated from one-half of an inch to one and one-half inches above the lower articular surface of the bone, the line of fracture being more or less transverse, although it may in some cases be slightly oblique; the characteristic deformity in this fracture is represented in Fig. 247.

The most important point in the treatment of this fracture is to effect complete reduction before the application of any splint; this is done by making extension from the hand, and at the same time, by extending and flexing the wrist and by manipulation, the deformity can usually be completely reduced. The arm should then be brought into the

position of supination, and a firm compress of lint is next placed over the lower end of the upper fragment on the

Fig. 247.



Fracture of the radius near its lower extremity.

Fig. 248.

Position of compresses in Colles's fracture.

Fig. 249.

Bond's splint.

palmar surface of the forearm; a second compress is then placed over the upper end of the lower fragment (Fig. 248), and a well-padded Bond splint (Fig. 249) is applied to the

palmar surface of the arm and held in place by the turns of a roller bandage. (Fig. 250).

Fig. 250.

Dressing for fracture of the lower end of the radius.

Many surgeons treat this fracture with the hand in a position between pronation and supination, the thumb pointing upward. A substitute for Bond's splint may be prepared by fastening a roller bandage obliquely upon a straight wooden splint as suggested by Dr. Hays. (Fig. 251.)

Fig. 251.

Substitute for Bond's splint.

Two straight splints with compresses are also employed in the treatment of this fracture, and a vast number of splints have been devised; among these may be mentioned those of Gordon, Coover, and the metal splint of the late Dr. R. J. Levis. The most important point in the treatment of this fracture is the complete reduction of the deformity at the first dressing, and if this has been satisfac-

torily done almost any splint may be used with a good result, and indeed some surgeons use no splint, applying only a compress over the palmar fragment, held in place by a strip of plaster, the arm being carried in a sling.

The after-treatment of these fractures consists in removing the splint and compresses after twenty-four or thirty-six hours and in sponging the surface of the skin with dilute alcohol, and the compresses and splint should then be reapplied in the same manner; the fracture should be dressed every second or third day for the first two weeks, and after this time it should be dressed at less frequent intervals. Union is usually quite firm at the end of four weeks, and the splint should be dispensed with at this time. A certain amount of stiffness of the wrist and fingers is apt to follow this fracture, which is usually soon overcome by passive motion and physiological use of the parts.

In *children* epiphyseal separations or fractures of the lower epiphysis of the radius are often met with, and their treatment is similar to that described above; a Bond splint with compresses or two straight splints with compresses being the most satisfactory dressing to employ in this injury, the dressings being retained for three weeks.

Fractures of the Carpal Bones.

These fractures are usually compound or open fractures, and are so frequently associated with extensive laceration of the arm and hand that operative measures have to be resorted to; but if such is not the case they are dressed, when compound, with an antiseptic dressing, and the hand and forearm are supported upon a well-padded palmar splint held in place by a roller bandage; more or less impairment in the motion of the wrist is apt to follow these fractures. In simple fractures of the carpal bones the use of an evaporating lotion for a few days, in connection with the splint just mentioned, will be found useful. The dressings should be retained for three or four weeks, and after their removal passive motion should be employed to overcome as far as possible the joint-stiffness resulting.

Fractures of the Metacarpal Bones.

These fractures are often met with as the result of direct or indirect force applied to the metacarpal bones. The treatment of fractures of the metacarpal bones consists in first reducing the deformity, which is usually an angular one, the projection of the angle being toward the back of the hand; this is reduced by pressure with the fingers, and the hand and forearm should then be placed upon a palmar splint (Fig. 252) with a pad of oakum or cotton under the

Fig. 252.



Agnew's splint for fracture of the metacarpal bones.

palm; a compress of lint is next placed over the seat of fracture, and the hand and forearm are bound to the splint by the turns of a roller bandage. (Fig. 253.) At the end

Fig. 253.

Dressing for fracture of the metacarpal bones.

of three weeks union at the seat of fracture is usually quite firm, and the splint should be dispensed with at this time.

Fractures of the Phalanges.

The treatment of fractures of the phalanges consists in reducing the displacement by extension and manipulation,

and in placing the finger in a moulded gutta-percha or pasteboard splint (Fig. 254), and securing the splint in position

Fig. 254.

Gutta-percha splint for fracture of phalanx. (Hamilton.)

by the turns of a roller bandage. When the proximal phalanx is fractured a narrow, padded, wooden splint extending from the end of the finger to the wrist should be applied upon the palmar surface of finger and hand, and a short dorsal splint should also be used; if there is a tendency to

Fig. 255.

Dressing for fracture of phalanx with anterior and posterior splints.

lateral displacement short lateral splints should also be employed, and the splints should be held in place by strips of plaster or by a roller bandage. (Fig. 255.)

Union in fractures of the phalanges is usually quite firm

at the end of three weeks, and the splints can be dispensed with at that time.

Fractures of the Lower Extremity.

Fractures of the Femur.

Fractures of the upper extremity of the femur are those involving the neck, great trochanter, and upper end of the shaft of the bone.

In dressing fractures of the upper extremity of the femur the patient should be placed in bed upon a firm mattress, and an extension apparatus made from adhesive plaster should be applied to the leg, extending as far as the knee-joint. The extension apparatus is constructed by taking a piece of adhesive plaster two and a half inches in width and long enough to extend from the outer side of the knee

Fig. 256.



Adhesive plaster extension apparatus applied to limb. (Ashhurst.)

to four inches below the sole of the foot, and from this point back to the inner side of the knee; in the centre of this strip is placed a block of wood, two and a half inches wide and four inches in length, with a perforation in its centre; the block and the inner surface of the strip on each side are next faced with a similar strip of adhesive plaster to a

point about an inch above each malleolus; a few straps are next wound around the wooden block to fix the previously applied straps; the strip of plaster is next warmed and applied to the sides of the leg and held in position by three strips of adhesive plaster carried around the leg at intervals (Fig. 256), and the plaster is made additionally secure by the application of a roller bandage applied to the foot and leg and carried up to the knee. Through the perforation in the block or stirrup is fastened a cord which passes over a pulley attached to the bed, and to this cord is attached the extending weight. The extension apparatus being applied, lateral support is given to the leg and thigh by sand-bags applied on either side; the outer sand-bag should extend from the foot to the axilla, and the inner one from the foot to the groin. A weight of five or ten pounds is attached to the extending cord, and the lower feet of the bed should be raised on blocks a few inches high to prevent the patient from slipping down in bed; a pad of oakum or cotton should also be placed under the tendo Achillis to relieve the heel from pressure. This dressing is kept in place for from four to six weeks, and if union has occurred the patient is kept in bed for a few weeks longer and is then allowed to be about using crutches. In the majority of cases of fracture of the neck of the femur fibrous union only takes place, and after employing the dressing before described for six weeks the patient is allowed to get up and go about on crutches. It often happens that the subjects in whom these fractures occur are old and feeble, and if it is found that restraint in bed with the dressings here described is not well borne, under such circumstances they should be discarded and the patient should be allowed to sit up in bed with the limb resting on a pillow, or in a chair, the treatment of the local condition having to be disregarded, attention being given to the patient's constitutional condition.

The application of a plaster-of Paris bandage to the leg, thigh and pelvis is also sometimes made use of in the treatment of fractures of the upper extremity of the femur; extension should be made from the foot while the bandage is being applied. (Fig. 257.) In fractures of the neck of

the femur and of the upper part of the shaft of the bone the anterior wire splint of Prof. N. R. Smith is sometimes

FIG. 257.

Plaster-of-Paris bandage applied to thigh. (HAMILTON.)

used with advantage; the limb being swung from the splint the patient is able to move in bed without causing him pain

FIG. 258.

Smith's anterior splint for fracture of the femur.

or disturbing the fragments. (Fig. 258.) In fractures in the upper portion of the femur where there is marked tilting for-

ward of the upper fragment Prof. Agnew employs extension made from the thigh and places the limb upon a double in-

Fig. 259.



Dressing of fracture of the femur with extension upon an inclined plane. (Agnew.)

clined plane and maintains this position during the treatment of the case. (Fig. 259.) With the same object in view, in place of the double inclined plane a double inclined frac-

Fig. 260.

Double inclined fracture-box.

ture-box may be employed, extension being made from the thigh by means of adhesive plaster strips applied above the knee, to which a weight is attached. (Fig. 260.)

Fractures of the Shaft of the Femur.

In the treatment of fractures of the shaft of the femur the dressings are applied to diminish as far as possible the shortening and to prevent angular or rotary displacement

of the fragments. In dressing these fractures the patient should be placed upon a fracture-bed or an ordinary bed with a firm hair mattress; an extension apparatus of adhesive plaster is applied and extension is made by a weight attached to this as previously described. Lateral support is given to the limb by the application of two wooden splints—the outer or long one extending from the axilla to the foot, the inner or short one extending from the groin to the foot. The splints at their upper extremity should be about six inches in width and at their lower extremity about three and a half inches The splints are wrapped in a splint cloth which extends from the foot to the groin, and after this has been placed under the limb the splints are fixed in their proper positions, the short one to the inner side, the long one to the outer side of the limb. Between the limb and the splints are interposed bran-bags; the outer bag should be long enough to extend from the axilla to the foot, the inner one from the groin to the foot. The splints and bran-bags are held in

FIG. 261.



Dressing for fracture of the shaft of the femur with splints and bran-bags. (ASHHURST.)

place by five or six strips of bandage passing under the limb and body and around the splints and bran-bags at intervals. The heel is saved from pressure by placing a wad of oakum or cotton under the tendo Achillis and after the splints have been brought into place the strips of ban-

dage are firmly tied to secure them and a weight of ten or twelve pounds is attached to the extending cord. The foot of the bed is raised to prevent the patient from slipping downward and to allow the weight of the body to act as a counter-extending force. After the application of the dressings the thigh should be slightly abducted. During the after-treatment of these fractures the surgeon should see that the splints and bran-bags are kept firmly in place and that the foot does not roll outward; this is accomplished by untying the strips and readjusting the bags and then bringing up the splints and securing them in position by fastening the strips. The extension apparatus usually does not require renewal during the course of treatment. The extension and splints are kept in place for four or six weeks and at this time union at the seat of fracture is usually quite firm, so that they may be removed, and the fracture is then supported by moulded pasteboard splints or by the application of a plaster-of-Paris splint for several weeks longer, and at the end of eight weeks it is safe to allow the patient to be up and around on crutches.

Many surgeons, in fracture of the shaft of the femur, prefer to use a long external sand-bag and a shorter internal one in place of the corresponing long and short splints and bran-bags, and, if care is observed to see that the sand-bags are kept accurately in contact with the limb and body, excellent results may be obtained by this form of dressing. After considerable experience with both methods of furnishing lateral support in the dressing of fractures of the shaft of the femur, I am well satisfied that angular deformity is less likely to result where the splints and bran-bags are employed.

The plaster-of-Paris dressing, including the foot, leg, thigh, and pelvis, is employed by some surgeons in the early treatment of fracture of the shaft of the femur, the limb being kept well extended until the plaster has thoroughly set. The double inclined plane and the anterior angular wire splint are also sometimes employed in the dressing of fractures of the shaft of the femur.

Fractures of the Shaft of the Femur in Children.

The treatment of these fractures in young children by extension by a weight and pulley and lateral splints is often unsatisfactory on account of the difficulty in keeping the patient quiet upon his back, and from the soiling of the dressings by the feces and the urine. In children two years of age and over I have never found much trouble in employing extension and lateral support by splints and bran-bags or sand-bags, and in these cases I make additional fixation at the seat of fracture, and guard against displacement of the fragments by the child sitting up in bed when not watched, by carefully moulding external and internal pasteboard or felt splints to the thigh, and holding them in place by the turns of a bandage. I have employed this form of dressing even in children under two years of age with the most satisfactory results.

In cases of fracture of the femur in children from a few months to a year or eighteen months of age, in whom it is difficult to obtain quietude, or who have to be moved to give them nourishment if they are taking the breast, the dressing which I have found most satisfactory consists in first applying a roller bandage from the foot to the groin, and then moulding to the outer half of the foot, leg, thigh, and also to half of the pelvis, a pasteboard or felt splint which is well padded with cotton, and held in position by the turns of a bandage carried from the foot to the pelvis and finished with circular turns about the pelvis. The splint should be so moulded as to include a little more than one-half of the circumference of the thigh and leg. If this splint becomes soiled it is easily replaced by a fresh one, and its removal and renewal is much easier than that of the plaster-of-Paris

Fig. 262.



Fracture of the femur treated by vertical extension.

splint which is recommended by some surgeons in these cases.

In young children fractures of the femur are often *incomplete* or *green-stick fractures;* and, even when complete, the shortening is usually not marked, as the line of fracture is apt to be transverse, the periosteum often not being completely ruptured, which tends to hold the fragments in position.

In green-stick fractures the deformity should be reduced by manipulation, even if it is necessary to convert the incomplete fracture into a complete one to accomplish this object.

Mr. Bryant recommends that fractures of the femur in young children be treated in the vertical position; the injured limb, together with the sound one, is flexed at a right angle to the pelvis and fixed with a light splint, and attached to a cradle or bar above the bed. (Fig. 262.)

If the plaster-of-Paris dressing is used, the limb should be first enveloped from the foot to the pelvis with a flannel bandage, and extension should be made while the plaster-of-Paris bandage is being applied and should be kept up until the bandage has become fixed. The plaster bandage should extend from the toes to the pelvis, and it is well to fix the hip-joint by carrying several turns of the bandage about the pelvis. To prevent the splint from absorbing the discharges and becoming offensive, the upper portion of it may be coated with shellac.

The time required for union in fractures of the femur in children is about three weeks, and the dressings may be removed at this time, but the child should not be allowed to use the limb for several weeks after this period.

Fractures of the Lower End of the Femur.

The fractures met with in this portion of the femur are supra-condyloid fractures, or those in which one condyle is separated, or comminuted fractures in which both condyles are separated; epiphyseal disjunctions of the lower end of the femur, met with in young subjects, may also be classed with fractures at this portion of the bone.

The dressing of supra-condyloid fractures, if there is short-

ening, should be similar to that employed in fractures of the shaft of the femur, consisting in the application of an extension apparatus, and bran-bags and splints or sand-bags to give lateral support; if, however, there is no marked shortening the dressing employed should be the same as that applied in fractures involving one or both condyles or epiphyseal separations.

The dressing employed in fracture of one or both condyles or in epiphyseal disjunction of the lower end of the femur consists in placing the limb in a long fracture-box extending from the foot to the upper third of the thigh, the box being well padded with a soft pillow, or a well-padded posterior splint, or a moulded pasteboard or felt gutter may be employed; if either of these dressings is employed, the splint or gutter should be long enough to extend from the lower part of the leg to the middle of the thigh.

If there is much effusion into the joint or soft parts, lead-water and laudanum should be applied over the region of injury for some days, until the swelling has subsided. At the end of two weeks it is well to place the limb in a plaster-of-Paris dressing, extending from the foot to the middle of the thigh. This dressing should be retained for four weeks, and at the end of this time the dressing should be removed, and if the union is sufficiently firm to allow the patient to go about on crutches, a fresh plaster-of-Paris splint should be applied extending from the middle of the leg to the middle of the thigh, or lateral splints of pasteboard may be substituted for the plaster dressing.

A certain amount of permanent impairment of the joint motion is apt to follow fractures involving one condyle or both condyles of the femur.

Fractures of the Patella.

The dressing of fractures of the patella consists, first, in the application of a roller bandage from the toes to the upper part of the leg; a well-padded posterior wooden splint long enough to extend from the middle of the leg to the

middle of the thigh, or an Agnew splint, which is provided with pegs for the attachment of strips of adhesive plaster (Fig. 263) is next placed under the limb. A small compress of lint is next placed above the upper fragment, and a

Fig. 263.



Agnew's splint for fracture of the patella.

similar compress is placed below the lower fragment; a strip of adhesive plaster one and a half inches in width and twenty-four inches in length has its middle portion applied over the compress, and its ends are then brought obliquely downward and fastened to the splint, or to the pegs if

Fig. 264.



Agnew's splint applied.

Agnew's splint be used; this may be reinforced by a second or third strip. The object of these strips is to bring the upper fragment down in contact with the lower fragment. A strip of plaster with the ends passing in the opposite direction is next placed over the lower compress, and the ends are fastened to the splint or pegs; this strip serves

only to steady the lower fragment, as it cannot be drawn upward to meet the upper fragment by reason of the inextensibility of its ligamentous attachment. (Fig. 264.) If the Agnew splint is employed the strips of plaster may be tightened by turning the pegs to which they are fastened without removing the splint.

The splint is next firmly fixed in contact with the limb by the turns of a roller bandage extending from the lower to the upper end of the splint. The limb should next be placed upon an inclined plane or in a long fracture-box with its foot elevated to relax the quadriceps femoris muscle. This dressing should be removed and reapplied in a few days, as the dressings become loose as the swelling about the seat of injury subsides, and after this disappears the dressings require renewal at less frequent intervals and usually at the end of three weeks the splint may be removed and a plaster-of-Paris bandage may be applied extending from the middle of the leg to the middle of the thigh. At the end of six weeks the patient may be allowed to walk upon the limb, the knee-joint being fixed with a plaster-of-Paris or pasteboard splint.

It is well, after the removal of the splints, for the patient to wear for some months a laced muslin knee-supporter, which gives some support to the knee-joint.

The union in fractures of the patella is usually fibrous, although in rare cases bony union has occurred.

A great variety of splints have been devised and used in the treatment of fractures of the patella, the main object of which is to fix the knee-joint and bring the fragments as nearly as possible in apposition. Malgaigne's hooks or Levis's modification of the same are employed by some surgeons to secure close apposition of the fragments. The method of treatment in fractures of the patella, which consists in exposing the fragments by an incision and drilling and suturing them with catgut or silver wire sutures, is also employed at the present time, the strictest antiseptic precautions being taken to prevent infection of the wound.

In cases of rupture of the fibrous union after fracture of the patella, which is not an uncommon accident, the treat-

LANE LIBRARY, STANFORD UNIVERSITY

ment of the case should be the same as that for a recent fracture of the patella.

Fracture of the Bones of the Leg.

In fractures of both bones of the leg the displacement is usually very marked; when one bone only is broken, the sound bone, acting as a splint, prevents much deformity, except in case of fracture at the lower end of the fibula, when the foot inclines to the injured side.

The dressing for fractures of both bones of the leg or for fracture of the tibia or fibula alone, except in cases where the lower portion of the fibula is the seat of injury, is best accomplished by the use of a fracture-box. (Fig. 265.)

Fig. 265.

Fracture-box with movable sides.

The displacement being overcome as far as possible by extension and manipulation, the leg is placed in a fracture-box, which is prepared for the reception of the limb by having the sides let down and having a soft pillow laid in it; the foot is next secured to the footboard by a loop of bandage passed around the foot, the ends being tied after passing through the slots in the footboard; a pad of oakum or cotton is placed under the tendo Achillis to relieve the heel from pressure, and a similar pad is placed between the sole of the foot and the footboard. (Fig. 266.) The sides of the box are then brought up and secured by two or three strips of bandage tied around the box. In using a fracture-box in the treatment of fractures of the bones of the leg the surgeon should see that the foot is kept well down to the footboard and is at a right angle with the leg, that there is no eversion of the knee and that the pillow is

full enough to make equable pressure upon the leg when the sides of the box are secured, and that the heel is not subjected to undue pressure—the use of a pad of oakum or

FIG. 266.

Application of the fracture-box.

cotton under the tendo Achillis being employed to prevent this complication. Where there is a tendency to tilting upward of the lower end of the upper fragment the lower fragment can be brought in line with this by raising the foot by a mass of oakum or cotton placed under the tendo Achillis and heel and so overcoming the deformity.

FIG. 267.

Plaster bandage applied to fracture of the leg.

The subsequent dressings of the cases are conducted by letting down the sides of the box and correcting any displacement, if present, by adjusting the limb and pads in their proper position, and again bringing up the sides of the

box and securing them. At the end of two or three weeks the fracture-box may be removed and a plaster-of-Paris dressing may be applied to the limb, which will allow the patient more freedom of movement in bed, or permit of his sitting up without disturbing the fragments (Fig. 267).

Union in fractures of the bones of the leg is usually quite firm in six weeks, but the patient should not be allowed to put his weight upon the limb in walking for at least eight weeks.

If the patient is restless, and finds his position with the fracture-box resting upon the bed irksome, the fracture-box may be swung from a frame fastened over the bed (Fig. 268).

Fig. 268.

Fracture-box suspended. (Agnew.)

The application of a plaster-of-Paris dressing as a primary dressing—the ordinary plaster-of-Paris bandage or the Bavarian dressings being applied—in fractures of the bones

FIG. 269.

Moulded binder's board splints for fractures of the leg.

of the leg, is adopted by some surgeons, and, if employed, the case should be under constant supervision for a few days, so that the dressing can be removed if a dangerous amount of swelling takes place. Moulded splints of felt or pasteboard are also sometimes applied in the treatment of these cases. (Fig. 269.)

In patients suffering with delirium tremens, or in maniacal patients, the use of a fracture-box in the treatment of fractures of the bones of the leg is often not satisfactory on account of the difficulty in restraining the movements of the patient,

and the consequent displacement of the fragments. In such cases it is well to apply a few strips of binder's board, well padded with cotton, to the limb, extending above and below the seat of the fracture, holding them in place by a few turns of a roller, and then to wrap the limb and foot in a soft pillow and hold this in place by the turns of a roller bandage applied with moderate firmness. This dressing allows the patient to move the limb without serious disturbance of the fragments, and, after the patient recovers from his attack, the leg may be placed in the fracture-box.

In fractures of the bones of the leg *in young children* the same difficulty is often experienced in keeping them quiet, and for this reason a fracture-box cannot be used with satisfaction. In dressing these cases, two lateral splints of pasteboard, moulded to the foot and leg and well padded with cotton, may often be employed with the best results. The splints should not be wide enough to meet on the anterior or posterior surface of the leg or foot. The splints, after being carefully adjusted, are held in place by the turns of a roller bandage; and, after these splints have been applied for two weeks, and all swelling has subsided at the seat of fracture, a plaster-of-Paris bandage may be substituted for them, which should be worn for three weeks; at the expiration of this time union is usually firm enough to dispense with all dressings.

Fractures of the Fibula.

In fractures of the fibula, with the exception of that fracture occurring at the lower end of the bone, the deformity is not marked, and they are usually dressed with a fracture-box applied as in the dressing of fractures of both bones of the leg, and at the end of two weeks a plaster-of-Paris dressing should be applied, and the patient may be allowed to get out of bed and move about on crutches. The union in a fracture of the fibula is usually quite firm at the end of five weeks, and all dressings may be dispensed with at that time.

Fracture of the Lower End of the Fibula.

This fracture usually occurs in the lower fifth of the bone and is often associated with a laceration of the internal lateral ligament of the ankle-joint or a sprain fracture of the internal malleolus and is usually accompanied by marked eversion of the foot. This fracture is commonly known as *Pott's fracture.*

In this fracture after reducing the displacement by extension and manipulation, the limb should be placed in a fracture-box provided with a soft pillow, the foot should be secured to the footboard and a pad of oakum or cotton should be placed under the tendo Achillis; before bringing up the sides of the box and securing them two firm compresses of lint or oakum should be placed in contact with the leg, one just above the inner malleolus, the other just below the outer malleolus. The sides of the box are next brought up and secured, and by the pressure of these compresses the foot is brought into an inverted position and the deformity is corrected.

The after-dressing of this fracture consists in letting down the sides of the box, and in inspecting the parts to see that the foot is kept in the proper position, and care should be taken to see that undue pressure is not made upon the skin by the compresses, which might result in ulceration; this may be avoided by sponging the skin with alcohol and changing the positions of the compresses slightly at each dressing. At the expiration of ten days the fracture-box and compresses may be removed and the limb may be put up in a plaster-of-Paris dressing including the foot and leg up to the knee. The patient may then be allowed to go about on crutches and at the end of five weeks all dressings may be dispensed with. A certain amount of stiffness and even permanent impairment in the motion of the ankle-joint often results from these fractures. This fracture is also dressed by means of *Dupuytren's splint*, which consists of a straight wooden splint long enough to extend from the condyles of the femur to end of the toes; this splint is provided with padding the thickest part of which, several

inches in thickness, should rest upon the skin just above the inner malleolus when the splint is applied to the inner side of the leg. The splint is applied to the inner surface of the leg with the thickest part of the pad resting upon the skin just above the inner malleolus, and is secured in position by the turns of a roller applied over the foot and at the upper part of the leg. After using this dressing for a few days if the displacement is satisfactorily corrected the splint may be removed and the leg may be placed in a fracture-box or in a plaster-of-Paris dressing. (Fig. 270.)

Fig. 270.



Dupuytren's splint applied.

This splint, when applied with sufficient firmness to correct the displacement, is not, as a rule, a comfortable dressing to the patient, so that in practice the use of the fracture-box and compresses will be found a more comfortable dressing and one equally satisfactory in correcting the deformity.

Fracture of the Bones of the Foot.

Fractures of the Tarsal Bones.

The *calcaneum* and *astragalus* are the tarsal bones most frequently fractured. The dressing of fractures of the *calcaneum* after reducing the displacement, which is not usually marked unless the posterior portion of the bone is involved, by manipulation, consists in placing the leg and foot in a fracture-box, and care should be taken to see that the foot is kept at a right angle to the leg. When the fracture involves the posterior portion of the bone and there is displacement

by the action of the muscles inserted into the fragment, the leg should be flexed upon the thigh and the foot should be extended; this position may be maintained by applying a well-padded curved splint to the anterior portion of the leg and foot and securing it in position by a bandage, or the same result may be obtained by applying a band or padded collar around the thigh, which is made fast by a cord or strap to the heel of a slipper applied to the foot. (Fig. 271.)

FIG. 271.

Apparatus for fracture of posterior portion of the calcaneum. (HAMILTON.)

Fractures of the *astragalus*, after reducing any deformity which is present by extension and manipulation, are dressed by placing the foot and leg in a fracture-box, care being taken to see that the foot is kept at a right angle to the leg. This precaution is important, as ankylosis not infrequently occurs after this fracture, and if the foot is in the proper position it is much more useful to the patient.

As soon as the swelling, which is usually very marked after fracture of the calcaneum or astragalus, subsides, the foot and leg should be put up in a plaster-of-Paris bandage. The amount of tension and the inability to reduce the displacement in cases of fracture of the astragalus may be indications for excision of the fractured bone. The time required for union in fractures of the tarsal bones is from five to six weeks.

Fractures of the Metatarsal Bones.

These fractures are dressed by placing the foot upon a well-padded plantar splint, and using compresses to hold the fragments in place if there is much displacement, the splint and compresses being held in position by a bandage; or they may be treated by placing the foot and leg in a fracture-box, the foot-board of the box acting as a plantar splint; the plaster-of-Paris dressing may also be used in these cases. The time required for union in fractures of the metatarsal bones is from three to four weeks.

Fractures of the Phalanges of the Toes.

These fractures are often compound and attended with so much laceration of the soft parts that immediate amputation is required; when, however, the fractures are simple, or in compound fractures where amputation is not required, the dressing consists in applying a plantar splint of wood or binder's board, extending beyond the toes and securing it in position by the turns of a roller bandage. When a single toe only is broken a moulded splint of gutta-percha or binder's board may be applied and a portion of the splint should extend some distance upon the sole of the foot to fix the proximal joint and also to give it a firm point of fixation; the moulded splint should be held in position by a narrow roller bandage or by strips of adhesive plaster. The time required for union in fractures of the phalanges of the toes is about three weeks.

Dressing of Compound or Open Fractures.

In the dressing of compound or open fractures the same dressings and splints which are generally used in the treatment of simple or closed fractures may be employed; the wound in the soft parts requires a special dressing and this should be so arranged as to secure free drainage and promote its prompt healing. In some cases of compound

fracture the treatment of the injures of the soft parts demands attention first, and in such cases the injury to the bones is for a time disregarded, care being taken to see that the fragments are kept quiet so as to prevent further damage to the soft parts until the wound is in such a condition that the proper manipulation to reduce the displacement and fix the fragments by splints and suitable dressings can be undertaken without interfering with the repair of the wound.

In the dressing of compound or open fractures the skin surrounding the wound should be first carefully cleansed and the wound should next be thoroughly irrigated with a 1 : 2000 bichloride solution or a 1 : 40 carbolic solution and any foreign bodies or loose fragments of bone should be removed, and if there is hemorrhage it should be controlled by securing the bleeding vessels with ligatures. The reduction of the displacement should next be accomplished by making extension and by manipulation (Fig. 272); if the

FIG. 272.

Method of reducing a compound fracture. (HAMILTON.)

fragments project from the wound, before this can be satisfactorily accomplished it may be necessary to enlarge the wound, and to resect one or both ends of the fractured bones, and in some cases it may be necessary to drill the ends of the fragments and introduce a strong wire or catgut suture to hold them in their proper positions. After reduction of

the displacement the wound should again be thoroughly irrigated with the antiseptic solution, and after making provision for drainage by the introduction of a drainage-tube or tubes, counter-openings being made to secure free drainage if necessary, the dressings should be applied.

The wound, if a small one, need not be closed with sutures; but, if extensive, a few catgut, silk, or silkworm-gut sutures may be applied to bring the edges of the wound into apposition, care being taken to avoid making undue tension; if the soft parts have been much lacerated or contused, it is better to introduce no sutures. A final irrigation of the wound through the drainage-tube is next made, and the wound is covered by a piece of protective, and the ordinary gauze dressing should be applied and covered by a number of layers of bichloride cotton, the whole dressing being held in position by a gauze bandage applied with moderate firmness.

The reduction of the fragments and the dressing of the wound having been accomplished as has been described, the splints and dressings appropriate for a similar fracture, if it were a simple or closed one, are next applied. If the surgeon has been able to render the wound aseptic, and has applied an antiseptic dressing, the compound fracture is often soon converted into a simple one, by the prompt healing of the wound, and the patient may exhibit no more constitutional disturbance than he would have with a similar simple or closed fracture. The re-dressing of a compound fracture dressed in this way need not be made for a week or ten days, unless there is a rise in the patient's temperature or the dressings become soaked with discharges from the wound, or they become uncomfortable to the patient by reason of swelling of the soft parts in the region of the wound. When the re-dressing of the fracture becomes necessary, the dressings are removed, and the drainage-tubes may be removed if no longer needed; the wound being re-dressed with an antiseptic dressing, the splints are reapplied, and, after the wound is healed, the subsequent dressing of the fracture should be the same as that of a simple fracture. The time required

for union in a compound fracture is usually much longer than in a corresponding simple fracture.

Many ingenious splints have been devised for the dressing of special compound fractures, but these were principally used before the introduction of the antiseptic method of wound-treatment, and as the treatment of these cases has been much simplified by its use, they possess no special advantage over the ordinary splints and dressings used in simple fractures.

The plaster-of-Paris dressing may be used as a primary dressing in compound fractures; the displacement being reduced and the wound being dressed with an antiseptic

FIG. 273.



Fenestrated plaster dressing for compound fracture of the leg. (STIMSON.)

gauze dressing, a plaster-of-Paris bandage is applied to the part so as to firmly fix the fragments; the joints on either side of the fracture should be fixed by the bandage, and the parts should be held in position until the plaster has set firmly. After the plaster has become firm, a fenestrum should be made over the position of the wound, so that it can be inspected or dressed through this when necessary. The ends of a piece of stout wire, bent into a semicircle, may be incorporated in the turns of the plaster bandage above and below the position of the fenestrum, to give it additional strength after the removal of a portion of the bandage to make the fenestrum. (Fig. 273).

If the plaster-of-Paris dressing is applied as a primary dressing in compound fractures the case should be carefully watched for a few days, and if much swelling occurs at the seat of fracture its removal and renewal is indicated; profuse discharge of serum may also soak the dressings and bandage so that its renewal is necessitated. Some surgeons, therefore, prefer to defer the application of the plaster-of-Paris dressing in compound fractures for a few weeks until the swelling has diminished and the wound is nearly or quite healed; the wound being covered with an antiseptic dressing the plaster bandage is applied and a fenestrum is made over the position of the wound if required.

Binder's-board or felt splints may also be employed in the dressing of compound fractures, being moulded to the parts after an antiseptic dressing has been applied to the wound, and held in position by the turns of a roller bandage.

The principal advantage in the use of these splints is the ease with which they can be removed and reapplied if frequent dressings of the fracture are necessary for any reason. They may be used during the course of treatment, or, after a few weeks when the swelling has diminished at the seat of fracture and the wound is well advanced toward repair, they may be discarded and a plaster-of-Paris dressing substituted. In compound fractures of the bones of the leg, after reducing the displacement and applying an antiseptic dressing to the wound, I usually apply moulded binder's board splints to either side of the leg, including the foot, and place the leg in a fracture-box for additional security, and after a few weeks I discard the binder's-board splints and apply a plaster-of-Paris dressing.

The *bran dressing* for compound fractures was formerly a popular dressing in this city, especially for compound fractures of the leg and thigh. It was applied by placing a piece of muslin or rubber cloth over the bottom and sides of a fracture-box and upon this was placed a layer of bran; the fractured leg was next placed in the box upon the layer of bran, the foot was then fastened to the footboard and the sides of the box were brought up and secured; bran was next poured into the box and firmly packed around and

over the limb. The bran absorbed the discharges which escaped from the wound and at the subsequent dressings the soiled bran was renewed without disturbing the limb and fresh bran was packed about the limb.

Sawdust which has been saturated with a solution of bichloride of mercury and dried may be used in the same manner as bran in the dressing of compound fractures and the former, which has been rendered antiseptic, has decided advantages over the bran dressing.

Continuous irrigation of compound fractures by a warm antiseptic solution either of bichloride of mercury 1 : 4000 or of carbolic acid 1 : 60 in cases in which so much contusion or laceration of the soft parts exists that the application of the ordinary dressings would be attended with the risk of gangrene, will be found a most satisfactory method of treatment. This dressing is applied by supporting the injured extremity upon a splint laid on a pillow covered by a rubber cloth, and a can or jar with a nozzle containing the solution is placed over the part and the irrigation is accomplished by allowing the fluid to run continuously over the wound; this irrigation may be kept up for days or weeks and when the vitality of the parts is assured, an antiseptic dressing with the ordinary splints or a plaster-of-Paris bandage may be applied.

PART IV.

DISLOCATIONS.

A DISLOCATION is the displacement of the articular surfaces of bones which enter into the formation of a joint.

Dislocations may be *complete*, *partial*, *simple*, *compound*, and *complicated*, and they are also known as *recent* and *old* dislocations, the latter terms being used not entirely with reference to the length of time the displacement of the articular surfaces of the bones has existed.

A complete dislocation is one in which no portions of the articular surfaces of the bones remain in contact with each other.

A partial dislocation is one in which portions of the articular surfaces of the bones still remain in contact with each other.

A simple dislocation is one in which there exists displacement in the relation of the articular surfaces of the bones with little injury to the soft parts adjacent to the joint, and the displaced ends of the bones do not communicate with the air by a wound in the soft parts.

A compound dislocation is one in which there exists displacement of the articular surfaces of the bones which communicates with the air through a wound in the soft parts.

A complicated dislocation is one in which in addition to the displacement of the articular surfaces of the bones, there exists a fracture, or a laceration of important bloodvessels, nerves, or muscles in proximity to the dislocation.

A recent dislocation is one in which the displacement of

the articulating surfaces of the bones has existed for such a period, that time has not been afforded for inflammatory changes to have taken place in the articular surfaces of the bones or in the adjacent tissues, which would seriously interfere with their reduction.

An old dislocation is one in which the displacement of the articulating surfaces of the bones has existed for some time, and in this variety of dislocation the displaced bones often form firm adhesions to the surrounding tissues.

Treatment of Dislocations.

The first indication in the treatment of dislocations is to return the displaced articular surfaces of the bones to their normal position and to retain them in this position by the use of suitable dressings The return of the articular surfaces of the bones to their normal position or *the reduction* of the dislocation, is accomplished by manipulation, extension, and counter-extension. The reduction of dislocations should be attempted as soon as possible after they have occurred.

The principal obstacles to the reduction of dislocations are muscular resistance and the anatomical peculiarities of the joints. The former is best overcome by the use of an anæsthetic given to the point where complete muscular relaxation is produced. The resistance offered by the changed relations of the articular surfaces and the ligaments is to be overcome by the surgeon making such manipulations, founded upon his knowledge of the anatomy of the parts, as will make the ligaments, muscles, and bones assist in the reduction of the dislocation.

In recent dislocations by the use of extension and manipulation, especially if an anæsthetic be employed, the reduction is usually accomplished without the use of much force, but in old dislocations, where absolute muscular shortening has taken place, the use of extending bands is often required, and in securing these bands to the limb the clove-hitch knot is useful. (Fig. 274.)

The treatment of dislocations after reduction consists in

placing the joint at complete rest by the application of suitable splints and bandages, and in treating any inflammatory complications if they arise, by the application of

FIG. 274.



Clove-hitch knot applied. (ERICHSEN.)

evaporating lotions, and in a week or two after the injured ligaments have been repaired, passive motion should be resorted to for restoring the function of the joint.

Special Dislocations.

DISLOCATIONS OF THE VERTEBRÆ.

Dislocations of the *lumbar* and *dorsal vertebræ*, as simple dislocations, are extremely rare accidents; they are occasionally met with, but are more often associated with fractures of the vertebræ in these regions; their occurrence in the cervical vertebræ is more common. The treatment of dislocations of the vertebræ, whether complicated with fracture or not, consists in attempting reduction by making extension and counter-extension with manipulation, and by this means in many cases the luxations can be reduced. If, however, the efforts at reduction are unsuccessful, permanent extension should be applied by means of a weight-extension apparatus from both legs, and from the shoulders and head. The after-treatment consists in keeping the patient at rest upon his back in bed upon a firm mattress, and if the cervical ver-

tebræ have been involved the head and neck should be supported by short sand-bags, and in case of the vertebræ below this point, the application of a plaster-of-Paris jacket may be used to give support and fixation to the parts. The general management of the case as regards complications is similar to that in cases of fracture of the vertebræ.

Dislocations of the coccyx are reduced by manipulation with the finger in the rectum and external manipulation at the same time. The only after-treatment required is rest in bed for a few days, and the administration of opium to keep the bowels quiet.

Dislocation of the Jaw.

This dislocation may consist in the displacement of one or both condyles of the jaw from the glenoid fossæ, consti-

Fig. 275.

Bilateral dislocation of the jaw. (Ashhurst.)

tuting the unilateral or bilateral dislocation of the jaw; the latter is the more common form of dislocation of the jaw

met with, and the deformity resulting is shown in Fig. 275.

The reduction of a dislocation of the lower jaw is accomplished as follows: The surgeon placing his thumbs, well protected by strips of bandage or a towel, on the molar teeth or behind them presses the angles of the jaw downward while he elevates the chin with his fingers, and by this manipula-

FIG. 276.

Method of reducing dislocation of the lower jaw. (HAMILTON.)

tion the condyles of the jaw usually slip back into place with a snap. After reduction of the dislocation the jaw should be fixed for a week or ten days by the application of a Barton's bandage or a four-tailed sling.

Dislocation of the Hyoid Bone.

A few cases of dislocations of the *hyoid* bone have been recorded; the treatment consists in throwing back the head as far as possible, to place the muscles of the neck upon the stretch, depressing the lower jaw and pressing the luxated bone into position.

DISLOCATIONS OF THE RIBS.

The *ribs* may be dislocated at their vertebral articulations, or at the junction with their costal cartilages. The treatment of these dislocations consists in reducing the displacements by manipulation and pressure and then in fixing the chest to

secure immobility of the ribs by strapping the affected side with strips of adhesive plaster, the same dressing being applied as in case of fracture of the ribs, the dressings being retained for three or four weeks.

Dislocation of the Sternum.

Dislocation or diastasis of the *sternum* may occur at the junction of the manubrium and gladiolus or at the junction of the ensiform cartilage and gladiolus. The reduction is effected by extension of the chest by bending the dorsal spine over a firm cushion placed under the back and by pressure upon the projecting bone; when the displaced bone has been reduced, a compress should be placed over the seat of injury and held in place by broad strips of adhesive plaster or by a bandage to keep the parts at rest. The dressing should be retained for three or four weeks.

In the few examples of dislocations of the ensiform cartilage which have been reported, the displacement of the cartilage has in some cases given rise to persistent vomiting, which was relieved by reduction of the displacement; it is, however, almost impossible to keep the fragment in place after reduction, and the vomiting gradually disappears after a time in these cases where it was impossible to keep the cartilage in its normal position.

Dislocations of the Pelvis.

Dislocations or diastasis of the bones of the *pelvis* may occur at the pubic or sacro-iliac symphyses.

These are generally serious injuries, as they are apt to be complicated by lesions of the pelvic viscera.

The reduction of these dislocations is effected by pressure and manipulation, and after reduction the parts should be supported by a compress held in place by a stout binder or by broad strips of adhesive plaster, the patient being kept quiet in bed, and the pelvis being supported by means of

sand-bags. The dressings should be retained for from four to six weeks.

Dislocations of the Clavicle.

Dislocations of the clavicle may occur either at the sternal or acromial end, and the latter injury some writers describe as a dislocation of the scapula, following the general rule that the distal bone is the one dislocated.

Dislocations of the sternal end of the clavicle may occur in a forward, backward, or upward direction, and the displacement is generally well marked. (Fig. 277.) The reduction of this dislocation is effected by placing the knee against the spine and drawing the shoulders outward and backward and pressing the displaced end of the clavicle

Fig. 277.

Dislocation of sternal end of clavicle forward. (Bryant.)

Fig. 278.

Dislocation of clavicle at acromial end. (Bryant.)

into place. The reduction is generally easy, but it is often difficult to keep the end of the bone in its proper position. To accomplish this, a compress should be placed over the end of the bone, and this should be secured in place by broad strips of adhesive plaster; the shoulders should be brought well backward and secured by a posterior figure-of-eight bandage of the chest, and the arm of the injured side

should be fastened to the side of the chest by spiral turns of a bandage. In some cases, in addition to the compress over the end of the bone, securing the arm of the injured side in the Velpeau position will be found all that is necessary to retain the bone in position.

Dislocation of the acromial end of the clavicle may be upward, downward, or backward. (Fig. 278.) The reduction is effected by manipulation of the arm and scapula and by pressure over the displaced end of the clavicle; the displacement is usually reduced without much trouble, but it is often a matter of difficulty to keep the end of the bone in its proper place.

The dressing consists in placing a compress over the acromial end of the clavicle and holding it in place by broad strips of adhesive plaster; the arm should at the same time be fixed in the Velpeau position. These dressings after reduction of dislocations of the clavicle should be kept in place for at least three weeks. Although in many cases a certain amount of deformity persists, the disability resulting from the injury is not often marked.

Dislocations of the Scapula.

Dislocation of the acromion processes of the scapula from the outer end of the clavicle, which has been described under dislocation of the acromial end of the clavicle, is classed by some writers as a scapular dislocation.

Dislocation or projection of the inferior angle of the scapula, due to its escape from under the latissimus dorsi muscle or relaxation of this muscle and of the serratus magnus, is sometimes described as a dislocation of the inferior angle of the scapula. The reduction of this deformity consists in the employment of manipulation and pressure to overcome the displacement, and the use of a compress held in place by broad strips of adhesive plaster to secure the bone in its proper position.

Dislocations of the Shoulder.

The head of the humerus may be dislocated downward, forward, or backward.

Subglenoid or downward dislocation of the head of the humerus is that variety of dislocation in which the head of the bone rests in the axilla. (Fig. 279.)

Fig. 279.



Subglenoid dislocation of the shoulder. (Stimson.)

Subcoracoid or forward dislocation of the head of the humerus is that variety of dislocation in which the head of the humerus rests beneath the coracoid process of the scapula. (Fig. 280.)

Subclavicular dislocation of the head of the humerus may be considered an aggravated form of the latter variety of

dislocation; the head of the humerus in this variety of dislocation rests beneath the clavicle.

FIG. 280.

Subcoracoid dislocation of the shoulder. (STIMSON.)

Subspinous or backward dislocation of the head of the humerus is that variety of dislocation in which the head of the humerus rests beneath the spine of the scapula. (Fig. 281.)

The reduction of dislocations of the humerus is effected by manipulation, by extension and counter-extension, and by a combination of these methods.

Manipulation in the reduction of *subglenoid dislocation* of the humerus is practised by first flexing the forearm upon the arm to relax the long head of the biceps muscle; the elbow is next seized and abducted so as to bring it to the side of the patient's head, thus relaxing the deltoid and supra-spinous muscles; the surgeon or an assistant next

places his hand upon the head of the humerus in the axilla, and, as the arm is drawn outward to a right angle with the

FIG. 281.

Subspinous dislocation of the head of the humerus. (ERICHSEN.)

body by the other hand, he pushes the head of the bone into the glenoid cavity.

In the reduction of *subglenoid* and *subclavicular* dislocations the manipulation is the same except that the arm is to be rotated outward before being carried downward.

In the reduction of *subspinous* dislocations after the arm has been abducted it should be rotated inward and direct pressure should be made upon the head of the bone as the arm is adducted. Reduction may also be effected by extension and counter-extension as in Cooper's method, where extension is made from the arm downward and counter-extension is made by the heel in the axilla. (Fig. 282.)

Reduction may also be accomplished by extension made upward, as in Mothe's method, the scapula being fixed by the foot or hand placed over the acromion process. (Fig. 283.)

After reduction of dislocations of the head of the humerus the arm should be bound to the side of the body by the

FIG. 282.

Reduction of shoulder by heel in the axilla. (ERICHSEN.)

FIG. 283.

Reduction of shoulder by extension upward.

turns of a spiral bandage of the chest, or should be held against the side by the application of a Velpeau bandage (Fig. 46, p. 55); this dressing should be removed at in-

tervals of a few days, and after ten days or two weeks all dressings should be dispensed with, passive motion should be employed and the patient should be allowed to move the arm.

Dislocations of the Elbow.

Dislocation of the Bones of the Forearm.

Dislocations of the bones of the forearm at the elbow may either be backward, forward, or lateral. The backward dislocation is the most common form. (Fig. 284.)

Fig. 284.

Dislocation of both bones of the forearm backward. (Liston.)

The reduction of *backward dislocations* is effected by making traction upon the forearm and at the same time making pressure upon the lower end of the humerus as the forearm is flexed upon the arm.

Or the reduction may be accomplished by bending the arm slowly and forcibly over the knee placed upon the inner surface of the elbow so as to press upon the radius and ulna, separating them from the humerus and freeing the coronoid process from its abnormal position. (Fig. 285.)

Lateral dislocations of the bones of the forearm at the elbow are reduced by making extension from the forearm, and at the same time making direct pressure on the displaced bones and counter-pressure on the lower end of the humerus.

Forward dislocations of the bones of the forearm at the elbow are reduced by making forced flexion at the elbow, together with extension or counter-extension, or by making

FIG. 285.

Reduction with the knee in the bend of the elbow. (HAMILTON.)

FIG 286.

Dressing after reduction of dislocations of the elbow.

forced extension of the forearm at the elbow, pressing the humerus backward and suddenly flexing the forearm.

The dressing, after the reduction of dislocations at the elbow, consists in the application of a well-padded anterior

FIG. 287.



Dislocation of head of the radius forward. (LISTON.)

right- or slightly obtuse-angled splint, to keep the forearm in a flexed position—the dressing being practically the same as that for fractures of the lower end of the humerus, with an anterior angular splint (Fig. 286). This dressing should be retained for two or three weeks, being removed at intervals of several days; after the removal of the splint, passive

motion should be practised, to prevent stiffness of the elbow-joint.

Dislocation of the Head of the Radius.

The head of the radius may be displaced forward, outward, or backward, the forward dislocation being the most frequent. (Fig. 287.) The reduction of these dislocations is effected by making extension from the forearm and counter-extension from the lower end of the humerus, and at the same time the head of the bone is pressed into its proper position. The dressing after reduction of the displacement consists in the application of a compress over the head of the bone, and the arm and forearm should be placed upon a well-padded anterior angular splint, which is secured by a roller bandage. The dressing is similar to that employed in fractures of the lower end of the humerus, in which an anterior angular splint is employed (Fig. 242, page 325). Difficulty is sometimes experienced in keeping the head of the bone in position after reduction, so that the use of the compress in addition to the use of the splint is often required. The arm should be kept upon the splint for three weeks, being redressed at intervals.

Dislocation of the Upper End of the Ulna.

The upper end of the ulna may be displaced backward, the olecranon projecting behind the condyles of the humerus, while the head of the radius occupies its normal position. The reduction of this displacement is effected in the same manner as that of both bones of the forearm backward, and the dressing after reduction is similar to that employed when both bones have been displaced.

Dislocations at the Wrist.

The lower end of the ulna may be dislocated from the radius forward, backward, or inward. The reduction of these displacements is effected by fixing the radius and pushing the ulna back into place. The dressing after reduction

consists in placing the wrist-joint at rest by the application of well-padded anterior and posterior straight splints. The splints should be retained for three or four weeks, dressings being made at intervals of two or three days.

Dislocations of the carpus upon the bones of the forearm may be forward (Fig. 288), or backward (Fig. 289). The

FIG. 288.

Dislocation of the carpus forward. (HAMILTON.)

FIG. 289.



Dislocation of the carpus backward. (HAMILTON.)

reduction in either variety of displacement is effected by extension from the hand and by pressure. After reduction of the displacement, which does not tend to recur, the hand and forearm should be placed upon a well-padded straight splint applied to the palmar surface of the hand and forearm. The splint should be retained for ten days or two weeks.

DISLOCATIONS OF THE BONES OF THE CARPUS.

The displacement of the individual bones of the carpus occasionally takes place, the *os magnum*, the *semilunare* and *pisiform* being the bones most usually displaced, although other bones of the carpus are sometimes dislocated. Reduc-

tion is effected by means of extension and pressure, and the part should afterward be dressed with a palmar splint and compresses.

Dislocations of the Metacarpal Bones.

The metacarpal bones may be dislocated upon the carpus; the bones most commonly displaced are those of the thumb, and of the index and middle fingers; the latter are usually displaced backward, while the metacarpal bone of the thumb may go either backward or forward.

Reduction is effected by extension and pressure. The dressing after reduction consists in the application of a palmar splint to the hand and forearm and a compress over the displaced bone. The dressings should be retained for two weeks.

Dislocations of the Fingers.

Dislocations of the phalanges of the hand usually take place at the metacarpo-phalangeal junction, but sometimes occur at the intra-phalangeal joints. The reduction is usually easily effected by extension (Fig. 290), or by push-

Fig. 290.

Backward dislocation of phalanx. Reduction by extension. (Hamilton)

ing the phalanx back until it stands perpendicularly upon the metacarpal bone, when by strong pressure upon its base, from behind forward, it is readily carried by flexion into its natural position.

Where difficulty is experienced in making extension in the reduction of these dislocations, the ingenious apparatus of the late Dr. Levis (Fig. 291), or the "Indian puzzle" apparatus (Fig. 292) may be employed with success.

FIG. 291.



Levis's apparatus for dislocation of the phalanges applied.

FIG. 292.

Extension by Indian puzzle. (BRYANT.)

In *dislocations of the proximal phalanx of the thumb backward* (Fig. 293), great difficulty in reduction is often

FIG. 293.



Dislocation of proximal phalanx of thumb backward. (FARABEUF.)

experienced from the head of the metacarpal bone slipping between the two heads of the short flexor muscle, or be-

tween the lateral ligaments. The interposition of the external sesamoid bone is considered by some surgeons to be the cause of difficulty in the reduction of this displacement.

In this dislocation reduction is effected by firmly pressing the metacarpal bone of the thumb strongly toward the palm of the hand to relax the two portions of the short flexor muscle. The thumb is next extended upon the wrist until its tip points to the elbow. An assistant now places his finger behind the proximal phalanx to prevent its slipping backward and by bringing the thumb down to the flexed position the bone slips into place. It sometimes happens that all efforts at reduction fail, and in such cases

FIG. 294.

Backward and upward dislocation of femur. (COOPER.)

FIG. 295.

Backward dislocation of femur. (COOPER.)

it may be necessary to divide one head of the short flexor muscle subcutaneously or through an open wound, before the displacement can be relieved.

The dressing of dislocations of the phalanges after reduction consists in the application of splints of wood, or moulded splints of binder's board or gutta-percha to fix the joint, which should be retained for ten days or two weeks.

Dislocations of the Hip.

The head of the femur is most frequently dislocated backward, downward or upward, although it may assume other positions in exceptional cases.

Posterior or *backward dislocations of the head of the femur* are either backward and upward, and are described as *iliac* or *dorsal*, the bone resting upon the dorsum of the ilium (Fig. 294). Or the dislocation may be backward, the head of the bone resting upon the ichiatic notch; these are known as *ischiatic* dislocations or dislocations of the femur *dorsal below the tendon* (of the obturator internus), according to Bigelow (Fig. 295).

Fig. 296.

Reduction of backward dislocation of femur. (Bigelow.)

The reduction of the posterior dislocations of the femur can generally be effected by manipulation. The patient being anæsthetized and placed upon his back, the surgeon grasps the leg at the ankle and knee, flexes the leg upon the thigh, and the thigh upon the pelvis; he then abducts the limb and rotates it outward, bringing it in a broad sweep across the abdomen, and by bringing it down to its natural position the head of the bone will slip into the acetabulum (Fig. 296).

Downward Dislocation of the Head of the Femur, or Downward and Forward Dislocation.—In this variety of

dislocation the head of the bone rests upon the thyroid foramen; this form of displacement is sometimes spoken of as a *thyroid* dislocation. (Fig. 297.)

The reduction of downward and forward dislocations of the head of the femur is effected by flexing the leg and thigh and bringing the limb into a position of abduction; it is then adducted and rotated inward in a broad sweep across the abdomen and brought down to its natural position, and the head of the bone slips into the acetabulum. (Fig. 298.)

FIG. 297.

Downward and forward dislocation of femur. (COOPER.)

FIG. 298.

Reduction of downward and for ward dislocation of he femur (BIGELOW.)

In making these manipulations the head of the bone sometimes slips back upon the dorsum of the ilium, converting the downward dislocation into a posterior one if

this accident occurs the displacement should be remedied by making the manipulation appropriate for the reduction of the latter dislocation.

Upward Dislocation, or the Dislocation Forward and Upward, of the Head of the Femur.—In this variety of dislocation the head of the bone rests upon the pubis; this form of displacement is also spoken of as a pubic dislocation. (Fig. 299.)

Fig. 299.

Forward and upward dislocation of the femur. (Cooper.)

The reduction of forward and upward dislocations of the head of the femur is effected by much the same manipulation as is employed in the reduction of downward and forward dislocations, except that in the pubic dislocation the flexed limb should be carried across the sound thigh at a higher point. The thigh being flexed the head of the bone is drawn down from the pubis; it is then semi-abducted and rotated inward to disengage the bone completely. While rotating inward and drawing on the thigh the knee should be carried inward and downward to its place by the side of its fellow, and the head of the bone will usually slip into the acetabulum.

As before stated various *anomalous displacements of the head of the femur* occasionally occur; the head of the bone may pass directly upward, or downward between the sciatic notch and thyroid foramen, or downward and backward on the body of the ischium, or downward and backward into the lesser sciatic notch, or downward, inward, and forward into the perineum. These anomalous displacements

usually occur where there has been extensive laceration of the capsular and Y-ligaments.

The dressing of cases, after reduction of dislocations of the head of the femur, consists in keeping the patient at rest in bed upon his back, and the limb should be kept at rest by sand-bags applied to either side of the limb, or the knees should be tied together.

The patient should be kept at rest for two or three weeks, and at the end of this time may be allowed to get out of bed and go about on crutches.

Dislocations of the Patella.

The patella may be dislocated *outward*, *inward*, or *upward*, or it may be rotated upon its own axis. The *outward* dislocation is the displacement most usually seen. (Fig. 300.)

Fig. 300.



Outward dislocation of the patella. (Duplay.)

Upward dislocation of the patella can only result from laceration of the ligamentum patellæ, and the treatment in such cases is similar to that for fracture of the patella.

The reduction of dislocations of the patella is effected by extending the leg upon the thigh, and flexing the thigh upon the pelvis to relax the quadriceps femoris muscle, when the patella can usually be forced back into place; in some cases alternate flexion and extension of the leg will accomplish the same result.

The dressing after reduction of the displacement consists in the application of a posterior straight splint or a moulded binder's-board or felt splint to keep the joint at rest; the splint should be worn for a week or ten days.

Dislocations of the Knee.

The head of the tibia may be dislocated *forward*, *backward*, or *laterally;* the latter dislocations are always incomplete, forward dislocation being the variety of displacement most commonly met with. (Fig. 301.)

Fig. 301.



Forward dislocation of the knee. (Bryant.)

The reduction of dislocations of the knee is effected by extension and counter-extension with forced flexion of the knee with pressure, aided by rocking movements. The treatment of cases of dislocation of the knee after reduction consists in fixing the knee-joint by the application of a straight posterior splint or a moulded splint of binder's board. As there is usually marked swelling following these injuries from violence to the joint-structures, the application of evaporating lotions for a few days will be found useful. As soon as the swelling has subsided the joint should be put up in a plaster-of-Paris dressing and this should be retained for four weeks.

Dislocation of the Semilunar Cartilages.

The displacement here consists in the slipping forward or backward and wedging of the semilunar cartilages between the femoral condyles and the tibia.

Reduction of the displaced cartilages can usually be effected by hyper-flexion of the knee followed by sudden full extension, or by alternately flexing and extending the

joint. Excision of the displaced cartilages is sometimes required in cases in which they cannot be reduced by manipulation.

The dressing of these cases after reduction of the displaced cartilages consists in the application of a posterior straight splint or a plaster-of-Paris dressing to fix the knee-joint; the splint should be worn for three or four weeks, and if there is a tendency to redisplacement the patient should wear a knee-cap of leather or muslin to partially fix the joint, with compresses so arranged as to make pressure upon the edge of the joint.

Dislocation of the Fibula.

Dislocations of the fibula may occur at either of its extremities, and the direction of the displacement may be *forward*, *backward*, or *upward*, dislocation of the head or upper extremity of the fibula being the most common, although all are rare forms of displacement.

The reduction of dislocations of the head of the fibula is effected by flexing the leg upon the thigh and making direct pressure and extension. Dislocations of the lower extremity of the fibula are reduced by manipulation and pressure. The dressing of cases after reduction of dislocations of the fibula consists in the application of a compress and moulded binder's board splint, and the dressing should be retained for three or four weeks.

Dislocations of the Ankle.

Dislocation of the foot upon the bones of the leg results from the separation of the articular surface of the astragalus from that of the tibia and fibula, and the displacement may be *forward*, *backward* (Fig. 302), or *lateral* (Fig. 303), the latter variety being often associated with fracture of the malleoli.

The reduction of dislocations of the ankle is effected by traction, combined with flexion and rotation of the ankle-

joint, the leg being first flexed upon the thigh to relax the tendo Achillis, and in some cases the subcutaneous division of this tendon is required before the reduction can be satisfactorily accomplished.

FIG. 302.

Dislocation of foot backward.
(BRYANT.)

FIG. 303.

Dislocation of foot inward.
(BRYANT.)

The dressing of dislocations of the ankle after reduction consists in the application of a fracture-box, or of pasteboard splints to fix the ankle, care being taken to see that the foot is fixed at a right angle to the leg, and in the application of evaporating lotions for a few days; after the swelling has subsided, a plaster-of-Paris dressing should be applied and retained for three or four weeks.

DISLOCATIONS OF THE TARSAL BONES.

The *astragalus* may be dislocated from the bones of the leg and from the other tarsal bones, being thrust *forward*, *backward*, *outward* (Fig. 304), or *inward*. The reduction of dislocations of the *astragalus outward* is effected by first flex-

ing the leg upon the thigh and making extension from the foot and rotating it at the same time, direct pressure being made upon the displaced bone; in some cases subcutaneous section of the tendo Achillis has assisted materially in the reduction of the displaced bone. *Backward dislocation of the astragalus* is usually irreducible, the patient, however, in many cases recovers with a useful foot. In cases of irreducible dislocations of the astragalus, excision of the astragalus may ultimately be required.

FIG. 304.

Dislocation of astragalus outward. (HAMILTON.)

After the reduction of dislocations of the astragalus, the foot and leg should be put at rest in a fracture-box, or by means of moulded splints of pasteboard or felt; evaporating lotions should also be employed to the region of the injury for a few days, and when the swelling has subsided, a plaster-of-Paris dressing should be applied and retained for three or four weeks.

Dislocations of the calcaneum and scaphoid upon the astragalus, or of the *calcaneum* upon the *astragalus* and *cuboid*, or upon the *astragalus* alone; of the *scaphoid* and *cuboid* upon the *calcis* and *astragalus;* or of the *cuboid*, *scaphoid*, or *cuneiform* bones, are occasionally met with.

Their reduction is effected by traction and direct pressure, and, after this has been accomplished, the parts should be put at rest by the application of a splint and compresses.

Dislocations of the Metatarsal Bones and Phalanges of the Toes.

These dislocations usually result from crushing forces which destroy the vitality of the soft parts so completely that amputation is required. Their reduction in cases of simple or uncomplicated dislocations is effected by traction, manipulation, and pressure. After reduction of the displacement, the parts should be kept in position by the application of splints and compresses.

Old Dislocations.

The reduction of old dislocations is attended with more difficulty and danger than that of recent dislocations, due to the permanent contraction and structural changes which

Fig. 305.



Reduction of old dislocation of the femur by pulleys. (Cooper.)

occur in the muscles, and to the abnormal adhesions which form between the displaced bone and the parts with which it is in contact. The reduction of old dislocations can usually be accomplished by the manipulations appropriate for recent dislocations of the same variety, but occasionally the use of more forcible extension is required, which is made by bands

and pulleys (Fig. 305), or by vertical extension (Fig. 306). The first step in the reduction of old dislocations consists in thoroughly breaking up the adhesions which have been formed between the displaced bone and the surrounding tissues; this has, in some cases, resulted in the laceration of

FIG. 306.



Reduction of old dislocation of hip by vertical extension. (BIGELOW.)

muscles, nerves, and bloodvessels, and in the fracture of the displaced bones or neighboring bones, so that the manipulations should be made with the least force that will accomplish the object desired. After the reduction of old dislocations, difficulty is sometimes experienced in maintaining the bone in its proper place, due to the changes which have occurred in the articular surfaces.

Compound Dislocations.

These are always grave injuries, and amputation or excision is often required. When, however, operative measures are not required, the reduction is effected in the same manner as in simple dislocations of corresponding parts, the greatest care being taken to render the wound aseptic, and to keep it in this condition by the application of a full antiseptic dressing.

Complicated Dislocations.

In dislocations complicated by fracture near the seat of displacement, the displaced bone should, if possible, be first reduced, and this in many cases is a matter of much difficulty as the fracture prevents the surgeon from using leverage otherwise present, in the reduction, and he has often to depend entirely upon pressure and manipulation to restore the displacement.

After reduction of the dislocation the fracture should be reduced and dressed.

Dislocation complicated by rupture of the main artery of the limb may require, after reduction of the displacement, exposure and ligation of the vessel or amputation of the limb. Rupture of an important nerve trunk complicating a dislocation may call for subsequent exposure and suturing of the divided nerve.

Spontaneous Pathological and Congenital Dislocations.

In the treatment of these varieties of dislocations after the reduction of the displacement by manipulation and pressure much difficulty is often experienced in maintaining the reduction. To effect the latter object the use of splints and bandages is employed and also the use of many ingenious forms of apparatus adapted to particular dislocations.

Tenotomy or *myotomy* are often required to prevent recurrence of the deformity, and continuous extension is also of much value in the treatment of these displacements.

PART V.

LIGATION OF ARTERIES.

In the application of a ligature to an artery in its continuity the surgeon should make his incision in the line which corresponds to the general course of the vessel and he should be thoroughly familiar with the anatomy and with the surgical landmarks of the part. A portion of the vessel, when possible, should be selected for the application of the ligature half an inch or an inch from any large collateral branch. The position of the incision being selected the surgeon steadies the skin with two fingers and makes an incision of the required length through it with a scalpel; the superficial fascia is next picked up on a director, any large superficial veins which come into view being displaced, and divided to an equal length with the incision in the skin; the deep fascia being exposed it should be nicked and divided upon a director; the inter-muscular space or the edge of the muscle or muscles which are the guide to the vessel should next be sought for, small vessels coming from the main vessel through these spaces will often serve as valuable guides to the position of the vessel. The surgeon next separates the tissues with the director, handle of the knife, or the finger until the sheath of the vessel is exposed; this is recognized by its communicated pulsation and by the absence of the smooth shining surface and pinkish-white color which the surface of the artery presents. The sheath of the artery should be picked up with forceps and nicked with the point of the knife applied flatwise; the incision into the sheath should be very limited, only large enough to allow the

aneurism needle to pass through it around the vessel; extensive dissections or separations of the sheath from the vessel should be avoided as the nutrition of the artery at the point of ligature may thus be impaired and sloughing and secondary hemorrhage may result. A distinct sheath is found only about the main arterial trunks, which is replaced in the smaller arteries by a layer of loose cellular tissue. The wall of the artery being exposed an aneurism needle is

FIG. 307.



Opening sheath. Passing ligature around the vessel. Tying artery. (BRYANT.)

FIG. 308.

Aneurism needle.

passed around the vessel, threaded with a catgut ligature, and withdrawn; the needle may be threaded before being passed, in which case the ligature is grasped with forceps and drawn through while the needle is withdrawn. The best material for ligatures is carefully prepared chromicized catgut. The needle should be passed away from important structures such as accompanying veins and nerves.

Before the ligature is tied the surgeon should satisfy himself that the ligature when tied will control the circulation

in the vessel below its point of application, by placing the tip of his finger upon the vessel and drawing upon the ends of the ligature so as to occlude the vessel at the point of application. Being satisfied as to this point the ligature is tied with a reef-knot, or a surgeon's-knot and reef-knot combined.

Some authorities recommend the application of two ligatures a short distance apart in the ligation of vessels in their continuity, and a division of the vessel between them, so that both ends can retract into the cellular sheath.

The ends of the ligature are cut short in the wound, which is irrigated and drained if necessary, and is closed by the application of a few sutures, and an antiseptic dressing is applied.

Ligation of Special Arteries.

Ligation of the Innominate Artery.

The innominate artery lies immediately behind the sterno-clavicular articulation, and is in relation in front with the innominate veins and pneumogastric nerve, on the inner side with the trachea, on the outer side and behind with the pleura.

Incision.—A V-shaped incision, each branch of which is two and a half or three inches in length, one of which lies over the anterior edge of the sterno-cleido-mastoid muscle, and the other parallel to and a little above the clavicle. (Fig. 309, *A*.) The incisions are carried down to the superficial fascia and a flap is dissected up. If the anterior jugular vein is met with it should be displaced. The sternal and clavicular attachments of the sterno-cleido-mastoid are next divided upon a director half an inch above the bone. The sterno-thyroid and sterno-hyoid muscles and the middle cervical fascia are next exposed, covered by the thyroid veins. The outer fibres of the sterno-hyoid and sterno-thyroid muscles are next divided, the thyroid vein being held aside, when upon tearing through the fascia with a director the common carotid artery is exposed and traced down to the

innominate artery; the innominate veins are pressed against the sternum with the finger and the artery is separated from its sheath about half an inch below its bifurcation, and the

FIG. 309.



Line of incision for—*A*, innominate artery; *B*, right subclavian artery; *C*, left subclavian artery; *D*, vertebral or inferior thyroid artery; *E*, axillary artery below clavicle. (STIMSON.)

aneurism needle is passed around the vessel from the outer side so as to avoid the vein, pneumogastric nerve, and pleura.

LIGATION OF THE SUBCLAVIAN ARTERY.

This artery may be tied at three points; in its *first* portion, between the trachea and scaleni muscles; in its *second* portion, behind the scaleni muscles, and in its *third* portion external to the scaleni muscles.

The left subclavian artery in its first portion is larger and more vertical in its direction than the right subclavian and is situated more posteriorly; from the difficulty in exposing

this portion, and from the possibility of injuring the thoracic duct, the ligation of this artery in its first portion has been seldom attempted.

Incision for the *first portion* of the subclavian artery is the same as that for the innominate (Fig. 309, *A*), and the ligature is passed from the outer side, the pneumogastric and phrenic nerves being pressed inward toward the carotid artery.

The right or left subclavian arteries are also seldom tied in their second portions, that is behind the scaleni muscles, but are frequently tied in their third portions, that is external to the scaleni muscles.

FIG. 310.



Ligation of subclavian and lingual arteries. (BRYANT.)

Incision for the *second portion* of the subclavian artery begins an inch external to the sterno-clavicular articulation half an inch above and parallel to the clavicle, and is three or four inches in length. (Fig. 309, *B* or *C*.) The steps of the operation are the same as for ligation of the third portion, and when the scalenus anticus muscle has been exposed it is divided upon a director; the phrenic nerve which lies upon its anterior aspect is to be avoided.

Incision for the *third portion* of the subclavian artery is the same as for the second portion. (Fig. 309, *B* or *C*.) The skin and platysma being divided, the jugular vein is exposed and drawn to one side or divided between the ligatures; the superficial fascia is next divided upon a director; the posterior belly of the omo-hyoid muscle is next found and drawn upward and outward; the outer border of the scalenus anticus is next felt for and followed down to the tubercle of the first rib—the artery lies against this, between it and the lowest bundle of the brachial plexus. The artery is next denuded with the director and the needle is passed from below, care being taken not to include the lowest bundle of the brachial plexus in the ligature. (Fig. 310.)

Ligation of the Vertebral Artery.

Incision for the ligation of the vertebral artery is three or three and a half inches in length, parallel with the anterior edge of the sterno-cleido-mastoid muscle, ending an inch above the clavicle. (Fig. 309, *D*.) The anterior edge of the sterno-cleido-mastoid being exposed the middle cervical fascia is divided and the carotid artery and jugular vein are exposed and drawn inward. The gap between the longus colli muscle and the scalenus anticus muscles is next felt for about an inch below the carotid tubercle; the fascia covering it is next torn through and the muscles are separated and the vertebral vein comes into view; this is held aside and the vertebral artery is exposed, and the ligature is then passed around it.

Ligation of the Inferior Thyroid Artery.

Incision for the inferior thyroid artery is the same as that for the vertebral artery. (Fig. 309, *D*.) The anterior edge of the sterno-cleido-mastoid muscle being exposed it is drawn outward, the middle cervical fascia is next divided and the carotid artery and internal jugular vein are drawn outward with a retractor. The head being flexed slightly, the surgeon feels for the carotid tubercle, and then separates the cellular tissue with a director and the artery should be found below the carotid tubercle. The needle should be passed between the artery and vein.

Ligation of the Common Carotid Artery.

The point of election for the ligation of the common carotid artery is just above the omo-hyoid muscle, about

Fig. 311.



Line of incision for common carotid artery at point of election. (Stimson.)

three-quarters of an inch below the bifurcation of the vessel, which takes place at a point on a line with the upper border of the thyroid cartilage.

Incision for the common carotid artery, is three inches in length along the anterior border of the sterno-cleido-mastoid muscle, the centre of which corresponds with the crico-thyroid space. (Fig. 311.)

Divide the skin, platysma and cellular tissue and aponeurosis, avoiding the superficial veins, and expose the anterior edge of the sterno-cleido-mastoid; seek for the inter-

FIG. 312.



Ligation of common carotid artery. Ligation of facial artery. (BRYANT.)

space between this muscle and the sterno-hyoid and sterno-thyroid muscles, draw the latter muscles inward and the artery will be exposed with the jugular vein external to it; the descendens noni nerve lying upon its sheath, which should be displaced outward. The sheath is next picked up and opened and the artery is separated from it with a director;

the artery lies internally, the internal jugular vein externally and somewhat more superficial, and the pneumogastric nerve lies between the two and is more deeply placed. (Fig. 312.)

FIG. 313.



Relation of the left common carotid artery above the omo-hyoid muscle. (ESMARCH.)

The sympathetic nerve is posterior to the vessel external to the sheath. The needle is passed from without inward, care being taken to avoid injury of the vein and nerve. (Fig. 313.)

LIGATION OF THE EXTERNAL CAROTID ARTERY.

Incision for the ligation of the external carotid artery is over the inner edge of the sterno-cleido-mastoid muscle from the angle of the jaw to a point corresponding to the middle of the thyroid cartilage. (Fig. 314, *B*.) The skin, platysma and cellular tissue being divided, the external jugular vein is drawn aside when encountered; the deep fascia being

opened, the facial and lingual veins will be exposed, which should be drawn to one side; the artery is next exposed covered by the hypoglossal nerve and the stylo-hyoid and digastric muscles. The vessel should next be isolated from the internal carotid artery and internal jugular vein, both of which lie along its outer side. The needle should be passed from without inward.

Ligation of the Internal Carotid Artery.

Incision the same as for the external carotid artery (Fig. 314, *B*); the vessel is external to the external carotid artery,

Fig. 314.



Line of incision for—*A*. Lingual artery. *B*. External and internal carotid artery. *C*. Occipital artery. *D*. Temporal artery. *E*. Facial artery. (Stimson.)

and in passing the needle the point should be directed from the internal jugular vein, that is from without inward.

Ligation of the Superior Thyroid Artery.

Incision about three inches in length along the anterior border of the sterno-cleido-mastoid muscle, starting a little lower down than that for the external carotid artery. The skin, superficial fascia, platysma, and deep fascia being divided, the cellular tissue in the sulcus between the upper portion of the larynx and the great vessels of the neck is broken up with the director and the vessel is exposed. The needle should be passed around the vessel from above downward.

Ligation of the Lingual Artery.

Incision a curved one two inches long, its concavity directed upward from the anterior edge of the sterno-cleido-mastoid muscle, half an inch above the great horn of the

Fig. 315.



Relations of the lingual artery. (Esmarch.)

hyoid bone, to a point one inch short of the median line of the neck. (Fig. 314, *A*.) Divide the skin and platysma, displacing the superficial veins, and open the deep fascia, when the submaxillary gland will be exposed; this is displaced upward with the handle of the knife and the tendon

of the digastric muscle attached to the hyoid bone, and the hypoglossal nerve will be exposed; next divide the fibres of the hyoglossus muscle midway between the hypoglossal nerve and the hyoid bone, and the lingual artery will be exposed. (Fig. 315.)

The needle should be passed around the vessel from above downward in order to avoid the nerve.

Ligation of the Facial Artery.

The facial artery passes over the inferior maxilla just in front of the anterior edge of the masseter muscle and is accompanied by the facial vein, which lies nearer to the muscle.

Incision either a horizontal one along the lower border of the maxilla or a vertical one an inch in length. (Fig. 314, *E.*) The skin, subcutaneous tissue, and fascia being divided, the artery is exposed and the needle should be passed around the vessel away from the vein.

Ligation of the Occipital Artery.

Incision two inches in length, starting from a point half an inch below and in front of the apex of the mastoid process carried obliquely backward parallel to the border of this process. (Fig. 314, *C.*) Divide the skin and fascia and expose the insertion of the sterno-cleido-mastoid muscle, which is also divided, and the aponeurosis of the splenius is exposed; this is also opened and the digastric groove is felt for, and when the belly of the digastric muscle is exposed the artery is brought into view by separating the cellular tissue in the anterior angle of the wound with a director. (Fig. 316.)

Ligation of the Temporal Artery.

Incision a transverse one, one inch in length, starting from the tragus of the ear forward over the zygomatic arch

(Fig. 313, *D*), or a vertical one of the same length a little in front of the tragus of the ear.

Divide the skin and expose the subcutaneous cellular tissue, which in this region is very dense and fibrous. This

Fig. 316.

Ligation of the occipital artery. (Skey.)

Fig. 317.

Ligation of the temporal artery. (Skey.)

tissue should be broken up with a director and the artery should be found in it about a quarter of an inch in front of the ear. (Fig. 317.) The temporal vein accompanies the artery and lies nearer to the ear, and in some cases the auriculo-temporal nerve is in close relation to the artery. The needle should be passed from behind forward.

Ligation of the Axillary Artery.

The axillary artery extends from the middle of the clavicle to the insertion of the teres major into the humerus; the axillary vein lies upon the inner side and in front of the artery. The axillary artery is tied either in its upper portion, just below the clavicle, or at its lower portion in the axilla.

Ligation of the Axillary Artery Below the Clavicle.

Incision four inches in length from the summit of the coracoid process inward a short distance below the clavicle

(Fig. 309, *E*), or an incision three inches in length commencing at a point one-half an inch from the sterno-clavicular articulation and carried obliquely downward toward the axilla.

The skin and subcutaneous tissue having been divided the deep fascia is exposed and opened, or the axillary artery may be reached by following the intermuscular space between the sternal and clavicular fibres of the pectoralis major which leads upward toward the clavicle and to the pectoralis minor; or the fibres of the pectoralis major being exposed are cut through and the costo-coracoid fascia is next torn through with a director, care being taken to avoid injury of the cephalic vein at the outer portion of the wound; the pectoralis minor is now seen, and after separating the cellular tissue with a director the axillary vein is seen crossing from the upper edge of the muscle to the clavicle; the

FIG. 318.



A. Incision for axillary artery in axilla. *B*. Incision for brachial artery. (STIMSON.)

vein almost completely covers the artery, which is exposed by drawing the vein inward. The needle is passed around the artery from within outward.

Ligation of the Axillary Artery in the Axilla.

Incision two and a half inches long, started at the upper part of the axilla and carried down the arm at the edge of

the coraco-brachialis muscle. (Fig. 318, *A.*) The skin only is divided in the first incision and the deep fascia is picked up and divided upon a director, and the fibres of the inner border of the coraco-brachialis muscle are exposed and held aside by a retractor, and the operator will see the median nerve, the musculo-cutaneous nerve, and the axillary artery. To the inner side of the artery are the axillary vein, ulnar

FIG. 319.



Relations of right axillary artery in axilla. (ESMARCH.)

and internal cutaneous nerves. The needle should be passed around the artery from the vein toward the coraco-brachialis muscle.

LIGATION OF THE BRACHIAL ARTERY.

Incision at the middle of the arm three inches long on a line corresponding to the inner edge of the biceps muscle. (Fig. 318, *B.*) The skin and cellular tissue being divided, care being taken not to injure the basilic vein, which should be drawn posteriorly, the deep fascia is next cut through and the fibres of the biceps muscle are exposed (Fig. 320);

this should be drawn forward and the sheath of the vessels enclosing the artery, veins, and median nerve is exposed;

FIG. 320.



Relations of right brachial artery at middle of arm. (ESMARCH.)

this is opened, the median nerve is pressed aside and the artery is separated from its veins and the needle is passed from the side of the nerve around the vessel.

FIG. 321.

Ligation of the brachial artery at bend of elbow. (BRYANT.)

In ligating the brachial artery the occasional high division of the vessel must be borne in mind.

Ligation of Brachial Artery at Bend of Elbow.

Incision two inches in length, along the inner border of the tendon of the biceps muscle. Divide the skin, superficial fascia, and the bicipital aponeurosis, under which the artery will be exposed, resting upon the brachialis anticus muscle. (Fig. 321.) The median nerve is to the inner side and some distance from the artery. The needle should be passed around the vessel, after isolating the veins, from within outward.

Ligation of the Radial Artery.

The radial artery extends in a straight line from a point half an inch below the centre of the fold of the elbow to the inner side of the styloid process of the radius.

The radial artery may be tied at its upper, middle, or lower third, or at the root of the thumb.

Ligation of the Radial Artery in the Upper Third of the Forearm.

Incision for the radial artery at its upper third is two and a half inches in length on a line drawn from the middle of the bend of the elbow to the ulnar side of the styloid process of the radius; the incision should begin one and a half inches below the bend of the elbow. (Fig. 322, *A*.) Divide the skin and superficial fascia, avoiding the superficial veins. When the deep fascia is exposed find the edge of the supinator longus muscle and divide the aponeurosis along its ulnar side and expose the fibres of the pronator radii teres muscle. The vessel lies in the interspace between these muscles surrounded by adipose tissue, and upon being exposed the veins should be isolated and the needle passed from without inward. The nerve lies so far external to the artery that it is not often exposed in the operation. (Fig. 323.)

Ligation of the Radial Artery in the Middle Third of the Forearm.

Incision two inches in length, following the same line as that for the upper third of the artery. After dividing the

skin, superficial and deep fascia, the artery is found in the interspace between the flexor carpi radialis on the inner side

FIG. 322.



Line of incision for—*A*. Radial artery in upper third. *B*. Radial artery in lower third. *C*. Ulnar artery in upper third. *D*. Ulnar artery in lower third. (STIMSON.)

FIG. 323.

Relations of right radial artery in the upper third of the forearm. (ESMARCH.)

FIG. 324.

Relations of right radial artery above the wrist. (ESMARCH.)

and the supinator longus on the outer side; the radial nerve at this part of the arm is in close relation with the vessel to the radial side, and the needle should be passed around the artery from without inward.

Ligation of the Radial Artery in the Lower Third of the Forearm.

Incision two inches in length following the same line (Fig. 322, *B*), ending one inch above the wrist. The skin, superficial fascia, and deep fascia being divided, the artery will be found between the tendon of the flexor carpi radialis on the inner side and the tendon of the supinator longus on the outer side. (Fig. 324.) The veins being separated the needle may be passed in either direction.

Ligation of the Radial Artery at the Root of the Thumb.

The radial artery may also be tied at the root of the thumb.

Incision one inch in length between the tendons of the extensor ossis metacarpi pollicis and extensor primi internodii pollicis on the outer side, and the tendon of the extensor secundi internodii pollicis on the inner side. The skin and superficial fascia being divided and the radial vein being displaced, the deep fascia is opened and the artery is exposed at the bottom of the wound; the needle may be passed in either direction.

Ligation of the Ulnar Artery.

The ulnar artery is tied at the junction of the upper and middle third of the forearm and at the lower third.

Ligation of the Ulnar Artery at the Junction of the Upper and Middle Thirds of the Forearm.

Incision three inches in length, starting four inches below the internal condyle of the humerus, on a line passing from the internal condyle of the humerus to the outer border of the pisiform bone. (Fig. 322, *C*.) Divide the skin and superficial fascia, and when the deep fascia has been exposed the interspace between the flexor carpi ulnaris and the flexor sublimis digitorum appears, enter this interspace and raise

the flexor sublimis digitorum and work transversely across the arm, and the artery will be found resting upon the deep flexor, with the ulnar nerve to the ulnar side. The needle should be passed from the nerve around the artery. (Fig. 325.)

FIG. 325.



Relations of the right ulnar artery at upper third of forearm. (ESMARCH.)

Ligation of the Ulnar Artery in the Lower Third of the Forearm.

Incision two inches in length a little to the radial side of the tendon of the flexor carpi ulnaris, which is attached to the pisiform bone, ending an inch above the wrist. (Fig. 322, *D.*) Divide the skin and superficial fascia and open the deep fascia, and the artery will be exposed, with accompanying veins, between the tendons of the flexor carpi ulnaris and flexor sublimis digitorum, the ulnar nerve being to the ulnar side of the vessel. The needle should be passed from within outward to avoid the nerve. (Fig. 326.)

FIG. 326.



Relations of right ulnar artery above the wrist. (ESMARCH.)

Ligation of the Interosseous Artery.

Incision similar to that employed in the ligation of the ulnar artery in its upper third.

Ligation of the Abdominal Aorta.

Incision in the linea alba from a point three inches above the umbilicus to a point three inches below it. The superficial structures being divided the peritoneum is opened upon a director, and the intestines are pressed aside and the aorta is exposed covered by peritoneum, with the filaments of the sympathetic nerve resting upon it, and the vena cava to the right side. Tear through the peritoneum and pass the needle from left to right around the vessel. After tying the ligature the ends should be cut short, and the external wound should be closed as in the ordinary laparotomy wound.

The vessel may also be exposed by an incision along the anterior border of the quadratus lumborum muscle, from the last rib to the crest of the ilium. The skin, lumbar muscles, and fascia transversalis being divided, the wound is held open with blunt hooks, so that the retro-peritoneal space is exposed and the aorta brought into view. The vessel being separated from the vena cava and nerves, the needle is passed around it and the ligature applied.

Ligation of the Common Iliac Artery.

The aorta divides into the two common iliac arteries on the left side of the fourth lumbar vertebra, and these arteries are usually about two inches in length, and bifurcate opposite the sacro-iliac synchondrosis to form the internal and external iliac arteries; the length of the common iliac artery, however, may vary considerably, being three or four inches in length in some cases.

Incision for ligation of the common iliac artery is four to six inches in length, beginning one-half inch above the middle

of Poupart's ligament, and is carried outward curving upward after passing the anterior superior spine of the ilium. (Fig. 327, *A*.)

Divide the skin, superficial fascia and aponeurosis of the external oblique muscle, and then divide the fibres of the internal oblique and transversalis muscles upon a director, and expose the transversalis fascia. This is opened at the lower

FIG. 327.



Line of incision for—*A*, common iliac artery. *B*, external iliac artery. *C*, femoral artery in Scarpa's triangle. (STIMSON.)

part of the wound, and the finger is introduced and the peritoneum is pressed back; the opening in the transversalis fascia is next enlarged, and the peritoneum is carefully drawn inward and upward with the fingers toward the inner edge of the wound. The operator next feels for the external iliac artery, and passes the finger along this until the common iliac artery is reached. The loose cellular tissue in which it is imbedded is next separated, and the needle is passed from within outward, to avoid the common iliac vein (Fig. 328),

which on the left side lies on the inner side of the artery, and on the right side it lies behind the artery. The ureter generally remains attached to the peritoneum; if not, it is seen crossing the bifurcation of the common iliac with the

FIG. 328.

Ligation of the common iliac artery. (LISTON.)

genito-crural nerve, and care should be taken to avoid injury of these structures if present.

The common iliac artery may also be exposed and tied by an incision made over the artery through the peritoneal cavity; the vessel being tied, the ends of the ligature are cut short, and the external wound is closed in the same manner as that resulting from the exposure of the abdominal aorta by incision through the peritoneum.

LIGATION OF THE INTERNAL ILIAC ARTERY.

Incision in the same line as for the common iliac artery, but it need not be quite so long. (Fig. 327, *A*.) The peri-

toneum being exposed, it is pushed upward and inward, and the internal iliac artery is exposed. The vessel is carefully isolated from the vein, which lies behind and on the inner side, and the needle is passed from within outward.

Ligation of the External Iliac Artery.

Incision three or four inches in length, half an inch above the middle of Poupart's ligament, made at first parallel to it and then curved upward. (Fig. 327, *B*.) The tissues of the abdominal wall being divided and the peri-

Fig. 329.



Relations of the right external iliac artery. (Esmarch.)

toneum exposed, it is pushed upward and inward in the same manner as for exposure of the common iliac artery. The artery lies at the inner border of the psoas muscle, the vein on its inner side, and the anterior crural nerve covered by the iliac fascia on the outer side; the genito-crural nerve passes obliquely across the artery. (Fig. 329.) The needle should be passed from within outward.

Ligation of the Gluteal Artery.

Incision three or four inches in length, from the posterior superior spinous process of the ilium to a point midway between the tuber ischii and the great trochanter. (Fig. 330, *A*.) After division of the skin and fascia, the fibres

Fig. 330.



Line for—*A*, gluteal artery. *B*, sciatic and internal pudic artery. (Stimson.)

of the gluteus maximus muscle are separated and held apart, and the deep fascia is divided, and the artery is sought for above the pyriformis muscle at the upper border of the great sacro-sciatic notch. It is accompanied by large veins, injury to which should be avoided in exposing the artery and passing the needle.

Ligation of the Sciatic and Internal Pudic Arteries.

Incision three or four inches in length, a little lower than that employed for exposure of the gluteal artery. (Fig. 330,

one-fourth of an inch above Poupart's ligament, and extending ten inches downward. Divide the skin and superficial fascia and the deep fascia and expose the sheath of the vessels; open this one-half an inch below Poupart's ligament and isolate the femoral artery from the femoral vein which lies to the inner side; the anterior crural nerve lies to the outer side. Pass the needle from within outward.

Ligation of the Femoral Artery at the Apex of Scarpa's Triangle.

Incision three inches long, the centre of which should be a little above the point where the sartorius muscle crosses a line drawn from the middle of Poupart's ligament to the inner condyle of the femur. (Fig. 332.) Divide the skin,

FIG. 332.



Lines of incision for the femoral artery. (STIMSON.)

superficial fascia and deep fascia, avoiding the internal saphenous vein, and expose the edge of the sartorius muscle, which may be recognized by the direction of its fibres. This muscle is drawn outward and the sheath of the vessels is exposed and opened; the vein lies on the inner side and somewhat behind the artery and the long saphenous nerve is on the outer side. (Fig. 333.) Pass the needle from within outward.

B.) Divide the skin, superficial fascia and fibres of the gluteus maximus muscle and deep fascia, and search for the vessels as they leave the great sciatic notch at the lower edge of the pyriformis muscle. The internal pudic artery enters the pelvis through the lesser sciatic notch, lying on the inner side of the sciatic artery during its passage over the spine of the ischium. The vessels are isolated and the needle is passed so as to avoid injury of the veins.

Ligation of the Femoral Artery.

The femoral artery may be ligated just below Poupart's ligament, at the apex of Scarpa's triangle, at the middle of the thigh, or in Hunter's canal.

Ligation of the Femoral Artery below Poupart's Ligament.

Incision beginning midway between the anterior superior spinous process of the ilium and the symphysis pubis,

Fig. 331.



Relations of the right femoral artery below Poupart's ligament. (Esmarch.)

one-fourth of an inch above Poupart's ligament, and extending ten inches downward. Divide the skin and superficial fascia and the deep fascia and expose the sheath of the vessels; open this one-half an inch below Poupart's ligament and isolate the femoral artery from the femoral vein which lies to the inner side; the anterior crural nerve lies to the outer side. Pass the needle from within outward.

Ligation of the Femoral Artery at the Apex of Scarpa's Triangle.

Incision three inches long, the centre of which should be a little above the point where the sartorius muscle crosses a line drawn from the middle of Poupart's ligament to the inner condyle of the femur. (Fig. 332.) Divide the skin,

FIG. 332.



Lines of incision for the femoral artery. (STIMSON.)

superficial fascia and deep fascia, avoiding the internal saphenous vein, and expose the edge of the sartorius muscle, which may be recognized by the direction of its fibres. This muscle is drawn outward and the sheath of the vessels is exposed and opened; the vein lies on the inner side and somewhat behind the artery and the long saphenous nerve is on the outer side. (Fig. 333.) Pass the needle from within outward.

Ligation of the Femoral Artery in the Middle of the Thigh.

Incision in the line above mentioned, its centre being a little above the middle of the thigh. Divide the skin, super-

Fig. 333.



Relations of right femoral artery at the apex of Scarpa's triangle. (Esmarch.)

Fig. 334.



Relations of the right femoral artery in the middle of the thigh. (Esmarch.)

ficial and deep fascia and expose the sartorius muscle which is drawn outward after the leg has been flexed; the sheath of the vessels is exposed and opened; the long saphenous nerve lies upon the artery and the femoral vein lies behind the artery; the saphenous vein lies more superficially and internal to the vessel. Pass the needle from within outward. (Fig. 334.)

Ligation of the Femoral Artery in Hunter's Canal.

Incision three inches in length along the tendon of the adductor magnus, the centre of which is at the junction of the lower and middle thirds of the thigh. (Fig. 332.) Divide the skin, superficial fascia and deep fascia, care being

taken not to injure the internal saphenous vein, which should be displaced and expose the sartorius muscle, which should be displaced downward and expose the aponeurosis which forms the anterior wall of the vascular canal; this should be opened upon a director, and the artery is uncovered and should be separated from the vein, which lies upon the outer side. The needle is passed from without inward.

Ligation of the Popliteal Artery.

Incision three or four inches in length, along the external border of the semi-membranosus muscle. Divide the

Fig. 335.



Relations of the right popliteal artery. (Esmarch.)

skin and superficial fascia, taking care not to injure the saphenous vein, and open the deep fascia. The edges of the

wound being held apart the adipose tissue is broken up with a director, and the internal popliteal nerve will be first

Fig. 336.



Ligation of popliteal artery. (Smith.)

exposed, and next the vein—both external to the artery. (Fig. 335.) The artery is isolated and the needle is passed from without inward. (Fig. 336.)

Ligation of the Anterior Tibial Artery.

The anterior tibial artery may be tied in the upper, middle, and lower thirds of the leg; the general direction of the artery corresponds with a line drawn from the middle of the space between the head of the fibula and the tubercle of the tibia to the middle of the anterior intermalleolar space.

Ligation of the Anterior Tibial Artery in the Upper Third of the Leg.

Incision two and a half to three inches in length, one and one-fourth inches external to the spine of the tibia. Divide the skin and superficial fascia, and when the deep fascia is exposed open it on a line corresponding to the intermuscular space between the tibialis anticus and the

extensor longus digitorum muscles. Separate the muscles and work down in this interspace, and the artery will be found with a vein on either side of it, and the anterior tibial nerve externally. (Fig. 337.) The needle should be passed from without inward, after isolating the veins.

Fig. 337.



Ligation of the anterior tibial artery at its upper third. (Stimson.)

Ligation of the Anterior Tibial Artery at its Middle Third.

Incision three inches in length in the same line as that for the upper portion of the vessel. After dividing the skin, superficial and deep fascia, the interspace between the tibialis anticus and the extensor longus digitorum muscles is opened and a third muscle comes in view, the extensor proprius pollicis The artery lies between the extensor proprius pollicis and the tibialis anticus muscles, and the anterior tibial nerve is to the outer side. The veins should be isolated and the needle should be passed from without inward.

Ligation of the Anterior Tibial Artery in its Lower Third.

Incision two inches in length, beginning three inches above the ankle-joint on the line of the artery. Divide the skin, superficial and deep fascia, and seek for the tendon of the extensor proprius pollicis muscle, the second tendon from the tibia. The artery is found in the interspace between this tendon and the tendon of the extensor longus digitorum muscle, the nerve being to the outer side. The veins are isolated from the artery, and the needle is passed from without inward.

Ligation of the Dorsalis Pedis Artery.

Incision one inch in length on a line drawn from the middle of the anterior inter-malleolar space to a point midway between the extremities of the first two metatarsal bones or along the outer border of the tendon of the extensor proprius pollicis. Divide the skin, superficial and deep fascia, and

Fig. 338.



Ligation of the dorsalis pedis artery. (Bryant.)

the artery will be found lying next to the inner tendon of the short extensor muscle of the toes. (Fig. 338.) The nerve is to the outer side. After separating the veins the needle is passed from without inward.

Ligation of the Posterior Tibial Artery.

The course of the posterior tibial artery is indicated by a line drawn from the middle of the popliteal space to a point

midway between the tendo Achillis and the internal malleolus of the tibia.

The posterior tibial artery may be ligated in its upper, middle, and lower thirds.

FIG. 339.



Lines of incision for the posterior tibial artery. (STIMSON.)

Ligation of the Posterior Tibial Artery at its Upper Third.

Incision three inches and a half in length, one-half inch from the inner edge of the tibia, beginning two inches from the upper edge of the tibia. (Fig. 339.) Divide the skin and superficial fascia, avoiding large superficial veins; next open the deep fascia and detach the origin of the soleus muscle from the tibia, and on raising it, its under surface will present a white shining sheath of tendinous material, beneath which will be seen a layer of fascia covering the tibialis posticus muscle. If search is made toward the middle of the leg, the artery will be found covered by the intermuscular fascia, the nerve being to the outer side. The needle is passed from without inward after the veins have been separated from the artery (Fig. 340).

Ligation of the Posterior Tibial Artery at its Middle Third.

Incision two and a half inches in length, parallel with the inner edge of the tibia and half an inch from its border.

Divide the skin, superficial and deep fascia, and the inner edge of the soleus will be exposed; press this outward and

Fig. 340.



Relations of the right posterior tibial artery in its upper third. (Esmarch.)

the artery with its veins will be exposed, also the posterior tibial nerve to the outer side. Pass the needle from without inward after separating the veins.

Ligation of the Posterior Tibial Artery Behind the Inner Malleolus.

Incision a curved one two inches in length, midway between the tendo Achillis and the internal malleolus. (Fig. 339.) Divide the skin and superficial fascia and lift the deep fascia upon a director and open it freely and the artery will be exposed with the tendons of the tibialis posticus and flexor longus digitorum muscles on the inner side and the posterior tibial nerve and the tendon of the flexor longus

FIG. 341.



Ligation of the posterior tibial artery behind inner malleolus. (BRYANT.)

pollicis muscle on the outer side. (Fig. 341.) After separating the veins from the artery the needle should be passed from without inward.

PART VI.

AMPUTATIONS.

The term amputation is now generally applied to the removal of a limb, and this may be removed through the bones, when the operation is spoken of as an amputation in the continuity of the limb; or it may be removed through its joints, and is then known as an amputation in the contiguity or a disarticulation.

Methods of Amputating.

Amputations may be performed by the *circular*, *flap*, *oval*, and *elliptical* methods; the *modified circular* operation, and *Teale's method* by rectangular flaps, are also employed.

Fig. 342.

Amputation by circular method. (Druitt.)

Circular Method.

In performing an amputation by this method the incision of the skin is made at a distance below the point where the bone is to be divided. An assistant grasps the limb and draws the skin evenly and firmly toward the root of the part and the surgeon passes the heel of the knife well into the tissues and makes a circular sweep around the limb and completes the division of the skin and cellular tissue with one motion of the knife. (Fig. 342.)

In some cases a cutaneous sleeve consisting of the skin and cellular tissue is dissected up and turned back, and sometimes it may be necessary to make a slit on one side of the flap to allow it to be turned up.

The second incision in an amputation by the circular method consists, after retraction of the skin, in making a circular cut through all of the tissues down to the bone. (Fig. 343.)

Fig. 343.

Division of muscles in circular amputation. (Smith.)

The third step in an amputation by the circular method consists, after retracting the skin and muscles and holding them back by a retractor, in the division of the bone with a saw.

Flap Method.

This method of amputating is susceptible of many variations. There may be one or two flaps of equal or unequal length; the flaps may be cut antero-posteriorly, laterally, or obliquely. (Fig. 344.) They may be made by transfixing the limb and cutting outward, or they may be cut from without inward, or they may be made to include the whole thickness of the tissues down to the bone, or merely the skin and superficial fascia, the deeper structures being divided by a circular incision. The flaps may have a curved outline

motion and then cuts obliquely forward until all the tissues are divided. The flap being turned up, he reënters his knife at the same point and passes it on the other side of the bone or bones and cuts the second flap in the same manner. (Fig. 345.) A retractor is next applied and the bone is divided with a saw.

The Oval Method.

The oval amputation is really a circular one in which the cuff of skin has been slit at one side and the angles rounded off. This is the form of amputation frequently performed at the metacarpo-phalangeal and metatarso-phalangeal joints, and is one of the methods of amputation at the shoulder-joint.

Elliptical Method.

This is a form of the oval method of amputation which is employed in amputations at the knee- and elbow-joints, the incision forming an ellipse coming below the joint on the front or outside of the limb, the resulting flap being folded upon itself.

Fig. 346.

Modified circular amputation. (Skey.)

Modified Circular Method.

In this method of amputation two oval skin flaps, antero-posterior or lateral, are turned up, and the muscles are next divided by a circular sweep of the knife down to the bone (Fig. 346).

Teale's Method by Rectangular Flaps.

In this method of amputation, two flaps are made of unequal length; the incisions are so planned that the shorter flap contains the main vessel or vessels. The flaps are cut of equal width and the length of the long flap should be one-half of the circumference of the limb at the point where the bone is to be divided; the length of the short flap should be one-eighth of the circumference of the limb. The flaps are cut from without inward, and embrace all of the tissues of the limb down to the bone. After the flaps have been dissected up, the bone is divided with a saw, and the long flap is folded over and sutured to the short flap (Fig. 347).

Fig. 347.



Teale's method of amputation. (Bryant.)

The disadvantage of this method of amputation is that in muscular limbs it requires the bone to be divided at a higher point than would otherwise be necessary.

Periosteal Flaps.

In any of the methods of amputation previously described the periosteum may be dissected up in two flaps attached to the muscles, or pushed up as a sleeve by means of a director or periosteotome before the bone is sawed. This procedure is most easily accomplished in young subjects. When these flaps are made and they are brought together, the periosteum covers the cut surface of the bone, to which it soon forms adhesions.

Instruments Required for Amputations.

The instruments required for amputations are knives of various shapes and sizes, saws, dissecting forceps, bone forceps, artery forceps, tenacula, hæmostatic forceps, scissors, periosteotome, tourniquets, Esmarch's bandage and strap, retractors, ligatures, sutures, and suture needles.

Amputating Knives.

The knives required for amputations vary according to the method of amputation and the part to be amputated.

Fig. 348.

Scalpel.

Fig. 349.

Straight bistoury.

In certain amputations a scalpel (Fig. 348) or straight bistoury may be used (Fig. 349), while in other cases the employment of amputating knives of various sizes will be found more satisfactory. For amputations of the thigh a knife with a blade of eight or nine inches is generally em-

ployed, and for smaller limbs a knife with a blade of six or seven inches in length; double-edged catlins are employed in

Fig. 350.

Amputating knife and catlin.

amputations of the leg and forearm to divide the inter-osseous tissues before applying the saw. The amputating knives now employed are constructed with solid metal handles so that they can be rendered thoroughly aseptic by immersion in boiling water before being used.

Amputating Saws.

Several kinds of amputating saws are in general use; one with a blade ten inches long by two and a half inches wide,

Fig 351.



Amputating saw.

Fig. 352.



Small amputating saw.

with a heavy back to give it additional firmness, is a very good variety of saw (Fig. 351). For amputations about the foot or hand a narrow saw with a movable back will be

found very convenient. (Fig. 352.) A bow saw with a metallic handle and a reversible blade is a very useful variety of saw, as it can be used either in amputations or in

Fig. 353.

Amputating saw with reversible blade.

excisions, and, being constructed entirely of metal, it can be easily rendered aseptic. (Fig. 353.)

Bone Forceps, or Cutting Pliers.

These instruments are used in smoothing off any rough edges of bone left after the use of the saw, or for the division of the small bones in amputations of the fingers and toes.

Fig. 354.

Bone forceps, or cutting pliers.

The forceps should be from ten to twelve inches in length, with blades from one to one and a half inches in length. (Fig. 354.)

Periosteotome.

The periosteotome, or raspatory, is employed for dissecting up a flap of periosteum, which, after sawing the bone, is drawn down over the sawed end of the bone. (Fig. 355.)

Fig. 355.



Periosteotome.

Artery Forceps and Tenacula.

These instruments are used for taking up the vessels, and one of the best forms of artery forceps is that known as the double-spring artery forceps. (Fig. 188, p. 252.) Tenacula

Fig. 356.



Ligature applied. (Esmarch.)

are also employed for the same purpose. *Hæmostatic forceps* will also be found most useful in cases of amputation, for the purpose of arrest of hemorrhage from small vessels after the tourniquet has been removed, the vessels being secured by ligatures before the hæmostatic forceps are removed.

Retractors.

These consist of pieces of muslin six or eight inches in width, one end of which is split into two or three tails; the former variety of retractor is employed where one bone is divided, as in amputations of the arm and thigh, and the latter in cases where two bones are divided, as in amputations of the forearm and leg. (Fig. 356.)

Ligatures.

The best material to employ for the ligature of vessels is juniper or chromicized catgut, the preparation of which has been described.

Sutures.

The materials employed for sutures in cases of amputation may be silkworm-gut, catgut, silk, or silver wire; deep or buried sutures of catgut in bringing together the edges of the

FIG. 357.

Deep or buried sutures of muscles.
(ESMARCH.)

FIG. 358.



Sutures of skin.
(ESMARCH.)

periosteal flaps, muscles, and fascia, are often employed with advantage in amputations (Fig. 357), the skin flaps being brought together with interrupted or continuous sutures of silk, catgut, silkworm-gut, or silver wire. (Fig. 358.)

Tourniquets.

For the control of hemorrhage during the amputation the Esmarch apparatus (Fig. 186), or Petit's tourniquet (Fig. 179) is employed; or the employment of both at the same time will often be found most satisfactory. The Esmarch bandage and tube being applied, after the removal of the bandage the tourniquet of Petit is loosely applied at a higher point, and after the main vessels have been secured the elastic strap is removed and the tourniquet is screwed down and controls the bleeding until the smaller vessels have been secured by ligatures.

Details of an Amputation.

The following are the steps of an amputation of the lower part of the thigh:

The skin is first thoroughly cleansed by rubbing it with turpentine and soap and water and is then washed with an antiseptic solution either of carbolic acid 1 : 40 or bichloride of mercury 1 : 2000. Provision is next made to prevent the loss of blood during the operation by the application of Esmarch's bandage and tube; the bandage being removed a tourniquet is placed over the femoral artery in Scarpa's triangle and loosely secured. The limb is again washed with bichloride solution. The instruments having been previously placed in an antiseptic solution, a rubber cloth covered with towels wrung out in a bichloride solution is placed under the limb. The variety of amputation having been decided upon, the flaps are cut and the muscles are divided down to the bone; the periosteum being dissected up, a two-tailed retractor is applied and the tissues are held back by an assistant while the surgeon divides the bone with the saw. When the bone has been divided the retractor is removed and the surface of the wound is irrigated with a 1 : 2000 bichloride solution. The femoral artery and vein are next found and secured with ligatures, and any branches which can be found are also secured. The elastic strap is removed after screwing down the tourniquet, and by letting up the

pressure on this, smaller vessels which bleed are picked up with artery forceps or hæmostatic forceps and secured. After all bleeding has been controlled the tourniquet is removed and the wound is again thoroughly irrigated with a 1 : 2000 bichloride solution. If there is much oozing from smaller vessels this solution should be as hot as the hands of the operator can comfortably stand, which will act promptly in controlling this variety of bleeding. The periosteal flaps, if they have been made, are brought together by two or three catgut sutures, and a drainage-tube is next introduced or two short tubes are introduced at either extremity of the wound

FIG. 359.



Stump showing application of sutures and drainage-tubes. (SMITH.)

and secured by sutures or safety-pins; the muscles should next be brought together by a few deep or buried sutures of catgut, and the skin flaps should then be brought into apposition by a number of interrupted sutures. The inner surface of the stump is next irrigated by a stream of bichloride solution introduced through the drainage-tube, and the surface of the stump is washed with the same solution; a piece of protective is next placed over the line of the wound and over this is placed a moist carbolized, bichloride, or

iodoform gauze dressing, and over this a number of layers of dry gauze; this is next covered by rubber tissue and a number of layers of bichloride cotton, or if the dry method of dressing is preferred the rubber tissue is omitted and a number of layers of bichloride cotton are laid over the gauze dressing, and the whole dressing is held in place by a recurrent bandage of the stump.

Re-dressing of Amputations.

The first dressing of an amputation, if strict antiseptic precautions have been observed at the time of operation, need not, as a rule, be made for a week or ten days, except in cases where the oozing is so profuse as to soak the dressings, or where consecutive hemorrhage has occurred, or the patient's condition shows that the wound is not running an aseptic course. The re-dressing of a stump can be accomplished without pain to the patient if the surgeon and his assistants are careful in their manipulations.

The dressings to be applied, the solutions for irrigation, and the instruments required should be prepared and at hand before the stump is exposed. The surgeon and his assistants should wash their hands carefully, and then dip them in a 1 : 2000 bichloride solution. The bandage retaining the dressings to the stump should be divided with bandage scissors without lifting the stump from the pillow upon which it rests. After the bandage has been divided and turned aside, the gauze dressing is next unfolded and turned down; an assistant now slips his hands under the stump and gently raises it from the dressings, and at the same time a rubber cloth covered with towels which have been wrung out in a 1 : 2000 bichloride solution is slipped under the stump and the soiled dressings are removed. The protective covering the incision is next removed and the surface of the stump is irrigated with a 1 : 2000 bichloride solution; the drainage-tubes are next examined and the cavity of the stump is irrigated with the bichloride solution through the tubes by means of a syringe or an irrigating apparatus.

If the wound is aseptic and there seems to be no further indication for the use of the drainage-tubes they may be removed and the track of the tube should be washed out with the antiseptic solution by the syringe or irrigator. The sutures are next examined and if the wound is firmly healed alternate sutures may be removed; if catgut or silkworm-gut sutures have been used they need not be disturbed at this dressing, and their removal may be postponed until a subsequent dressing.

The wound should next be covered with a piece of protective, and a gauze dressing should be applied consisting of a number of layers of bichloride cotton, and the dressings should be held in place by a recurrent bandage of the stump. In holding the stump the assistant should hold it firmly to prevent muscular spasm, and after the dressings have been secured it should be placed upon a clean pillow prepared for its reception. The same procedures are adopted at subsequent dressings, and if the wound has run an aseptic course, two or three dressings, at most, will be required.

Special Amputations.

Amputations of the Hand.

Amputations of the Fingers.

The fingers may be amputated in the continuity of the phalanges or in their contiguity, and, as a rule, as it is important to save as much as possible of the finger, the former method is generally to be employed instead of disarticulation at a higher point. The incisions should be so planned that the cicatrix does not occupy the plantar surface; the larger flap should, therefore, be taken from the palmar aspect of the finger. In amputating the phalanges of the fingers in their *continuity* the circular method (Fig. 363, *B*) or a short dorsal flap and a long palmar flap may be employed. In *disarticulating* a phalanx it is best to enter the joint with a narrow knife from the dorsal side, and after

having carried it through the joint, to cut a long palmar flap, keeping close to the bone. (Fig. 360.) In locating

FIG. 360.



Amputation of finger: long palmar flap. (ERICHSEN.)

the position of the phalangeal joints it is well to remember that the prominence of the knuckle, when the finger is flexed is formed entirely of the head of the proximal, and

FIG. 361.

Phalanges flexed.

FIG. 362.



Guides to articulations of the finger. (SMITH.)

not of the base of the distal phalanx (Fig. 361), and also that the folds on the palmar surface of the finger do not correspond exactly to the joints. (Fig. 362.)

Amputation of the Finger through Metacarpo-phalangeal Articulation.

In this variety of amputation an incision is made from a point of the dorsal surface of the metacarpal bone a quarter of an inch above the articulation, which is carried through the interdigital web and back upon the palmar sur-

FIG. 363.



A. Disarticulation of phalanx, palmar flap. *B*. Amputation in continuity, circular. *C*. Metacarpo-phalangeal disarticulation. *D*. Amputation of metacarpal bone in continuity. *E*. Disarticulation of little finger. *F*. Disarticulation of fifth metacarpal bone. *G*. Amputation at the wrist, circular. *H*. Amputation at the wrist. (STIMSON.)

face to a point a quarter of an inch above the flexor fold (Fig. 363, *C*). A similar incision beginning and ending at the same points is made upon the opposite side of the finger. The flaps are dissected back, and the lateral ligaments, tendons, and remainder of the capsule are divided. The

finger may also be amputated at the metacarpo-phalangeal joint by making an incision on one side and dissecting the flap back to the joint, then dividing the lateral ligament, opening the joint and carrying the knife across this, dividing the tendons and lateral ligament on the other side and cutting a flap from within outward.

Removal of the head of the metacarpal bone if desired may be accomplished by the use of cutting pliers (Fig. 364),

FIG. 364.

Removal of head of metacarpal bone. (SKEY.)

but, as a rule, this procedure is not to be recommended, for, although the deformity is diminished, the strength of the hand is also diminished.

In amputating the *little* and *index* fingers a full lateral flap may be cut on the free side and an incision is next carried across the palmar surface to the angle of the web and thence back to the joint, which is opened and the disarticulation is effected. (Fig. 363, *E.*)

In amputations of the finger at the phalangeal joints or at the metacarpo-phalangeal joints two vessels usually re-

quire ligaturing, and after these are secured a catgut drain or a small drainage-tube is introduced and the flaps are brought together by a few interrupted sutures.

Amputations of the Metacarpal Bones.

In amputating the metacarpal bones it is advisable to leave the carpal ends of the bones to avoid opening the wrist-joint, except in the case of the first and fifth metacarpal bones, which do not communicate with the others and with the synovial sacs.

The incisions for the removal of the metacarpal bones are the same as for the removal of a finger at the metacarpo-phalangeal joint, the incision being prolonged backward as far as necessary over the dorsal surface of the bone. (Fig. 363, *D.*) After the metacarpal bone has been bared for a sufficient distance, it is cut through with bone-pliers or disarticulated, and the distal end is raised from its bed and carefully separated from the soft parts, care being taken to avoid injury of the structures of the palm of the hand.

In amputating the fifth metacarpal bone the incision should be made along the inner border of the hand and carried down to the bone between the skin and the abductor minimi digitii muscle. (Fig. 365.) The lower end of the incision passes over the knuckle to the web of the finger and backward under the palmar surface to join the first incision.

Amputation of the entire thumb with its metacarpal bone is effected by making an oval flap from the palmar surface; in case of the left thumb the joint may be opened by an oblique incision on the dorsal surface of the hand, beginning a little in front of the joint and being carried down to the web between the thumb and forefinger; the palmar flap is then made by

FIG. 365.



Incision for removal of the fifth metacarpal bone. (SMITH.)

thrusting the knife upward to its point of entrance and cutting downward and outward. In amputating the right thumb with its metacarpal bone it is better to make the palmar flap first by transfixion, the dorsal flap being made subsequently.

Amputation of the hand at the carpo-metacarpal joint is occasionally performed, or between the rows of carpal bones, but is not as a rule to be recommended, as the carpal bones are apt subsequently to become diseased and require removal, so that it is better to amputate at the radio-carpal joint.

Amputations at the Wrist.

Circular Method.

The skin of the forearm near the wrist being retracted by an assistant, a circular incision of the skin and cellular

Fig. 366.



Amputation at the wrist. Erichsen.

tissue is made half an inch below the point of the styloid process of the radius. (Fig. 363, *G*.) The skin and cellular tissue are next dissected back as far as the joint, which is opened and the disarticulation is completed.

Antero-posterior Flap Method.

This method is also employed in amputations at the wrist-joint; an incision carried downward is made on the back of the hand from one styloid process to the other; the hand being flexed the tendons are divided and the joint opened, and the palmar flap, which should extend as far as the base of the metacarpal bones, is cut from within outward. (Fig. 366.) Amputation at the wrist is sometimes done by cutting a single flap from the palm, the joint being opened by a transverse incision on the back of the hand from one styloid process to the other.

Lateral Flap Method.

This method (Fig. 363, *H*) is also sometimes employed in amputation at the wrist, and may be employed with advantage in cases of laceration of the hand, in which the injury to the tissues prevents the formation of the flaps used in the other methods of amputation.

Amputations of the Forearm.

The forearm may be amputated by the circular or flap methods, or by making rectangular flaps (Teale's method).

Circular Method.

At the lower portion of the forearm the circular method of amputation is to be preferred. A circular incision of the skin and cellular tissue is made and a cuff is dissected up, the muscles and interosseous membrane being cut through; a three-tailed retractor is next applied and the bones are divided with a saw.

Mixed Method.

Amputation of the forearm by the mixed method, which consists in first dissecting up two antero-posterior oval flaps of skin and cellular tissue and then dividing the muscles by a circular incision, is also a satisfactory operation. (Fig. 367.)

In amputations at the upper portion of the forearm, *antero-posterior*, or *lateral* flaps, cut from without inward or by transfixion, or *rectangular* flaps may be made with advantage.

FIG. 367.

Amputation of the forearm by mixed method. (BRYANT.)

The principal vessels requiring the application of ligatures in amputations of the forearm are the radial, ulnar, and interosseous arteries.

AMPUTATIONS AT THE ELBOW.

The methods of amputation employed at the elbow are the anterior flap, lateral flap, and circular.

Anterior Flap Method.

A flap three inches in length with its base parallel to and half an inch below the condyles of the humerus, is cut either by transfixion or from without inward. The joint is next opened and the lateral ligaments divided and the olecranon is exposed and the attachment of the triceps is separated and a posterior flap is cut from without inward, or from within outward a little below the line of the condyles. (Fig. 368, *A*.)

Lateral Flap Method.

In amputation at the elbow-joint lateral flaps may be employed, cut either from without inward or by transfixion. (Fig. 368, *B*.) An external flap three inches in length is made on the outer side of the arm, starting from a point a finger's breadth below the bend of the elbow, by transfixion

or by cutting from without inward; a shorter internal flap is next cut in the same manner, and the joint is opened and the disarticulation effected. (Fig. 369.)

Fig. 368.



Amputation at the elbow-joint. *A*. Anterior flap method. *B*. External flap method. *C*. Circular method. (Stimson.)

Fig. 369.



Lateral flap method of amputation at the elbow-joint. (Smith.)

Fig. 370.



Circular amputation at the elbow. (Smith.)

Circular Method.

An incision dividing the skin and cellular tissue is made around the limb three inches below the line of the condyles of the humerus (Fig. 368, *C*), the skin is dissected up and a circular incision made through the muscles, the joint is opened and the disarticulation effected. (Fig. 370.)

Amputations of the Arm.

The arm may be removed at any point below the attachment of the muscles at the axilla, by either the circular, flap, oval, or modified circular methods.

Circular Method.

This operation is usually employed in removing the arm in its lower third: a circular incision of the skin and muscles

Fig. 371.



Circular amputation of the arm.

is first made, and when the cuff has been dissected up, a circular division of the muscles is made, and after applying the retractor the bone is sawed through. (Fig. 371.)

Flap Method.

From the central position of the bone in the arm the flap method in amputating the arm is preferred by many operators. The arm being grasped by the hand the point of a medium-sized amputating knife is thrust through the arm so as to pass over the humerus and make its exit at a corresponding point in the skin on the opposite side; a flap of sufficient length is cut from within outward. The knife is next passed behind the bone and a posterior flap is cut in

the same manner (Fig. 372); the bone is next cleared of muscular tissue and the flaps are retracted and it is divided with a saw.

FIG. 372.

Amputation of the arm by flap operation. (BRYANT.)

Lateral flaps may be made in this amputation in the place of the antero-posterior flaps, and they are cut from within outward in the same manner.

Oval, or Modified Oval Method.

This method of amputating the arm is also employed with advantage. An oval flap of skin and cellular tissue is made and dissected up, and the muscular tissue is divided by a circular incision. Or two oval flaps of skin and cellular tissue are cut and dissected up, and the muscles are next divided by a circular sweep of the knife.

In all amputations of the arm it is well to remember the possibility of a high division of the brachial artery, and to see that the abnormal vessel is properly secured, if present.

In high amputations of the arm there is sometimes not room enough to apply Esmarch's strap or a tourniquet to the arm itself to control the hemorrhage during the operation, and in such cases the strap may be passed from the axilla around the outer end of the clavicle, as is done to control the bleeding during amputation at the shoulder-joint. (Fig. 373).

Fig. 171.



Esmarch's bandage applied in the amputation of the arm. SMITH.

AMPUTATIONS AT THE SHOULDER-JOINT.

Several methods of operation are employed in amputating at the shoulder-joint, such as the oval method, or Larrey's method, flap method, Lisfranc's or Dupuytren's method, and

Fig. 172.



Amputation at the shoulder-joint. *A.* Oval, or Larrey's method. *B.* Double flap, or Lisfranc's method. STIMSON.

Spence's method. (Fig. 374.) The control of the bleeding from the axillary artery during the operation is a matter of the first importance, and it may be arrested by pressure made upon the subclavian artery, as it crosses the first rib, with the thumb, or the padded handle of a large key, or by the fingers of an assistant grasping the axillary flap and compressing the vessel after the head of the bone has been disarticulated, or by the use of an elastic strap applied around the axilla and shoulder. (Fig. 373.)

Oval, or Larrey's Method.

In this method of amputation the point of the knife is entered just below the acromion process, and a deep incision

FIG. 375.

Amputation at the shoulder-joint by Larrey's method.

three inches in length is made down to the head of the bone in the axis of the arm ; from the middle of this incision two

others are made obliquely downward to the points where the anterior and posterior folds of the axilla end in the tissues of the arm; the latter incision should be only deep enough to divide the skin and superficial fascia. The flaps are then dissected up until the head of the bone is well exposed, and, after opening the capsule and dividing the muscles inserted into the neck and tuberosities of the humerus, which division may be facilitated by rotating the head of the bone outward and inward, the disarticulation is effected by adducting the elbow; the knife is next passed downward behind the bone and made to cut outward in the line of the cutaneous incisions—an assistant controlling the artery before it is divided, by grasping the axillary tissues behind the knife with his fingers.

Flap, or Dupuytren's Method.

In this method of amputation at the shoulder-joint the flaps may be cut either by transfixion, or from without in-

FIG. 376.

Amputation at the shoulder-joint, Dupuytren's method. (BRYANT.)

ward; the large flap embraces the greater part of the deltoid muscle, and the smaller or short flap is cut from the inside of the arm after the head of the bone has been disarticulated. When cut by transfixion, the point of the knife

should be entered an inch in front of the acromion process and pushed across the outer aspect of the head of the humerus, and should be brought out at the posterior fold of the axilla; the knife is made to cut downward until a large deltoid flap is formed. This flap is turned up, and the head of the bone is disarticulated; the knife being placed behind it, a short flap is cut out, keeping close to the bone so that the vessel is divided with the last cut of the knife. (Fig. 376.) An assistant should control the vessel by grasping the axillary tissues with his fingers behind the knife.

Double Flap, or Lisfranc's Method.

In this method of amputation at the shoulder-joint the point of the knife is entered at the outer side of the coracoid process, and is carried across the outer aspect of the head of the humerus and brought out a little below the posterior border of the acromion process, and a long flap is cut downward. This flap is turned up and the attachments of the head of the bone are divided and it is disarticulated. The knife is again entered behind the bone, and a long posterior flap is cut from within outward. (Fig. 374, *B.*)

FIG. 377.

Amputation at the shoulder-joint. Spence's method. (STIMSON.)

Spence's Method.

In this method of amputation at the shoulder-joint an incision is made down to the head of the humerus immediately in front of the coracoid process, and is continued downward through the clavicular fibres of the deltoid and pectoralis major muscles until the attachment of the latter to the humerus is reached. (Fig. 377.) The incision is now carried backward to the posterior fold of the axilla. A second incision, including only the skin and cellular tissue, is next made from the anterior portion of the first incision

across the inside of the arm to meet the incision on the outer part. The outer flap thus formed is turned up and the head of the bone is disarticulated, and the operation is completed by dividing the remaining tissues on the axillary aspect.

Many other methods of removing the arm at the shoulder-joint have been devised and employed, including the circular method.

Amputation above the Shoulder-joint.

This form of amputation consists in the removal of the arm with a part or the whole of the scapula and sometimes a portion of the clavicle.

As this form of amputation is required in cases in which the laceration of the parts has passed beyond the shoulder-joint, or in cases of growths involving the tissues beyond the joint, no definite rule can be laid down for the incisions; the only rule being as far as possible to make the incisions in such a manner that the least possible amount of skin is sacrificed, so that a sufficient covering for the wound can be obtained.

Amputations of the Foot.

Amputations of the Toes.

The phalanges of the toes may be removed in the same manner as those of the fingers. It is better to amputate at the metatarso-phalangeal articulations than to attempt to remove them at the joints in front of this articulation, except in the case of the great toe, as the preservation of a portion of a toe is rather a discomfort than an advantage, except in the instance mentioned. All incisions should be made so that the resulting cicatrix does not occupy the plantar surface, and it is well to remember that the web of the toes is considerably below the position of the metatarso-phalangeal joint. (Fig. 378.)

The toes are usually removed by an incision on the dorsal

surface a little above the joint, which is carried down the bone for about an inch and then diverges into the web, and

FIG. 378.

Relations of web and metatarso-phalangeal joint of toes. (STIMSON.)

FIG. 379.



Incisions for amputation of toes and metatarsal bones. (STIMSON.)

is carried under the toe and back on the other side to the point of divergence. (Fig. 378.)

Amputation of Two Adjoining Toes.

The dorsal incision should be made in the inter-metatarsal space just above the level of the joint (Fig. 379, *B*) and carried down to the beginning of the web; then over the toe to the beginning of the adjoining web, and under the

plantar surface of both toes in the line of the digito-plantar fold, through the web and back to the point of divergence.

Amputation of the Great Toe.

This may be accomplished by means of the racket-shaped incision employed in amputation of the other toes or by means of a lateral flap. In the latter case the knife is made to enter the joint by cutting through the commissure, and the operation is completed by carrying the knife through the joint and along the outer or inner side of the bone, forming a flap of the required size. (Fig. 380.)

In this amputation a short dorsal flap and long plantar flap may be employed, or a long internal flap may be used.

Amputation of All the Toes.

To amputate all the toes, make a dorsal incision from the head of the fifth to the head of the first metatarsal

FIG. 380.



Amputation of the great toe. (SMITH.)

FIG. 381.



Incision for amputation of all the toes. (SMITH.)

bone; the incision should be a curved one passing just in front of the joints. (Fig. 381.) Dissect up the flap and open the joints, dividing the lateral ligaments, and pass the

knife behind the phalanges and cut a flap from the plantar surface.

Amputation of the Metatarsal Bones.

It is better in these amputations to leave the tarsal head of the metatarsal bone in place and divide the bone, or in other words to do an amputation in continuity to prevent opening up the tarsal articulations.

Amputation of the Metatarsal Bone of the Great Toe.

The incision begins upon the dorsal surface of the metatarsal bone, a little below the point at which the bone is to be divided, and is carried down below the metatarso-phalan-

Fig. 382.



Amputation of the great toe and first metatarsal bone. (Smith.)

geal joint, then diverges and passes under the toe and comes back again to the point of divergence. (Fig. 379, *C*.) The bone is exposed and cut through with cutting forceps and is then lifted up and dissected loose from the tissues. (Fig. 382.)

Amputation of the Fifth Metatarsal Bone.

The incision for the removal of the fifth metatarsal bone is made over the bone a little below the metatarso-tarsal articulation, and is carried down and curved around the toe (Fig. 379, *D*), and after the bone is exposed by dissecting back the flaps, it is divided, or the joint is opened and it is dissected out.

Amputation Through the Metatarsal Bones.

In performing this amputation an incision is made across the dorsum of the foot and a short dorsal flap is dissected up; the metatarsal bones are next divided with a saw and a long plantar flap is cut from within outward by entering the knife behind the ends of the bones.

Tarso-metatarsal Amputations.

In all amputations of the foot involving the tarsus the surgeon should be thoroughly familiar with the anatomy of the foot and the surgical landmarks of the different articulations. I shall refer to those laid down by Mr. Bryant, which are as follows:

"On the *inner* side of the foot not far from the inner malleolus the tubercle of the scaphoid (Fig. 383, *A*), is to be felt as a marked prominence; about one-half an inch in front of this will be found the articulation with the cuneiform bone (*B*), and one inch in front of this the joint which the surgeon will have to open in Lisfranc's or Hey's operation (*C*); just above the tubercle of the scaphoid will be found the articulation with the astragalus, the line of Chopart's amputation (*D*). On the *outer* side of the foot, one inch below the external malleolus, a sharply defined projection will always be felt. which is the peroneal tubercle (*E*), one-half an inch in front of this will be found the joint which separates the os calcis from the cuboid (*F*), this joint forming the outer circle to Chopart's amputation. Half an inch in front again or one inch from the tubercle, the

prominence of the fifth metatarsal bone is always to be felt (*H*), the line above this prominence indicating the articula-

FIG. 383.



Surgical guides to the foot as expressed by anatomy. (BRYANT.)

FIG. 384.



Incision for—*A*. Lisfranc's amputation. *B*. Chopart's amputation. (STIMSON.)

tion with the cuboid bone, which forms the outer boundary of the incision for Hey's or Lisfranc's amputations.

Tarso-metatarsal Amputation (*Lisfranc's*).

The incision for this amputation is a curved one carried across the dorsum of the foot from the base of the fifth to the base of the first metatarsal bone. (Fig. 384, *A*.) The incision should involve the skin only, its centre lying half

an inch or more below the centre of the line of the articulations, and it should begin and end at the sides of the foot at their junction with the sole. A plantar flap should be marked out by a curved incision crossing the sole of the foot near the origin of the toes, starting and ending at the same points as the dorsal incision.

The dorsal flap is next dissected back to the line of the articulations; the tendons, muscular fibres, and fascia being divided, the joints between the tarsal and metatarsal bones are opened with a stout, narrow-bladed knife. (Fig. 385.)

Fig. 385.

Amputation at tarso-metatarsal joint (Lisfranc's). (Skey.)

Difficulty is sometimes experienced in opening the joint between the head of second metatarsal bone and the second cuneiform bone, which occupies a position higher on the foot than the other articulations. The disarticulation may also be facilitated by forcibly depressing the anterior portion of the foot. After the joints have all been opened, the knife is passed behind the ends of the metatarsal bones and a plantar flap is cut from within outward, following the line of the incision previously marked out. The plantar flap may be cut from without inward if preferred.

Tarso-metatarsal Amputation (Hey's).

The line of incision and the steps of this operation are similar to those in Lisfranc's amputation, with the exception that Hey sawed off the projecting portion of the internal cuneiform bone after disarticulating the metatarsal bones. This modification, although it improves the appearance of the stump, possesses no advantages over the previous procedure.

Medio-tarsal, or Chopart's Amputation.

In this amputation the disarticulation is through the joints formed by the astragalus and calcaneum behind and the scaphoid and cuboid in front. An incision is made from

FIG. 386.



Line of incision for—*A*. Chopart's amputation. *B*. Syme's amputation. *C*. Section of bone in Syme's amputation. *D*. Subastragaloid amputation. (STIMSON.)

the tubercle of the scaphoid across the dorsum of the foot an inch in front of the head of the astragalus to the lower and outer border of the cuboid. (Fig. 386, *A*.) The plantar flap is next marked out by an incision beginning and ending at the same points as the first incision and crossing

the sole of the foot four or five finger-breadths nearer the toes. The dorsal flap is next dissected up, and after the tendons and fascia have been divided the joint is opened

Fig. 387.

Chopart's amputation. (Bryant.)

and a plantar flap is cut from within outward following the line of the previously marked out plantar incision. (Fig. 387.)

Subastragaloid Amputation.

In this amputation all the bones of the foot are removed except the astragalus. An incision is made beginning an inch below the tip of the external malleolus which is carried forward to the base of the fifth metatarsal bone; it is then carried over the dorsum of the foot to the calcaneo-cuboid articulation. (Fig. 386, *D.*) The joints between the scaphoid and astragalus and between the astragalus and calcis are opened, and the latter bone is carefully dissected out; the ligaments are divided and the astragalus only is allowed to remain in place.

Amputations at the Ankle-joint.

Syme's Amputation at the Ankle-joint.

In this amputation, the foot being at a right angle to the leg, an incision is made from the centre of one malleolus

directly across the sole of the foot to the centre of the opposite malleolus. (Fig. 386, *B*) The tissues of the heel are next carefully dissected from the bone by keeping the knife close to the osseous surface until the tuberosity of the os calcis is fairly turned. The two extremities of the first incision are then joined by a transverse one across the instep, and, the joint being opened, the lateral ligaments are divided to complete the disarticulation. (Fig. 388.) The

FIG. 388.



Syme's amputation at the ankle-joint. (SKEY.)

knife is next used to clear the malleoli, and they are next removed by the saw in the line indicated. (Fig. 386, *C*.)

Pirogoff's Amputation at the Ankle-joint.

In this amputation the posterior portion of the os calcis is retained. The incision is carried from the tip of the inner malleolus, over the instep, half an inch in front of the anterior edge of the tibia, to a point half an inch in

front of the tip of the outer malleolus; a second incision, crossing the sole of the foot and carried down to the bone,

Fig. 389.



Pirogoff's amputation. *A*, cutaneous incision. *B*, line of section of bones. (Stimson.)

is next made. (Fig. 389, *A*.) The plantar flap is dissected back for a quarter of an inch, the joint is opened by dividing

Fig. 390.



Application of saw to calcis in Pirogoff's amputation. (Erichsen.)

the lateral ligaments, and the astragalus is disarticulated, and the malleoli are exposed. A narrow saw is next applied to the upper and posterior part of the calcaneum behind the astragalus, and it is divided obliquely downward in the line of the plantar incision. (Fig. 390.) The malleoli and a thin slice of the tibia are next removed with the saw as in Syme's amputation. (Fig. 386, *C.*) Some surgeons do not remove the malleoli, but press the sawed surface of the os calcis between them when it is possible to do so. The position of the os calcis in relation to the tibia after union has occurred, is shown in Fig. 391.

Fig. 391.

Union between calcaneum and tibia in Pirogoff's amputation. (Hewson.)

Roux's Amputation at the Ankle-joint.

In this method of amputation an incision is made at the outer edge of the tendo Achillis, a little above its insertion, which is carried forward under the outer malleolus, and across the instep half an inch in front of the anterior edge of the tibia, and back to a point just in front of the inner malleolus; the incision is carried from this point downward and partly across the sole of the foot, and then back to the point of origin of the original incision. (Fig. 392.) The flaps are dissected up for a short distance, the ankle-joint is then opened, and the disarticulation is effected, and the internal flap is carefully dissected from the bones.

Other methods of amputation of the foot are sometimes emloyed; such, for instance, as that advocated by *Hancock*, who has combined Pirogoff's amputation with the subastragaloid method, bringing the sawed surface of the os calcis in contact with a transverse section of the astragalus.

Hancock has advocated the propriety of amputating in the foot without regard to the position of the tarsal joints, cutting

the flaps of sufficient length and dividing the bones with a saw.

Tripier has also modified the subastragaloid amputation by leaving the upper part of the calcaneum, which he saws through on a level with the sustenaculum tali, and at right angles to the axis of the leg; the external incisions are made as in Chopart's amputation.

FIG. 392.



Incisions in Roux's amputation.

In the method advocated by *Mikulicz* the astragalus and calcaneum are removed, the ends of the tibia and fibula are sawed off, and the sawed surface of the scaphoid and cuboid are approximated to these, the stump resulting resembling the foot of *pes equinus*.

AMPUTATIONS OF THE LEG.

The leg may be amputated at its lower, middle, or upper third, the rule being to save as much of the limb as possible, but as regards the application of prothetic apparatus, I think the stumps resulting from amputations in the middle and upper thirds will be found more satisfactory. It is well also in sawing the bones to divide the fibula at a slightly higher point than the tibia.

Amputation at the Lower Third of the Leg.

At this position the leg may be amputated by the circular, modified circular, or elliptical method.

Circular Method.

A circular incision is made through the skin and connective tissue just above the malleoli and the cuff is dissected up for a sufficient distance, and a circular incision of the tendons and muscles is next made and the tissues being retracted the bones are divided with a saw.

Modified Circular Method.

In this method of amputation of the leg a circular incision of the skin and connective tissue and two short lateral incisions are made and the flaps are dissected up to the end of the incisions, and a circular division of the muscles is next made. (Fig. 393, *A*.) Or oval skin flaps are made and dissected up, and the tissues are next divided down to the bone by a circular incision and the bones are divided with a saw. (Fig. 395.)

Elliptical Method.

In this method of amputation the incision is in the form of an ellipse, its lower end crosses the heel below the insertion of the tendo Achillis and the upper end of the incision is about an inch above the anterior articular edge of the tibia. (Fig. 394, *B*.)

Long Anterior Flap Method.

An anterior flap equal in length to the diameter of the leg at its base is marked out by a curved incision through the skin beginning at the posterior edge of the tibia on the inner side, a little below the point at which the bones are to be divided, and is carried over the leg to a point directly

opposite over the fibula. (Fig. 394, *A*.) The anterior muscles are divided transversely half an inch above the

Fig. 393.



Amputation of the leg. *A*. Modified circular method. *B*. Rectangular flap. *C*. Antero-posterior flaps. (Stimson.)

Fig. 394.



Amputation of the leg. *A*. Long anterior flap. *B*. Supra-malleolar long posterior flap. *C*. At upper third. (Stimson.)

lower end of the flap and are dissected from the bones to the base of the flap.

FIG. 395.

Oval skin flaps with circular division of the muscles.

The posterior flap is then made by entering the knife behind the bones at the point of the original incision and cutting directly outward.

Long Anterior Rectangular Flap Method. (*Teale.*)

In this method of amputation of the leg an incision equal in length to half of the circumference of the leg is made from the point at which the bones are to be divided on one side of the leg and is carried across the limb and back upon the opposite side to a point opposite the point of starting. The flap thus marked out is dissected up to its base and a posterior flap one-fourth the length is next cut by a transverse incision down to the bones and is dissected back to the line of the origin of the first incision. (Fig. 393, *B.*) The long flap is next doubled back and its edges secured to the posterior flap, or the long flap may be cut from the posterior surface of the leg and the short flap from the anterior surface.

Antero-posterior Flap Method.

A long anterior flap including half of the circumference of the limb may be cut from without inward, composed of skin, connective tissue, and muscles, and a short posterior flap cut from within outward may also be employed. This method is often employed in amputations in the upper portion of the leg. (Fig. 393, *C.*)

Lateral Flap Method.

In the lower and middle thirds of the leg the method of amputation by means of lateral skin flaps may be employed with advantage. In this method an incision is made over the spine of the tibia and an oval flap, embracing one-half of the circumference of the leg, composed of the skin and connective tissue, is dissected up; starting from the same point a similar flap is cut upon the opposite side of the leg and dissected up; the muscles at the upper extremity of the flaps are next divided by a circular incision and the bones are divided with a saw.

External Flap Method. (*Sédillot.*)

In this method of amputation of the leg the point of the knife is entered a finger's breadth external to the spine of the tibia and carried outward, grazing the fibula and is brought out as far as possible to the inner side; a flap three or four inches in length is then cut from within outward; the extremities of the incision are next united by an incision across the inner side of the limb involving the skin only; any remaining muscular tissue is next divided and the bones are sawed, and the long external flap is brought over the ends of the bones and fastened to the edges of the incision on the inner side of the limb. Prof. Ashhurst modifies this operation by cutting the long external flap from without inward, and makes also a short internal flap in the same manner. By either method the resulting stump is a good one, with the ends of the bone covered by the tissues of the external flap.

AMPUTATIONS AT THE KNEE-JOINT.

Amputations at the knee-joint may be done either by the circular or elliptical incision or by means of flaps, and may consist in simple disarticulations or sections through the condyles of the femur.

Elliptical or Oval Method.

In this operation an incision crossing the spine of the tibia five finger-breadths below the lower extremity of the patella is carried around the back of the leg three finger-breadths higher than in front; the tissues on the front of the leg are dissected up until the tendon of the patella is exposed; the leg is then flexed and the ligament of the patella is divided; the capsular ligament and the lateral and crucial ligaments are next divided, care being taken not to injure the popliteal vessels with the point of the knife. The tibia is next drawn forward and the knife is passed behind its posterior border, and the remaining soft parts are divided from within outward.

Anterior Flap Method.

In this method of amputation a long cutaneous flap is formed; the incision beginning half an inch below the articulation is carried five inches below the patella; crossing the anterior surface of the leg it is carried back to the condyle of the femur on the opposite side. This flap is dissected up and the ligament of the patella is divided, and the disarticulation is effected. A short posterior flap, uniting the anterior incision one inch below its extremities, is next cut by transfixion or from without inward. (Fig. 396, *A*.)

Amputation through the Condyles of the Femur.

In this amputation, which is known as *Carden's* amputation, an anterior flap, whose lower extremity is three finger-breadths below the patella, is cut and the disarticulation is effected, and the posterior soft parts are divided. The patella is removed and the condyles next sawed through just above the edge of the articular cartilage. (Fig. 396, *B*.)

Lateral Flap Method.

In this operation an incision is made just below the patella, which is carried down the spine of the tibia for

Fig. 396.



Amputations at the knee-joint and lower third of thigh. *A*. Long anterior flap. *B*. Amputation through condyles. *C*. Modified flap at lower third of thigh. (Stimson.)

Fig. 397.

Amputation at knee-joint by lateral flaps. (Smith.)

three inches, and is then carried backward to the middle of the leg to a point opposite the beginning of the incision; a similar flap is cut on the opposite side of the leg, and the flaps are dissected up to the line of the articulation, and when this point is reached the joint is opened and the disarticulation is effected. The patella is not removed. (Fig. 397.)

Gritti's Amputation at the Knee-joint.

In this operation a long rectangular anterior flap is first cut and dissected up, and after the disarticulation has been effected the skin covering the posterior surface of the knee is cut from within outward. The condyles of the femur are next removed by a saw above the edge of the articular cartilage, and the articular surface of the patella is removed by the saw or cutting forceps. The patella is next brought down so that its sawed surface

is in contact with the sawed surface of the condyles, and the flaps are brought together. (Fig. 398, *A.*)

Amputations of the Thigh.

Modified Flap Method.

Two semilunar flaps of skin and connective tissue, the upper extremity of which are several inches above the condyles of the femur, are cut and dissected up, and the muscles are next divided by a circular incision, and the bone is cut through with the saw. (Fig. 396, *C.*)

Long Anterior Flap Method.

In this operation an incision is made on the anterior aspect of the thigh, marking out a flap whose length is equal to one-third, and whose width at its base is equal to two-thirds, of the circumference of the limb. The anterior muscles are next divided obliquely upward and backward, so that the flap shall not be too thick, and the posterior muscles are cut transversely and the bone is divided with a saw. (Fig. 398, *B.*)

Amputation in the lower third of the thigh may also be effected by employing a *long anterior* and *short posterior* flap. The anterior flap is cut, its lower extremity extending down to the lower edge of the patella, and after dissecting up the skin and cellular tissues to the upper extremity of the patella, the muscles are cut obliquely up to the point at which the bone is to be divided. A short posterior flap is next cut, and the soft parts being retracted, the bone is sawed through. (Fig. 398, *C.*)

Amputation of the Thigh by Transfixion.

In amputations of the thigh the flaps may also be cut by transfixion, either *lateral* or *antero-posterior* flaps being employed. (Fig. 399.)

FIG. 398.



A. Gritti's amputation at the knee. *A'*. Lines of division of the bones. *B*. Amputation of the thigh, long anterior flap. *B'*. Division of the bone. *C*. Amputation at the lower third of the thigh. *C'*, Division of the bone *D*. Disarticulation at the hip-joint.

Amputation of the Thigh through the Trochanters.

When, for any reason, it is inadvisable to amputate at the hip-joint, an amputation may be made through the trochanters, a long anterior and short posterior flap being employed with a circular division of the muscles.

FIG. 399.

Amputation of thigh by flaps cut by transfixion.

AMPUTATIONS AT THE HIP-JOINT.

In amputations at the hip-joint it is important that provision be made for the control of hemorrhage during the

FIG. 400.



Abdominal tourniquet.

operation, and this is accomplished by the use of an abdominal tourniquet (Fig. 400), or by the use of Davy's lever making compression upon the common iliac artery from the rectum, or by compression of the femoral artery by the fingers of an assistant, or by the preliminary ligation of the femoral artery just below Poupart's ligament. Esmarch's

FIG. 401.



Esmarch's elastic strap applied to control hemorrhage during amputation at the hip-joint.

elastic strap may also be employed for the control of bleeding during amputation at the hip-joint, the strap being applied in such a manner that it occupies the position of the turns of a spica bandage of the groin. (Fig. 401.)

The most satisfactory method of controlling the bleeding during amputation at the hip-joint, or at the trochanters, is that recommended by Wyeth, which consists in the use of two stout steel pins twelve inches in length, and a piece of rubber tubing one-half of an inch in diameter and five or six yards in length. The point of one pin is inserted into the tissues one and a half inches below the anterior spine of the ilium, and is passed through the tissues external to the neck of the femur, and its point is made to project from the skin

just back of the great trochanter; the second pin is passed through the skin an inch below the level of the groin internal to the saphenous opening, and is carried through the adductor muscles and its point made to emerge half an inch in front of the tuberosity of the ischium; the rubber tubing is next wound around the thigh above the pins and securely tied.

The methods of amputation at the hip-joint are the *oval*, *antero-posterior flap*, and *lateral flap*, and *modified circular* methods.

Oval Method.

This is performed by entering the point of a strong knife into the tissues below the anterior superior spinous process of the ilium and making two oblique incisions, one forward and downward and the other backward, both incisions meeting on a transverse line on the inner side of the thigh. The muscles are next divided on a little higher line, and when the joint is exposed disarticulation is effected from the outside and any remaining tissue is divided.

Antero posterior Flap Method.

In this method the point of a long amputating knife is thrust into the tissues about two finger-breadths below the anterior superior spinous process of the ilium, and is pushed through the tissues grazing the hip-joint and is brought out on the opposite of the thigh close to the junction of the scrotum. The knife is next carried downward close to the bone and an anterior flap of sufficient length is cut from within outward. This flap is held up by an assistant and the head of the bone is disarticulated, and the knife being passed behind the bone, a posterior flap of equal length is cut from within outward. (Fig. 402.)

Guthrie's method of amputation at the hip-joint consists in cutting the flaps from without inward, a smaller knife being used for this purpose and the posterior flap being cut first.

FIG. 402.



Amputation at the hip-joint by antero-posterior flaps. (HOLMES.)

Modified Circular Method.

In this operation short antero-posterior flaps of skin and connective tissue are cut and dissected up, and the muscles

FIG. 403.

Amputation at the hip-joint by external and internal flaps. (BRYANT.)

are divided by a circular incision on the level of the joint, and the disarticulation of the head of the femur is next effected.

Lateral Flap Method.

In this operation two flaps are cut from the inner and outer side of the thigh by transfixion, or by cutting from without inward and exposing the joint, which is opened and the disarticulation of the head of the femur is effected as in the previous methods. (Fig. 403.)

INDEX.

ABDOMINAL aorta, ligation of, 413
bandage, 28
Abscess, acute, 263
chronic, 264
cold, 264
deep-seated, opening of, 263
dressing of, 263
opening of, 263
sinuses from, 265
Absorbent cotton, 131
A. C. E. anæsthetic mixture, 202
Acromion process of scapula, fracture of, 320
Actual cautery, 155
Acupressure, 254
first method of, 254
second method of, 254
third method of, 255
fourth method of, 255
fifth method of, 255
sixth method of, 256
seventh method of, 256
Acupuncture, 152
needles, 152
Acute abscess, 263
Adhesive plaster, 134
Agnew's splint for fracture of patella, 346
American bandage of foot, 72
Amputating knives, 433, 434
saws, 434
Amputation or amputations, 428
at ankle-joint, 464
Pirogoff's, 465
Roux's, 467
Syme's, 464
of arm, 450
circular, 429
details of, 438
at elbow, 448
elliptical, 431
of fingers, 441
metacarpo-phalangeal, 443
Amputations, flap, 429
of foot, 460
Hancock's, 467
Hey's, 463
Lisfranc's, 461
Mikulicz's, 468
Tripier's, 467
of forearm, 447
of hand, 441
carpo-metacarpal, 446
at hip-joint, 477 *et seq.*
Guthrie's, 479
instruments for, 433
at knee-joint, 472
Carden's, 473
Gritti's, 474
of leg, 468 *et seq.*
Sédillot's, 472
medio-tarsal, 463
of metacarpal bones, 445
of metatarsal bones, 459
methods of, 428
modified circular, 431
oval, 431
periosteal flaps in, 433
by rectangular flaps, 432
re-dressing of, 440
above shoulder-joint, 456
at shoulder-joint, 452 *et seq.*
Dupuytren's, 454
Larrey's, 453
Lisfranc's, 455
Spence's, 455
subastragaloid, 464
sutures in, 437
tarso-metatarsal, 460
Hey's, 463
Lisfranc's, 461
Teale's method, 432
of thigh, 475 *et seq.*
of toes, 456
at wrist, 446
Anæsthesia from cocaine, 193
from cold, 192

Anæsthesia, local, 192
from nitrous oxide gas, 194
from rapid respiration, 193
from rhigolene, 192
Anæsthetic mixture, A. C. E., 202
Anæsthetics, 192
in tracheotomy, 279
Aneurism needle, 394
Ankle-joint, amputations at, 464
Pirogoff's, 465
Roux's, 467
Syme's, 464
dislocations of, 387
Anterior tibial artery, ligation of, 422
Antisepsis, 99
Antiseptic bandages, 118
dressings, improvised, 118
gauze, 115
poultice, 142
Aorta, abdominal ligation of, 413
compressor, Lister's, 245
Aqua ammonia, 149, 151
Aristol, 108
Arm, amputations of, 450
Arterial hemorrhage, 241
control of, permanent, 249
temporary, 241
transfusion, 167
Arteriotomy, 164
Artery or arteries—
anterior tibial, ligation of, 422
axillary, ligation of, 405
brachial, ligation of, 407
common carotid, ligation of, 399
iliac, ligation of, 413
dorsalis pedis, ligation of, 424
external carotid, ligation of, 401
iliac, ligation of, 416
facial, ligation of, 404
femoral, ligation of, 418 *et seq.*
forceps for, 436
gluteal, ligation of, 417
inferior thyroid, ligation of, 399
innominate, ligation of, 395
internal carotid, ligation of, 402
iliac, ligation of, 415
pudic, ligation of, 417
interosseous, ligation of, 413
ligation of, 393 *et seq.*
lingual, ligation of, 403
occipital, ligation of, 404
popliteal, ligation of, 421
posterior tibial, ligation of, 424
radial, ligation of, 409
sciatic, ligation of, 417
Artery, subclavian, ligation of, 396
superior thyroid, ligation of, 403
temporal, ligation of, 404
ulnar, ligation of, 411
vertebral, ligation of, 398
wounded, ligation of, 258
Artificial respiration, 170
direct method of, 171
Howard's method of, 171
Marshall Hall's method of, 175
Sylvester's method of, 174
Asepsis, 99
Aseptic operation, details of, 122
preparation for, 120
surgery, materials used in, preparation of, 109
Aspiration, 177
Aspirator, 177
Astragalus, dislocation of, 388
fracture of, 355
Auto-transfusion, 168
Axillary artery, ligation of, 405

BACILLUS pyocyaneus, 100
pyogenes fœtidus, 100
Bandage or bandages—
abdominal, 28
antiseptic, 118
Barton's, 32
modified, 33
of chest, anterior figure-of-eight, 60
posterior figure-of-eight, 60
circular, 19, 22
compound, 23
crossed, of eye, 41
of both eyes, 42
demi-gauntlet, 48
Desault's, 56
dimensions of, 17
Esmarch's, 247
figure-of-eight, 22
of elbow, 51
of knee, 68
of both knees, 69
of leg, 75
of neck and axilla, 54
of foot, American, 72
covering heel, 72
not covering heel, 73
French, 73
spica, 71
four-tailed, of chin, 27
of head, 27

Bandage, gauntlet, 46
Gibson's, 34
glue and oxide of zinc, 96
gum and chalk, 95
handkerchief, 28
cord, 29
cravat, 29
oblong, 28
square, 28
triangle, 29
for hands and feet, 78
of head, 32
and neck, 40
oblique, 44
Liebreich's, 78
of lower extremity, 65
many-tailed, 26 *et seq.*
of neck, 32
oblique, 19
of angle of jaw, 36
occipito-facial, 43
paraffin, 96
plaster-of-Paris, 83
application of, 84
preparation of, 84
removal of, 93
trapping of, 92
Pott's, 80
recurrent, 22
of head, 37
of stump, 77
roller, 14
rubber, 81
scissors, 18
of Scultetus, 79
silicate of potassium, 95
of sodium, 95
special, 76
spica, 21
of foot, 71
of groin, ascending, 65
descending, 66
of shoulder, ascending, 52
descending, 53
of thumb, 49
spiral, 19, 20
of chest, 59
of finger, 45
reversed, 20
of lower extremity, 74
of penis, 76
of upper extremity, 50
starched, 94
suspensory and compressor, of breast, 62
of both breasts, 63
Bandage of trunk, 59
of upper extremity, 45
Velpeau's, 55
winder, 14
Bandaging, 13
rules for, general, 17 *et seq.*
Barton's bandage, 32
modified, 33
handkerchief, 31
Bavarian dressing, 90
Bedsores, 272
Bellocq's canula, 261
Beta-naphthol, 105
Bichloride cotton, 119
of mercury, 103
gauze, 115
Binder's-board splints, 98, 300
in compound fractures, 360
Bis-axillary cravat, 36
Bladder, hemorrhage from, 262
washing out of, 215
Blood, transfusion of, 164
direct, 165
indirect, 166
Bloodletting, 157
Bond's splint, 332
Bone forceps, 435
Boric acid, 107
Boro-salicylic lotion, 107
Bougies, 209
bulbous, 209
filiform, 209
œsophageal, 180
rectal, 191
Bouisson's suture, 230
Brachial artery, ligation of, 407
Bran bags, 301
dressing in compound fractures, 360
Bread poultice, 141
Breast, strapping of, 137
suspensory and compressor bandage of, 62
double, 63
Bruises, 270
Bulbous bougies, 209
Buried suture, 223
Burns, 271
Button suture, 226

CALCANEUM, fracture of, 354
Cantharidal collodion, 150
Cantharis, 150
Capillary hemorrhage, 241

Capillary hemorrhage, treatment of, 257
Capsicum, 149
Carbolic acid, 104
Carbolized gauze, 117
Carbuncle, strapping of, 140
Carden's amputation at knee-joint, 473
Carpal bones, fracture of, 334
Carpo-metacarpal amputation of hand, 446
Carpus, dislocation of, 378
Carotid artery, common, ligation of, 399
external, ligation of, 401
internal, ligation of, 402
Cartilages, costal, fractures of, 310
semi-lunar, dislocations of, 386
Catgut, chromic acid, 111
for drainage, 113
juniper, 111
ligatures, 111
sutures, 111
Catheter, elbowed, 208
female, 212
introduction of, 212
flexible, 207
introduction of, 209
metallic, 206
prostatic, 207
soft rubber, 208
tying in, 213
Cauterization in arterial hemorrhage, 251
Cautery, actual, 155
Charcoal poultice, 142
Chest, figure-of-eight bandage of, anterior, 60
posterior, 60
spiral bandage of, 59
strapping of, 137
T-bandage of, double, 26
single, 24, 25
Children, fractures of the femur in, 343
leg in, 352
Chin, four-tailed bandage of, 27
Chloride of zinc, 106
Chloroform, 148, 151, 200
administration of, 200
apparatus, Clover's, 201
Chopart's amputation of foot, 463
Chromic acid catgut, 111
Chronic abscess, 264
Circular amputation, 429
bandage, 19, 22
Clavicle, dislocations of, 368
of acromial end of, 369
of sternal end of, 368
fracture of, 314 *et seq.*
in children, 318
modified, Velpeau's dressing for, 319
Sayre's dressing for, 316
Velpeau's dressing for, 317
Cleanliness, surgical, 102, 120
Clinical thermometer, 189
Closed fracture, 291
Clove-hitch, 364
Clover's chloroform apparatus, 201
Cocaine, anæsthesia from, 193
Coccyx, dislocations of, 365
fracture of, 311
Cold abscess, 264
anæsthesia from, 192
in arterial hemorrhage, 249
compresses, 146
Cold-water dressings, 146
Colles's fracture, 332
Collodion, cantharidal, 150
Condyles of femur, fractures of, 344
Congenital dislocations, 392
Consecutive hemorrhage, 241
Constitutional treatment of hemorrhage, 241
Continued suture, 223
Contused wounds, 268
Contusions, 270
Comminuted fracture, 291
Common carotid artery, ligation of, 399
iliac artery, ligation of, 413
Complete dislocation, 362
fracture, 290
Complicated dislocation, 362, 392
fracture, 292
Compound bandages, 23
dislocation, 362, 392
fracture, 291
Binder's board splints in, 360
bran dressing in, 360
dressing of, 356
felt splints in, 360
plaster-of-Paris dressing in, 359
sawdust dressing in 361
Compresses, 132, 302
cold, 146
in hemorrhage, 242
hot, 143
Coracoid process of scapula, fracture of, 320

Coronoid process, fracture of, 328
Corrosive sublimate gauze, 115
Costal cartilages, fractures of, 310
Cotton, 130
absorbent, 131
bichloride, 119
Counter-irritation, 147
Creolin, 107
Crossed bandage of eye, 41
of both eyes, 42
Cruro-pelvic triangle, 30
Cupping, 158
dry, 159
-glass, 159
wet, 160
Cutting pliers, 435
Czerny suture, 231

DEEP incisions, 158
Demi-gauntlet bandage, 48
Desault's bandage, 56
first roller, 56
second roller, 57
third roller, 58
Diastasis of sternum, 367
Diffused suppuration, 265
Digital compression in hemorrhage, 241
Dilators, tracheal, 277
Director, tracheotomy, 276
Dislocation or dislocations, 362
of acromion process of scapula, 369
of ankle, 387
of astragalus, 388
of carpal bones, 378
of carpus, 378
of clavicle, 368
of coccyx, 365
complete, 362
complicated, 362, 392
compound, 362, 392
congenital, 392
dressing of, 362
of elbow, 374
of femur, 382
anomalous, 384
downward and forward, 382
forward and upward, 384
pubic, 384
thyroid, 383
of fibula, 387
of fingers, 379
of head of radius, 377
of hip, 382
of hyoid bone, 366
Dislocation of the inferior angle of scapula, 369
of jaw, 365
of knee, 386
of metacarpal bones, 379
of metatarsal bones, 390
old, 363, 390
vertical extension in, 391
partial, 362
of patella, 385
pathological, 392
of pelvis, 367
of phalanges of toes, 390
of proximal phalanx of thumb, 380
recent, 362
of ribs, 366
of scapula, 369
of semilunar cartilages, 386
of shoulder, 370
reduction of, 371
simple, 362
spontaneous, 392
of sternum, 367
of tarsal bones, 388, 389
of toes, 390
treatment of, 363
of upper end of ulna, 377
of vertebræ, 364
of wrist, 377
Dorsal dislocation of femur, 382
Dorsalis pedis artery, ligation of, 424
Double cyanide of mercury and zinc 109
gauze, 116
ligature, 236
roller bandage, 16
spring artery forceps, 252
Drainage, catgut for, 113
horsehair for, 113
-tubes, 112
glass, 113
rubber, 112
Dressing or dressings—
antiseptic, 101
improvised, 118
moist method in, 125
reapplication of, 125
Bavarian, 90
cold water, 146
of compound fractures, 356
of dislocations. See under each dislocation.
dry sterilized, 120
of fractures. See under each fracture.

Dressing, fixed, 82
gauze, 115
preparation of, 115
hardening, 82
moss, 117
plaster-of-Paris, interrupted, 86
sawdust, 117
of septic wounds, 128
of wounds, 266
Dry cupping, 159
dressings in wounds, 101
sterilized dressings, 120
Dupuytren's amputation at shoulder-joint, 454
splint, 353

ELASTIC ligatures, 240
Elbow, amputations of, 448
dislocations of, 374
figure-of-eight bandage of, 51
Elbowed catheter, 208
Electrolysis, 186
Elliptical amputation, 431
Endoscope, 214
Enema, glycerin, 191
Enemata, 191
nutritious, 191
Epiphyseal fracture, 294
separation, 294
Epistaxis, 259
Erichsen's ligature, 239
Esmarch's bandage, 247
elastic strap, 246
Ether, 195
administration of, 195
first insensibility from, 197
inhaler, 196
Exploring needle, 184
trocar, 184
Extension, vertical, in old dislocations, 391
External carotid artery, ligation of, 401
iliac artery, ligation of, 416
Eye, Liebreich's bandage for, 78
crossed bandage of, 41
Eyes, crossed bandage of both, 42

FACIAL artery, ligation of, 404
Fascia, strains of, 275
Faradization, 187
Felt splints, 98, 300
in compound fractures, 360
Female catheter, 212
Femoral artery, ligation of, 418 *et seq.*
hernia, truss for, 205
Femur, dislocations of, 382
Femur, dislocations, anomalous, 384
backward, 382
reduction of, 382
downward and forward, 382
reduction of, 383
dorsal, 382
reduction of, 382
below tendon, 382
forward and upward, 384
reduction of, 384
iliac, 382
ischiatic, 382
posterior, 382
pubic, 384
reduction of, 384
thyroid, 383
reduction of, 383
fracture of, 337
in children, 343
condyles of, 344
dressing of, 337
green-stick, 344
lower end of, 344
shaft of, 340
upper extremity of, 337
Fermenting poultice, 142
Fibula, dislocations of, 387
fracture of, 352
of lower end of, 353
Figure-of-eight bandage, 22
of chest, anterior, 60
posterior, 61
of elbow, 51
of knee, 68
of both knees, 69
of leg, 75
of neck and axilla, 54
Filiform bougie, 209
Fingers, amputation of, 441 *et seq.*
dislocations of, 379
spiral bandage of, 45
Fissured fracture, 291
Fixed dressings, 82
Flap amputation, 429
Flaps, periosteal in amputation, 433
Flaxseed poultice, 141
Flexible catheters, 207
Fomentations, hot, 142
Foot, American bandage of, 72
amputations of, 460
Chopart's, 463
Hancock's, 467
Hey's, 463
Lisfranc's, 461
Mikulicz', 468
Tripier's, 467

Foot, bandage of, not covering heel, 73
covering heel, 72
-bath, mustard, 149
fractures of bones of, 354
spica bandage of, 71
Forced respiration, 176
Forceps, artery, 436
double spring, 252
bone, 435
hæmostatic, 247, 276, 436
torsion, 251
tracheal, 278
Forearm, amputation of, 447
fracture of bones of, 329
green-stick fracture of, 331
Foreign bodies, tracheotomy for, 284
Fracture or fractures, 290
of acromion process of scapula, 320
anæsthetics in, 296
of astragalus, 355
-bed, 298
-box, 301
double inclined, 340
of body of scapula, 320
of bones of foot, 354
of leg in children, 352
of calcaneum, 354
of carpal bones, 334
of clavicle, 314 *et seq.*
closed, 291
of the coccyx, 311
comminuted, 291
complete, 290
complicated, 292
compound, 291
binder's-board splints in, 360
bran dressing in, 360
continuous irrigation in, 361
dressing of, 356
felt splints in, 360
plaster-of-Paris dressing in, 359
sawdust dressing in, 361
of coracoid process of scapula, 320
of coronoid process of ulna, 328
of costal cartilages, 310
direction of, 293
dressing of, 290 *et seq.*
provisional, 296
epiphyseal, 294
examination of, 295
of femur, 337
Fracture of femur in children, 343
of fibula, 352
fissured, 291
of forearm, 329
green-stick, 290
of humerus, 321
of lower extremity, 324
of shaft, 322
of upper extremity, 321
of hyoid bone, 308
impacted, 292
incomplete, 290
indented, 290
of jaw, 304, 305
of larynx, 308
of leg, 348
longitudinal, 293
of malar bone, 304
of maxilla, lower, 305
upper, 304
of metacarpal bones, 335
of metatarsal bones, 356
multiple, 292
of nasal bones, 302
of neck of radius, 329
of scapula, 320
oblique, 293
of olecranon, 327
open, 291
dressing of, 356
partial, 290
of patella, 345
of pelvis, 311
of phalanges of fingers, 335
of toes, 356
Pott's, 353
punctured, 290
of radius, 329, 331
of head, 329
of lower end, 331
reduction of, 298
repair of, 294
of ribs, 309
of sacrum, 311
of scapula, 320
setting of, 298
of shaft of femur, 340
simple, 291
of skull, 313
of sternum, 310
of trachea, 308
transverse, 293
of trunk, 309
of ulna, 327
of upper extremity, 314
varieties of, 290

Fracture of vertebræ, 312
of zygoma, 304
French bandage of foot, 73

GALVANO-CAUTERY, 186
Gastrostomy, sutures for, 233
Gastrotomy, sutures for, 235
Gauntlet bandage, 46
Gauze, bichloride of mercury, 115
carbolized, 117
corrosive sublimate, 115
double cyanide of mercury and zinc, 116
dressings, 115
preparation of, 115
iodoform, 116
Gély's suture, 229
Gibson's bandage, 34
Glass drainage-tube, 113
Glover's suture, 223
Glue and oxide of zinc bandage, 96
Gluteal artery, ligation of, 417
Golding-Bird's tracheal dilator, 277
Granny knot, 221
Green-stick fracture, 290
of bones of forearm, 331
Gritti's amputation at knee-joint, 474
Groin, spica bandage of, ascending, 65
descending, 66
double, 67
T-bandage of, 24, 25
Gum and chalk bandage, 95
Gunshot wounds, 269
Guthrie's amputation at hip-joint, 479
Gutta-pecha splints, 300

HÆMOSTATIC forceps, 247, 276, 436
Hancock's amputation of foot, 467
Hand, amputation of, 441
carpo-metacarpal, 446
Hands, cleansing of, 120
removal of plaster-of-Paris from, 93
Handkerchief bandages, 28
Barton's, 31
Hardening dressings, 82
Hare-lip suture, 224
Hatter's felt splints, 98
Head, bandages of, 32
four-tailed, 27
oblique, 44
recurrent, 37
Head and neck bandage, 40
V-bandage of, 37
Hemorrhage, arterial, 241
cauterization in, 251
cold in, 249
control of, permanent, 249
temporary, 241
hot water in, 249
ligation in, 252
position in, 249
pressure in, 250
torsion in, 251
styptics in, 250
from bladder, 262
capillary, 241
treatment of, 257
compresses in, 242
consecutive, 241
deep suture in, 253
digital compression in, 241
Esmarch's elastic strap in, 246
intermediary, 241
parenchymatous, treatment of, 257
primary, 241
from rectum, 262
secondary, 241
treatment of, 257
Spanish windlass in, 244
treatment of, 241
constitutional, 241
local, 241
from urethra, 261
venous, 241
treatment of, 256
Hernia, femoral, truss for 205
inguinal, truss for, 204
irreducible, truss for, 206
umbilical, truss for, 205
Hey's amputation of foot, 463
Hip-joint, amputations at, 477 *et seq.*
Guthrie's, 479
Hip, dislocations of, 382. See Femur.
Hoey's clamp, 245
Hood's truss, 204
Horsehair for drainage, 113
Hot compresses, 143
fomentations, 142
water, 147
in arterial hemorrhage, 249
Howard's method of artificial respiration, 171
Humerus, dislocation of, subclavicular, 370
subcoracoid, 370
subglenoid, 370

Humerus, dislocation of, subspinous, 371
fracture of, 321
of lower extremity of, 324
of shaft of, 322
of upper extremity of, 321
Hydrogen peroxide, 106
Hypodermic syringe, 183
injections, 182
Hyoid bone, dislocation of, 366
fracture of, 308

ICE-BAG, 147
Iliac artery, common, ligation of, 413
external, ligation of, 416
internal, ligation of, 415
dislocation of femur, 382
Impacted fracture, 292
Incised wounds, 266
Incisions, deep, 158
Incomplete fracture, 290
India-rubber suture, 224
Indented fracture, 290
Inferior thyroid artery, ligation of, 399
Inguinal hernia, truss for, 204
Injections, hypodermic, 182
intra-venous, of milk, 169
of saline solution, 168
rectal, 191
urethral, 216
Innominate artery, ligation of, 395
Instruments for amputation, 433
sterilizing of, 121
Internal carotid artery, ligation of, 402
iliac artery, ligation of, 415
pudic artery, ligation of, 417
Intermediary hemorrhage, 241
Interosseous artery, ligation, of, 413
Interrupted suture, 222
Intestinal anastomosis, sutures for, 232
Intra-venous injection of milk, 169
of saline solution, 168
Intubation of larynx, 286
-tube, 286
extractor, 287
Iodoform, 105
gauze, 116
Irreducible hernia, truss for, 206
Irrigating apparatus, 123
Irrigation, 143
continuous, 144
in compound fractures, 361
Irrigation, mediate, 145
Ischiatic dislocation of femur, 382
Isinglass plaster, 135
Issue pea, 153
Issues, 152

JACKET, plaster-of Paris, application of, 86
Jaw, bandage of angle of, oblique, 36
dislocations of, 365
lower, fracture of, 305
splint for, 307
upper, fracture of, 304
Jobert's suture, 231
Joints, strapping of, 139
Jugular vein, external, bleeding from, 163
Juniper catgut, 111
Junk bags, 301
Jury-mast, application of, 89
Jute, 131

KNEE, figure-of-eight bandage of, 68
of both, 69
-joint, amputations at, 472
Carden's, 473
Gritti's, 474
dislocations of, 386
Knives, amputating, 433, 434
Knot, granny, 221
reef or flat, 219
Staffordshire, 221
surgeon's, 220
Kreolin, 107

LACERATED wounds, 267
Larrey's amputation at the shoulder-joint, 453
Laryngotomy, 284
Laryngo-tracheotomy, 285
Larynx, fractures of, 308
intubation of, 286
Leather splints, 97, 300
Leech, mechanical, 162
Leeching, 160
Lembert's suture, 228
Leg, amputation of, 468 *et seq.*
Sédillot's, 472
figure-of-eight bandage of, 75
fracture of, 348
in children, 352
Liebreich's bandage, 78
Ligation of abdominal aorta, 413
of anterior tibial artery, 422
in arterial hemorrhage, 252

Ligation of arteries, 393 *et seq.*
of axillary artery, 405
of brachial artery, 407
of common carotid artery, 399
iliac artery, 413
of dorsalis pedis artery, 424
of external carotid artery, 401
iliac artery, 416
of facial artery, 404
of femoral artery, 418
of gluteal artery, 417
of inferior thyroid artery, 399
of innominate artery, 395
of interosseous artery, 413
of internal carotid artery, 402
iliac artery, 415
pudic artery, 417
of lingual artery, 403
of occipital artery, 404
of popliteal artery, 421
of posterior tibial artery, 424
of radial artery, 409
of sciatic artery, 417
of subclavian artery, 396
of superior thyroid artery, 403
of temporal artery, 404
of ulnar artery, 411
of vertebral artery, 398
of wounded arteries, 258
Ligature or ligatures, 437
catgut, 111
double, 236
elastic, 240
Erichsen's, 239
quadruple, 237
securing of, 219
single, 235
subcutaneous, 237
for vascular growths, 235
Lingual artery, ligation of, 403
Lint, 129
Lisfranc's amputation of foot, 461
at shoulder-joint, 455
Lister's aorta compressor, 245
Local anæsthesia, 192
Longitudinal fracture, 293
Lower extremity, bandages of, 65
spiral reversed, 74
Luxations. See Dislocations.

MACKINTOSH, 114
Malar bone, fracture of, 304
Marshall Hall's method of artificial respiration, 175
Massage, 188
Maxilla, lower, fracture of, 305
upper, fracture of, 304
Mechanical leech, 162
Medio-tarsal amputation, Chopart's, 463
Mercier's catheter, 208
Mercury bichloride, 103
Metacarpal bones, amputation of, 445
dislocations of, 379
fracture of, 335
Metallic catheters, 206
Metatarsal bones, amputation of, 459
dislocation of, 390
fractures of, 356
Mikulicz, amputation of foot, 468
Milk, intra-venous injection of, 169
Minor surgery, 99
Moist dressings in wounds, 102
Moss dressing, 117
Motion, passive, 188
Moulded plaster-of-Paris splints, 91
Mouth-gag, 286
-to-mouth inflation, 170
Moxa, 153
Multiple fracture, 292
Muscles, strains of, 275
Muslin, oiled, 131
Mustard, 148
foot-bath, 149
papers, 149
plaster, 148

NASAL bones, fracture of, 302
Neck, bandages of, 32
Needle or needles, acupuncture, 152
aneurism, 394
exploring, 184
-holder, 219
mounted, 218
seton, 154
surgical, 218
Nitrate of silver, 151
Nitrous oxide gas, 194
Nose, T-bandage of, double, 26
Nutritious enemata, 191

OAKUM, 130
poultice, 142
Oblique bandage, 19
Oblique bandage of angle of jaw, 36
of head, 44
fracture, 293
Occipital artery, ligation of, 404
Occipito-facial bandage, 43
Œsophageal bougie, 180
Oiled muslin, 131
silk, 131
Old dislocation, 363, 390
Olecranon, fractures of, 327

Open fracture, 291
dressing of, 356
Operation, aseptic, details of, 122
preparation for, 120
of patient for, 121
Oval amputation, 431

PAPER, paraffin, 132
parchment, 114, 132
splints, 300
waxed, 132
Paquelin's thermo-cautery, 156
Paraffin bandage, 96
paper, 132
Parchment paper, 114, 132
Parenchymatous hemorrhage, treatment of, 257
Partial dislocation, 362
fracture, 290
Passive motion, 188
Pasteboard splints, 98
Patella, dislocations of, 385
fracture of, 345
Pathological dislocations, 392
Pelvis, dislocation of, 367
fractures of, 311
Penis, spiral reversed bandage of, 76
Periosteal flaps in amputation, 433
Periosteotome, 435
Permanganate of potassium, 108
Peroxide of hydrogen, 106
Petit's tourniquet, 243
Phalanges of fingers, dislocation of, 379
fractures of, 335
of toes, dislocations of, 390
fractures of, 356
Pirogoff's amputation at ankle-joint, 465
Plaster or plasters, 134
adhesive, 134
isinglass, 135
mustard, 148
resin, 134
rubber adhesive, 134
soap, 135
spice, 149
Plaster-of-Paris bandage, 83
application of, 84
preparation of, 84
removal of, 93
saw for, 93
shears for, 94
trapping of, 92
dressing in compound fractures, 359
Plaster-of-Paris bandage in fracture of femur, 339
interrupted, 86
jacket, application of, 86
suspensory apparatus for, 87
removal of, from hands, 93
splints, 300
moulded, 91
Plate suture, 226
Poisoned wounds, 269
Popliteal artery, ligation of, 421
Porous felt splints, 98
Porte-moxa, 153
Position in arterial hemorrhage, 249
Posterior tibial artery, ligation of, 424
Potassium permanganate, 108
Pott's bandage, 80
fracture, 353
Poultice or poultices, 140
antiseptic, 142
bread, 141
charcoal, 142
fermenting, 142
flaxseed, 141
oakum, 142
Powder-burns, 270
Pressure in arterial hemorrhage, 250
Primary hemorrhage, 241
roller, 301
Prostatic catheters, 207
Protective, 113
Provisional dressings of fracture, 296
Pubic dislocation of femur, 384
Pudic artery, internal, ligation of, 417
Puncturation, 158
Punctured fracture, 290
wounds, 268
Pyoktanin, 108

QUADRUPLE ligature, 237
Quilled suture, 225
Quilt suture, 224

RACK for fractures, 302
Radial artery, ligation of, 409
Radius, dislocation of head of, 377
fracture of, 329
head of, 329
lower end of, 331
neck of, 329
Rapid respiration, anæsthesia from, 193

Raw-hide splints, 97
Recent dislocations, 362
Rectal bougies, 191
injections, 191
tube, 190
Rectum, hemorrhage from, 262
Recurrent bandage, 22
of head, 37
of stump, 77
Reduction of dislocations, 363
of fractures, 298
Reef knot, 219
Resin plaster, 134
Respiration, artificial, 170
direct method of, 171
Howard's method, 171
Marshall Hall's method of, 175
Sylvester's method of, 174
forced, 176
Retractors, 133, 437
three-tailed, 133
two-tailed, 133
Rhigolene, anæsthesia from, 192
Ribs, dislocations of, 366
fractures of, 309
Roller bandage, 14
double, 16
preparation of, 15
single, 16
primary, 301
Roux's amputation at ankle-joint, 467
Rubber adhesive plaster, 134
bandage, 81
drainage-tube, 112
tissue, 114, 132
Rubefacients, 147

SACRUM, fractures of, 311
Saline solution, intra-venous injection of, 168
Sand bags, 301
Saphena vein, internal, bleeding from, 164
Saw, amputating, 434
for plaster-of-Paris bandage, 93
Sawdust dressing, 117
in compound fractures, 361
Sayre's dressing for fracture of clavicle, 316
Scalds, 271
Scapula, dislocations of, 369
acromial process of, 369
inferior angle of, 369
fracture of, 320
Scapula, fracture of body of, 320
neck of, 320
Scarification, 157
Scarificator, 160
Sciatic artery, ligation of, 417
Scissors, bandage, 18
skin-grafting, 185
Scultetus, bandage of, 79
Secondary hemorrhage, 241
treatment of, 257
sutures, 217
Sédillot's amputation of leg, 472
Semilunar cartilages, dislocation of, 386
Septic wounds, dressing of, 128
Seton, 153
needle, 154
Setting of fractures, 298
Shears for plaster-of-Paris bandage, 94
Shotted suture, 227
Shoulder, dislocations of, 370
reduction of, 371
-joint, amputation above, 456
amputation at, 452 *et seq.*
Dupuytren's, 454
Larrey's, 453
Lisfranc's, 455
Spence's, 455
spica bandage of ascending, 52
descending, 53
Signorini's tourniquet, 245
Silicate of potassium bandage, 95
of sodium bandage, 95
Silk, 110
oiled, 131
Silkworm-gut, 111
Simple dislocation, 362
fracture, 291
Sinapism, 148
Single ligature, 235
roller bandage, 16
Sinuses from abscesses, 265
Skin-grafting, 185
Skull, fractures of, 313
Slings, 26
Soap plaster, 135
Soft rubber catheter, 208
Sounds, 209
Spanish windlass, 244
Spence's amputation at the shoulder-joint, 455
Spica bandage, 21
of foot, 71
of groin, ascending, 65
descending, 66

Spica bandage of groin, double, 67
of shoulder, ascending, 52
descending, 53
of thumb, 49
Spice plaster, 149
Spiral bandage, 19, 20
of chest, 59
of finger, 45
reversed bandage, 20
of lower extremity, 74
of penis, 76
of upper extremity, 50
Splints, 299
angular, anterior, 325
internal, 323
binder's board, 98, 300
in compound fractures, 360
Bond's, 332
Dupuytren's, 353
felt, 300
in compound fractures, 360
gutta-percha, 300
hatter's felt, 98
leather, 97, 300
moulded binder's-board, 351
paper, 300
pasteboard, 98
plaster of-Paris, 300
moulded, 91
porous felt, 98
raw-hide, 97
wooden, 299
Sponges, 109
Spontaneous dislocations, 392
Sprain-fracture, 274
Sprains, 273
Staffordshire knot, 221
Staphylococcus pyogenes albus, 100
aureus, 100
Starched bandage, 94
Sterilization, chemical, in dressing of wounds, 101
Sterilized dressings, dry, 120
Sterilizing of instruments, 121
oven, 119
Sternum, diastasis of, 367
dislocations of, 367
fractures of, 310
Stomach-pump, 180
-tube, 178
Strains of fascia, 275
of muscles, 275
Strangury, 151
Strapping, 135
of breast, 137
Strapping of carbuncle, 140
of chest, 137
of joints, 139
of testicle, 136
of ulcers, 138
Streptococcus pyogenes, 100
Stump, recurrent bandage of, 77
Styptics in arterial hemorrhage, 250
Subastragaloid amputation, 464
Subclavian artery, ligation of, 396
Subclavicular dislocation of humerus, 370
Subcoracoid dislocation of humerus, 370
Subcutaneous ligature, 237
Subglenoid dislocation of humerus, 370
Subspinous dislocation of humerus, 371
Sulphocarbolate of zinc, 106
Superior thyroid artery, ligation of, 403
Suppuration, diffused, 265
Surface thermometer, 189
Surgeon's knot, 220
and reef knot combined, 220
Surgery, aseptic, materials used in, preparation of, 199
minor, 99
Surgical cleanliness, 102, 120
needles, 218
Suspensory apparatus for plaster-of-Paris jacket, 87
and compressor bandage of breast, 62
of both breasts, 63
Sutures or suture, 217, 437
of approximation, 217
Bouisson's, 230
buried, 223
button, 226
catgut, 111
of coaptation, 217
continued, 223
Czerny, 231
deep, in hemorrhage, 253
for gastrostomy, 233
for gastrotomy, 235
Gély's, 229
glover's, 223
hare-lip, 224
India-rubber, 224
for intestinal anastomosis, 232
interrupted, 222
Jobert's, 231
Lembert's, 228

Sutures, plate, 226
quilled, 225
quilt, 224
of relaxation, 217
removal of, 228
secondary, 217
securing of, 219
shotted, 227
tongue-and-groove, 227
twisted, 224
Sylvester's method of artificial respiration, 174
Syme's amputation at ankle-joint, 464
Syringe, hypodermic, 183

T-BANDAGE, 23
of chest, double, 26
single, 24, 25
double, 25
of groin, 24, 25
of nose, double, 26
single, 23
Tarsal bones, dislocation of, 388, 389
Tarso-metatarsal amputations, 460
Teale's amputation, 432
Temporal artery, ligation of, 404
Tenaculum, 252, 436
Tent, 133
Testicle, strapping of, 136
Thermo-cautery, Paquelin's, 156
Thermometer, clinical, 189
surface, 189
Thigh, amputation of, 475 *et seq.*
Three-tailed retractors, 133
Thumb, dislocation of proximal phalanx of, 380
spica bandage of, 49
Thyroid artery, inferior, ligation of, 399
superior, ligation of, 403
dislocation of femur, 383
Tibial artery, anterior, ligation of, 422
posterior, ligation of, 424
Tissue, rubber, 114, 152
Toes, amputation of, 456
dislocations of, 390
fractures of, 356
Tongue-and-groove suture, 227
Torsion in arterial hemorrhage, 251
forceps, 251
Tourniquets, 243, 438
abdominal, 477
Petit's, 243
Signorini's, 245
Trachea, fracture of, 308
Tracheal dilators, 277
Golding-Bird's, 277
Trousseau's 277
forceps, 278
Tracheotomy, 275
anæsthetics in, 279
director, 276
for foreign bodies, 284
position of patient in, 279
operation of, 279
tubes, 278
Transfusion, arterial, 167
of blood, 164
direct, 165
indirect, 165
Trapping of plaster-of-Paris bandage, 92
Transverse fracture, 293
Tripier's amputation of foot, 467
Trocar, exploring, 184
Trousseau's tracheal dilator, 277
Trunk, bandages of, 59
fractures of, 309
Truss or trusses, 202
application of, 203 *et seq.*
for femoral hernia, 205
Hood's, 204
for inguinal hernia, 204
for irreducible hernia, 206
for umbilical hernia, 205
Tube, rectal, 190
tracheal, 278
Turpentine, 148
stupe, 148
Twisted suture, 224
Two-tailed retractors, 133

ULCERS, strapping of, 138
Ulna, dislocation of lower end of, 377
upper end of, 377
fracture of, 327
coronoid process of, 328
olecranon process of, 327
Ulnar artery, ligation of, 411
Umbilical hernia, truss for, 205
Upper extremity, bandage of, 45
fractures of, 314
spiral reversed bandage of, 50
Urethra, hemorrhage from, 261
Urethral injections, 216
Urethroscope, 215

V-BANDAGE of head, 39
Vaccination, 181
Vascular growths, ligatures for, 235

Vein, jugular, bleeding from, 163
saphena, bleeding from, 164
Velpeau's bandage, 55
dressing for fracture of clavicle, 317
Venesection, 162
Venous hemorrhage, 241
treatment of, 256
Vertebræ, dislocations of, 364
fractures of, 312
Vertebral artery, ligation of, 398
Vesicants, 150

WASHING out the bladder, 215
Water-bed, 312
Waxed paper, 132
Wet cupping, 160
Wooden splints, 299
Wounds, aseptic, dressing of, 101
Wounds, chemical sterilization in, 101
contused, 268
dressing of, 266
dry dressing in, 101
gunshot, 269
incised, 266
lacerated, 267
moist dressings in, 102
modified, 102
poisoned, 269
punctured, 268
septic, dressing of, 128
Wrist, amputation at, 446
dislocations of, 377

ZINC, chloride of, 106
sulpho-carbolate of, 106
Zygoma, fracture of, 304

ALLEN (HARRISON). A SYSTEM OF HUMAN ANATOMY. WITH AN INTRODUCTORY SECTION ON HISTOLOGY, by E. O. Shakespeare, M.D. Comprising 813 double-columned quarto pages, with 380 engravings on stone on 109 plates, and 241 woodcuts in the text. In six sections, each in a portfolio. Section I. (Histology), Section II. (Bones and Joints), Section III. (Muscles and Fasciæ), Section IV. (Arteries, Veins and Lymphatics), Section V. (Nervous System), Section VI. (Organs of Sense, of Digestion and Genito-Urinary Organs, Embryology, Development, Teratology, Post-Mortem Examinations, General and Clinical Indexes). Price per section, $3 50. Also, bound in one volume, cloth, $23. *Sold by subscription only*.

AMERICAN SYSTEM OF DENTISTRY. In treatises by various authors. Edited by Wilbur F. Litch, M.D., D.D.S. In three very handsome super-royal octavo volumes, containing 3180 pages, with 2863 illustrations and 9 full-page plates. *Now ready*. Per volume, cloth, $6; leather, $7; half Morocco, $8. *For sale by subscription only*. Apply to the publishers.

AMERICAN SYSTEMS OF GYNECOLOGY AND OBSTETRICS. In treatises by the most eminent American specialists. Gynecology edited by Matthew D. Mann, A.M., M.D., and Obstetrics edited by Barton C. Hirst, M.D. In four large octavo volumes comprising 3612 pages, with 1092 engravings, and 8 colored plates. Per volume, cloth, $5; leather, $6; half Russia, $7. *For sale by subscription only*. Prospectus free on application to publishers.

ASHHURST (JOHN, Jr.) THE PRINCIPLES AND PRACTICE OF SURGERY. FOR THE USE OF STUDENTS AND PRACTITIONERS. New (fifth) and revised edition. In one large and handsome octavo volume of 1144 pages, with 642 woodcuts. Cloth, $6; leather, $7.

ASHWELL (SAMUEL). A PRACTICAL TREATISE ON THE DISEASES OF WOMEN. Third edition. 520 pages. Cloth, $3 50.

A SYSTEM OF PRACTICAL MEDICINE BY AMERICAN AUTHORS. Edited by William Pepper, M.D., LL.D. In five large octavo volumes, containing 5573 pages and 198 illustrations. Price per volume, cloth, $5 00; leather, $6 00; half Russia, $7 00. *Sold by subscription only*. Address the publishers.

ATTFIELD (JOHN). CHEMISTRY; GENERAL, MEDICAL AND PHARMACEUTICAL. Twelfth edition, specially revised by the Author for America. In one handsome 12mo. volume of 782 pages, with 88 illustrations. Cloth, $2 75; leather, $3 25.

BALL (CHARLES B.) DISEASES OF THE RECTUM AND ANUS. In one 12mo. vol. of 417 pages, with 54 illus. and 4 colored plates. Cloth, $2 25. See *Series of Clinical Manuals*, p. 13.

BARLOW (GEORGE H.) A MANUAL OF THE PRACTICE OF MEDICINE. In one 8vo. volume of 603 pages. Cloth, $2 50.

BARNES (ROBERT). A PRACTICAL TREATISE ON THE DISEASES OF WOMEN. Third American from 3d English edition. In one 8vo. vol. of about 800 pages, with about 200 illus. *Preparing*.

BARNES (ROBERT and FANCOURT). A SYSTEM OF OBSTETRIC MEDICINE AND SURGERY, THEORETICAL AND CLINICAL. The Section on Embryology by Prof. Milnes Marshall In one large octavo volume of 872 pages, with 231 illustrations Cloth, $5; leather, $6.

Catalogue of Books

PUBLISHED BY

LEA BROTHERS AND COMPANY.

The books in the annexed list will be sent by mail, post-paid, to any Post Office in the United States, on receipt of the printed prices. No risks of the mail, however, are assumed, either on money or books. Gentlemen will, therefore, in most cases, find it more convenient to deal with the nearest bookseller.

LEA BROTHERS & CO.,
Nos. 706 and 708 Sansom Street, Philadelphia.

Periodicals. 1891.

THE MEDICAL NEWS,

A WEEKLY JOURNAL OF MEDICAL SCIENCE,

EDITED BY GEORGE M. GOULD, M.D.

Published every Saturday, containing 24-28 large double-columned quarto pages of reading matter in each number.

FOUR DOLLARS ($4) per annum, in advance.

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES.

EDITED BY EDWARD P. DAVIS, A.M., M.D.

Published every Month.

112 large octavo pages, fully illustrated in each number. Price

REDUCED TO FOUR DOLLARS PER ANNUM.

COMBINATIONS AT REDUCED RATES.

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES, $4.00 THE MEDICAL NEWS, $4.00	Together $7.50
THE MEDICAL NEWS VISITING LIST for 1891 (see below and on page 16), $1.25 With either or both above periodicals, 75 cents. THE YEAR-BOOK OF TREATMENT for 1891 (see page 16), $1.50 With either or both above periodicals, 75 cents.	Together $8.50

THE MEDICAL NEWS VISITING LIST.

This LIST, which is by far the most handsome and convenient now attainable, has been thoroughly revised for 1891. A full description will be found on page 16. It is issued in four styles. Price, each, $1 25. For Special Combination Rates with periodicals and the Year-Book of Treatment see above. Thumb letter Index for quick use 25 cents extra.

BUTLIN, (HENRY T.) DISEASES OF THE TONGUE. In one pocket-size 12mo. vol. of 456 pp., with 8 col. plates and 3 woodcuts. Limp cloth, $3 50. See *Series of Clinical Manuals*, p. 13.

CARPENTER (WM. B) PRIZE ESSAY ON THE USE OF ALCOHOLIC LIQUORS IN HEALTH AND DISEASE. New Edition, with a Preface by D. F. Condie, M.D. One 12mo. volume of 178 pages. Cloth, 60 cents.

—— PRINCIPLES OF HUMAN PHYSIOLOGY. A new American, from the eighth English edition. In one large 8vo. volume.

CARTER (R. BRUDENELL) AND FROST (W. ADAMS) OPHTHALMIC SURGERY. In one pocket-size 12mo. volume of 559 pages, with 91 engravings and one plate. Cloth, $2 25. See *Series of Clinical Manuals*, p. 13.

CHAMBERS (T. K.) A MANUAL OF DIET IN HEALTH AND DISEASE. In one handsome 8vo. vol. of 302 pages. Cloth, $2 75.

CHAPMAN (HENRY C). A TREATISE ON HUMAN PHYSIOLOGY. In one octavo volume of 925 pages, with 605 illustrations. Cloth, $5 50; leather, $6 50.

CHARLES (T. CRANSTOUN). THE ELEMENTS OF PHYSIOLOGICAL AND PATHOLOGICAL CHEMISTRY. In one handsome octavo volume of 451 pages, with 38 woodcuts and one colored plate. Cloth, 3 50.

CHURCHILL (FLEETWOOD). ESSAYS ON THE PUERPERAL FEVER. In one octavo volume of 464 pages. Cloth, $2 50.

CLARKE (W. B.) AND LOCKWOOD (C. B.) THE DISSECTOR'S MANUAL. In one 12mo. volume of 396 pages, with 49 illustrations. Cloth, $1 50. See *Students' Series of Manuals*, p. 14.

CLASSEN'S QUANTITATIVE ANALYSIS. Translated by Edgar F. Smith, Ph.D. In one 12mo. vol. of 324 pp., with 36 illus. Cloth, $2 00.

CLELAND (JOHN). A DIRECTORY FOR THE DISSECTION OF THE HUMAN BODY. In one 12mo. vol. of 178 pp. Cloth, $1 25.

CLOUSTON (THOMAS S.) CLINICAL LECTURES ON MENTAL DISEASES. With an Abstract of Laws of U. S. on Custody of the Insane, by C. F. Folsom, M.D. In one handsome octavo vol. of 541 pages, illustrated with woodcuts and 8 lithographic plates. Cloth, $4 00. Dr. Folsom's *Abstract* is also furnished separately in one octavo volume of 108 pages. Cloth, $1 50.

CLOWES (FRANK). AN ELEMENTARY TREATISE ON PRACTICAL CHEMISTRY AND QUALITATIVE INORGANIC ANALYSIS. New American from the fourth English edition. In one handsome 12mo. volume of 387 pages, with 55 illustrations. Cloth, $2 50.

COATS (JOSEPH). A TREATISE ON PATHOLOGY. In one vol. of 829 pp., with 339 engravings. Cloth, $5 50; leather, $6 50.

COHEN (J. SOLIS). DISEASES OF THE THROAT AND NASAL PASSAGES. Third edition, thoroughly revised. In one handsome octavo volume. *Preparing.*

COLEMAN (ALFRED). A MANUAL OF DENTAL SURGERY AND PATHOLOGY. With Notes and Additions to adapt it to American Practice. By Thos. C. Stellwagen, M.A., M.D., D.D.S. In one handsome 8vo. vol. of 412 pp., with 331 illus. Cloth, $3 25.

CONDIE (D. FRANCIS). A PRACTICAL TREATISE ON THE DISEASES OF CHILDREN. Sixth edition, revised and enlarged. In one large 8vo. vol. of 719 pages. Cloth, $5 25; leather, $6 25.

COOPER (B.B.) LECTURES ON THE PRINCIPLES AND PRACTICE OF SURGERY. In one large 8vo. vol. of 767 pages. Cloth, $2 00.

CORNIL (V.) SYPHILIS: ITS MORBID ANATOMY, DIAGNOSIS AND TREATMENT. Translated, with notes and additions, by J. Henry C. Simes, M.D., and J. William White, M.D. In one 8vo. volume of 461 pages, with 84 illustrations. Cloth, $3 75.

CULLERIER (A.) AN ATLAS OF VENEREAL DISEASES. Translated and edited by FREEMAN J. BUMSTEAD, M.D., LL.D. A large quarto volume of 328 pages, with 26 plates containing about 150 figures, beautifully colored, many of them life-size. Cloth, $17.

DALTON (JOHN C.) DOCTRINES OF THE CIRCULATION OF THE BLOOD. In one handsome 12mo. vol. of 293 pp. Cloth, $2.

——— A TREATISE ON HUMAN PHYSIOLOGY. Seventh edition, thoroughly revised, and greatly improved. In one very handsome 8vo. vol. of 722 pages, with 252 illustrations. Cloth, $5; leather, $6.

DANA (JAMES D.) THE STRUCTURE AND CLASSIFICATION OF ZOOPHYTES. With illust. on wood. In one imp. 4to. vol. Cl., $4.

DAVENPORT (F. H.) DISEASES OF WOMEN. A Manual of Non-Surgical Gynæcology. For the use of Students and General Practitioners In one handsome 12mo. volume of 306 pages with 105 illustrations. Cloth, $1 50.

DAVIS (F. H.) LECTURES ON CLINICAL MEDICINE. Second edition In one 12mo. volume of 287 pages. Cloth, $1 75.

DE LA BECHE'S GEOLOGICAL OBSERVER. In one large 8vo. vol. of 700 pages, with 300 illustrations. Cloth, $4.

DRAPER (JOHN C.) MEDICAL PHYSICS. A Text-book for Students and Practitioners of Medicine. In one handsome octavo volume of 734 pages, with 376 illustrations. Cloth, $4.

DRUITT (ROBERT). THE PRINCIPLES AND PRACTICE OF MODERN SURGERY. A new American from the 12th London edition, edited by Stanley Boyd, F.R C.S. In one large octavo volume of 965 pages, with 373 illustrations. Cloth, $4; leather, $5.

DUNCAN (J. MATTHEWS). CLINICAL LECTURES ON THE DISEASES OF WOMEN. Delivered in St. Bartholomew's Hospital. In one octavo volume of 175 pages. Cloth, $1 50.

DUNGLISON (ROBLEY). MEDICAL LEXICON; A Dictionary of Medical Science. Containing a concise explanation of the various subjects and terms of Anatomy, Physiology, Pathology, Hygiene, Therapeutics, Pharmacology, Pharmacy, Surgery, Obstetrics, Medical Jurisprudence and Dentistry; notices of Climate and of Mineral Waters; Formulæ for Officinal, Empirical and Dietetic Preparations; with the accentuation and Etymology of the Terms, and the French and other Synonymes. Edited by R. J. Dunglison, M.D. In one very large royal 8vo. vol. of 1139 pages. Cloth, $6 50; leather, $7 50; half Russia, $8.

EDES' TEXT-BOOK OF THERAPEUTICS AND MATERIA MEDICA. In one 8vo. volume of 544 pages. Cloth, $3 50; leather, $4 50.

EDIS (ARTHUR W.) DISEASES OF WOMEN. A Manual for Students and Practitioners. In one handsome 8vo. vol. of 576 pp., with 148 illustrations. Cloth, $3; leather, $4.

ELLIS (GEORGE VINER). DEMONSTRATIONS IN ANATOMY. Being a Guide to the Knowledge of the Human Body by Dissection. From the eighth and revised English edition. In one octavo vol. of 716 pages, with 249 illustrations. Cloth, $4 25; leather, $5 25.

EMMET (THOMAS ADDIS). THE PRINCIPLES AND PRACTICE OF GYNÆCOLOGY, for the use of Students and Practitioners. Third edition, enlarged and revised. In one large 8vo. volume of 880 pages, with 150 original illustrations. Cloth, $5; leather, $6.

ERICHSEN (JOHN E.) THE SCIENCE AND ART OF SURGERY. A new American, from the eighth enlarged and revised London edition. In two large octavo volumes containing 2316 pages, with 984 illus. Cloth, $9; leather, $11.

FARQUHARSON (ROBERT). A GUIDE TO THERAPEUTICS. Fourth American from Fourth English edition, revised by Frank Woodbury, M.D. In one 12mo. volume of 581 pages. Cloth, $2 50.

FINLAYSON (JAMES). CLINICAL DIAGNOSIS. A Handbook for Students and Practitioners of Medicine. Second edition. In one 12mo. volume of 682 pages, with 158 illustrations. Cloth, $2 50.

FLINT (AUSTIN). A TREATISE ON THE PRINCIPLES AND PRACTICE OF MEDICINE. Sixth edition, thoroughly revised and largely rewritten by the Author, assisted by William H. Welch, M.D., and Austin Flint, Jr., M.D. In one large 8vo. volume of 1160 pages, with illustrations. Cloth, $5 50; leather, $6 50.

—— A MANUAL OF AUSCULTATION AND PERCUSSION; of the Physical Diagnosis of Diseases of the Lungs and Heart, and of Thoracic Aneurism. Fifth edition, revised by James C. Wilson, M.D. In one handsome 12mo. volume of 274 pages, with 12 illustrations. Cloth, $1 75. *Just ready.*

—— A PRACTICAL TREATISE ON THE DIAGNOSIS AND TREATMENT OF DISEASES OF THE HEART. Second edition, enlarged. In one octavo volume of 550 pages. Cloth, $4 00.

—— A PRACTICAL TREATISE ON THE PHYSICAL EXPLORATION OF THE CHEST, AND THE DIAGNOSIS OF DISEASES AFFECTING THE RESPIRATORY ORGANS. Second and revised edition. In one octavo volume of 591 pages. Cloth, $4 50.

—— MEDICAL ESSAYS. In one 12mo. vol., pp. 210. Cloth, $1 38.

—— ON PHTHISIS: ITS MORBID ANATOMY, ETIOLOGY, ETC. A series of Clinical Lectures. In one 8vo. volume of 442 pages. Cloth, $3 50.

FOLSOM (C. F.) An Abstract of Statutes of U. S. on Custody of the Insane. In one 8vo. vol. of 108 pp. Cloth, $1 50. Also bound with *Clouston on Insanity.*

FOSTER (MICHAEL). A TEXT-BOOK OF PHYSIOLOGY. Fourth and revised American from the fifth English edition. In one large octavo vol. of about 925 pages, with about 300 illust. *In press.*

FOTHERGILL'S PRACTITIONER'S HANDBOOK OF TREATMENT. Third edition. In one handsome octavo volume of 664 pages. Cloth, $3 75; leather, $4 75.

FOWNES (GEORGE). A MANUAL OF ELEMENTARY CHEMISTRY (INORGANIC AND ORGANIC). New edition. Embodying Watts' *Physical and Inorganic Chemistry.* In one royal 12mo. vol. of 1061 pages, with 168 illus., and one colored plate. Cloth, $2 75; leather, $3 25.

FOX (TILBURY) and T. COLCOTT. EPITOME OF SKIN DISEASES, with Formulæ. For Students and Practitioners. Third Am. edition, revised by T. C. Fox. In one small 12mo. volume of 238 pages. Cloth, $1 25.

FRANKLAND (E.) and JAPP (F. R.) INORGANIC CHEMISTRY. In one handsome octavo vol. of 677 pages, with 51 engravings and 2 plates. Cloth, $3 75; leather, $4 75.

FULLER (HENRY). ON DISEASES OF THE LUNGS AND AIR PASSAGES. Their Pathology, Physical Diagnosis, Symptoms and Treatment. From 2d Eng. ed In 1 8vo. vol., pp. 475. Cloth, $3 50.

GANT (FREDERICK JAMES). THE STUDENT'S SURGERY. A Multum in Parvo. In one square octavo volume of 845 pages, with 159 engravings. Cloth, $3 75.

GIBBES (HENEAGE). PRACTICAL PATHOLOGY. In one very handsome octavo volume of 314 pages, with 60 illustrations, mostly photographic. Cloth, $2 75. *Just ready.*

GIBNEY (V. P.) ORTHOPÆDIC SURGERY. For the use of Practitioners and Students. In one 8vo. vol. profusely illus. *Prepg.*

GIBSON'S INSTITUTES AND PRACTICE OF SURGERY. In two octavo volumes of 965 pages, with 34 plates. Leather, $6 50.

GOULD (A. PEARCE). SURGICAL DIAGNOSIS. In one 12mo. vol. of 589 pages. Cloth, $2 See *Students' Series of Manuals*, p. 14.

GRAY (HENRY). ANATOMY, DESCRIPTIVE AND SURGICAL. Edited by T. Pickering Pick, F.R.C.S. A new American, from the eleventh English edition, thoroughly revised, with additions, by W. W. Keen, M.D. To which is added Holden's "Landmarks, Medical and Surgical." In one imperial octavo volume of 1098 pages, with 685 large and elaborate engravings on wood. Cloth, $6; leather, $7; very handsome half Russia, raised bands, $7 50. The same edition is also issued with veins, arteries, and nerves distinguished in colors. Price, cloth, $7 25; leather, $8 25; half Russia, $8 75.

GRAY (LANDON CARTER). A PRACTICAL TREATISE ON THE DISEASES OF THE NERVOUS SYSTEM. In one handsome octavo volume of about 600 pages. *Preparing.*

GREEN (T. HENRY). AN INTRODUCTION TO PATHOLOGY AND MORBID ANATOMY. New (sixth) American, from the seventh London edition. In one handsome octavo volume of 540 pages, with 167 illustrations. Cloth, $2 75.

GREENE (WILLIAM H.) A MANUAL OF MEDICAL CHEMISTRY. For the Use of Students. Based upon Bowman's Medical Chemistry. In one 12mo. vol. of 310 pages, with 74 illus. Cloth, $1 75.

GRIFFITH (ROBERT E.) A UNIVERSAL FORMULARY, CONTAINING THE METHODS OF PREPARING AND ADMINISTERING OFFICINAL AND OTHER MEDICINES. Third and enlarged edition. Edited by John M. Maisch, Phar.D. In one large 8vo. vol. of 775 pages, double columns. Cloth, $4 50; leather, $5 50.

GROSS (SAMUEL D.) A SYSTEM OF SURGERY, PATHOLOGICAL, DIAGNOSTIC, THERAPEUTIC AND OPERATIVE. Sixth edition, thoroughly revised. In two imperial octavo volumes containing 2382 pages, with 1623 illustrations. Strongly bound in leather, raised bands, $15.

—— A PRACTICAL TREATISE ON THE DISEASES, INJURIES and Malformations of the Urinary Bladder, the Prostate Gland and the Urethra. Third edition, thoroughly revised and much condensed, by Samuel W. Gross, M.D. In one octavo volume of 574 pages, with 170 illus. Cloth, $4 50.

—— A PRACTICAL TREATISE ON FOREIGN BODIES IN THE AIR PASSAGES. In one 8vo. vol. of 468 pages. Cloth, $2 75.

GROSS (SAMUEL W.) A PRACTICAL TREATISE ON IMPOTENCE, STERILITY, AND ALLIED DISORDERS OF THE MALE SEXUAL ORGANS. New (fourth) edition. Edited by F. R. Sturgis, M.D. In one handsome octavo volume of 165 pages, with 18 illustrations. Cloth, $1.50.

HABERSHON (S. O.) ON THE DISEASES OF THE ABDOMEN, AND OTHER PARTS OF THE ALIMENTARY CANAL. Second American, from the third English edition. In one handsome 8vo. volume of 554 pages, with illus. Cloth, $3 50.

HAMILTON (ALLAN McLANE). NERVOUS DISEASES, THEIR DESCRIPTION AND TREATMENT. Second and revised edition. In one octavo volume of 598 pages, with 72 illustrations. Cloth, $4.

HAMILTON (FRANK H.) A PRACTICAL TREATISE ON FRACTURES AND DISLOCATIONS. New (eighth) edition, revised and edited by Stephen Smith, A.M., M.D. In one handsome 8vo. vol. of 832 pages, with 507 illustrations. Cloth, $5 50; leather, $6 50. *Just ready.*

HARE (HOBART AMORY). A TEXT-BOOK OF PRACTICAL THERAPEUTICS, with Special Reference to the Application of Remedial Measures to Disease and their Employment upon a Rational Basis. With articles on various subjects by well-known specialists. New (second) edition. In one handsome octavo volume of 650 pages. Cloth, $3 75; leather, $4.75. *Just ready.*

HARTSHORNE (HENRY). ESSENTIALS OF THE PRINCIPLES AND PRACTICE OF MEDICINE. Fifth edition. In one 12mo. volume, 669 pages, with 144 illustrations. Cloth, $2 75; half bound, $3.

—— A HANDBOOK OF ANATOMY AND PHYSIOLOGY. In one 12mo. volume of 310 pages, with 220 illustrations. Cloth, $1 75.

—— A CONSPECTUS OF THE MEDICAL SCIENCES. Comprising Manuals of Anatomy, Physiology, Chemistry, Materia Medica, Practice of Medicine, Surgery and Obstetrics. Second edition. In one royal 12mo. volume of 1028 pages, with 477 illustrations. Cloth, $4 25; leather, $5 00.

HERMANN (L.) EXPERIMENTAL PHARMACOLOGY. A Handbook of the Methods for Determining the Physiological Actions of Drugs. Translated by Robert Meade Smith, M.D. In one 12mo. vol. of 199 pages, with 32 illustrations. Cloth, $1 50.

HILL (BERKELEY). SYPHILIS AND LOCAL CONTAGIOUS DISORDERS. In one 8vo. volume of 479 pages. Cloth, $3 25.

HILLIER (THOMAS). A HANDBOOK OF SKIN DISEASES. 2d ed. In one royal 12mo. vol. of 353 pp., with two plates. Cloth, $2 25.

HIRST (BARTON C.) AND PIERSOL (GEORGE A.) HUMAN MONSTROSITIES. Magnificent folio, containing about 120 pages of text and illustrated with engravings and about 60 large photograhic plates from nature. In four parts, price each, $5. *Limited edition, for sale by subscription only. In press.*

HOBLYN (RICHARD D.) A DICTIONARY OF THE TERMS USED IN MEDICINE AND THE COLLATERAL SCIENCES. In one 12mo. vol. of 520 double-columned pp. Cloth, $1 50; leather, $2.

HODGE (HUGH L.) ON DISEASES PECULIAR TO WOMEN, INCLUDING DISPLACEMENTS OF THE UTERUS. Second and revised edition. In one 8vo. volume of 519 pages. Cloth, $4 50.

HOFFMANN (FREDERICK) AND POWER (FREDERICK B.) A MANUAL OF CHEMICAL ANALYSIS, as Applied to the Examination of Medicinal Chemicals and their Preparations. Third edition, entirely rewritten and much enlarged. In one handsome octavo volume of 621 pages, with 179 illustrations. Cloth, $4 25.

HOLDEN (LUTHER). LANDMARKS, MEDICAL AND SURGICAL. From the third English edition. With additions, by W. W. Keen, M.D. In one royal 12mo. vol. of 148 pp. Cloth, $1.

HOLLAND (SIR HENRY). MEDICAL NOTES AND REFLECTIONS. From 3d English ed. In one 8vo. vol. of 493 pp. Cloth, $3 50.

HOLMES (TIMOTHY). A SYSTEM OF SURGERY. With notes and additions by various American authors. Edited by John H. Packard, M.D. In three very handsome 8vo. vols. containing 3137 double-columned pages, with 979 woodcuts and 13 lithographic plates. Cloth, $18; leather, $21; very handsome half Russia, raised bands, $22 50. *For sale by subscription only.*

—— A TREATISE ON SURGERY. Its Principles and Practice. A new American from the fifth English edition. Edited by T. Pickering Pick, F.R.C.S. In one handsome octavo volume of 1008 pages, with 428 engravings. Cloth, $6; leather, $7.

HORNER (WILLIAM E.) SPECIAL ANATOMY AND HISTOLOGY. Eighth edition, revised and modified. In two large 8vo. vols. of 1007 pages, containing 320 woodcuts. Cloth, $6.

HUDSON (A.) LECTURES ON THE STUDY OF FEVER. In one octavo volume of 308 pages. Cloth, $2 50.

HUTCHINSON (JONATHAN). SYPHILIS. In one pocket size 12mo. volume of 542 pages, with 8 chromo-lithographic plates. Cloth, $2 25. See *Series of Clinical Manuals*, p. 13

HYDE (JAMES NEVINS). A PRACTICAL TREATISE ON DISEASES OF THE SKIN. New (second) edition. In one handsome octavo volume of 676 pages, with 85 engravings and 2 colored plates. Cloth, $4 50; leather, $5 50.

JONES (C. HANDFIELD). CLINICAL OBSERVATIONS ON FUNCTIONAL NERVOUS DISORDERS. Second American edition. In one octavo volume of 340 pages. Cloth, $3 25.

JULER (HENRY) A HANDBOOK OF OPHTHALMIC SCIENCE AND PRACTICE. In one 8vo. volume of 460 pages, with 125 woodcuts, 27 chromo-lithographic plates test types of Jaeger and Snellen and Holmgren's Color-blindness test. Cloth, $4 50; leather, $5 50.

KING (A. F. A.) A MANUAL OF OBSTETRICS. New (fourth) edition. In one 12mo. volume of 432 pages, with 141 illustrations. Cloth, $2 50.

KLEIN (E.) ELEMENTS OF HISTOLOGY. Fourth edition. In one pocket-size 12mo. volume of 376 pages, with 194 engravings. Cloth, $1 75. See *Students' Series of Manuals*, p. 14.

LANDIS (HENRY G.) THE MANAGEMENT OF LABOR. In one handsome 12mo. volume of 329 pages, with 28 illus. Cloth, $1 75.

LA ROCHE (R.) YELLOW FEVER. In two 8vo. vols. of 1468 pages. Cloth, $7.

—— PNEUMONIA. In one 8vo. vol. of 490 pages. Cloth, $3.

LAURENCE (J. Z.) AND MOON (ROBERT C.) A HANDY-BOOK OF OPHTHALMIC SURGERY. Second edition, revised by Mr. Laurence. In one 8vo. vol. pp. 227, with 66 illus. Cloth, $2 75.

LAWSON (GEORGE). INJURIES OF THE EYE, ORBIT AND EYELIDS. From the last English edition. In one handsome octavo volume of 404 pages, with 92 illustrations. Cloth, $3 50.

LEA (HENRY C.). CHAPTERS FROM THE RELIGIOUS HISTORY OF SPAIN; CENSORSHIP OF THE PRESS; MYSTICS AND ILLUMINATI; THE ENDEMONIADAS; EL SANTO NIÑO DE LA GUARDIA; BRIANDA DE BARDAXI. In one 12mo. volume of 522 pages. Cloth, $2.50.

—— SUPERSTITION AND FORCE; ESSAYS ON THE WAGER OF LAW, THE WAGER OF BATTLE, THE ORDEAL AND TORTURE. Third edition, thoroughly revised and greatly enlarged. In one handsome royal 12mo. vol. pp. 552. Cloth, $2 50.

—— STUDIES IN CHURCH HISTORY. The Rise of the Temporal Power—Benefit of Clergy—Excommunication. New edition. In one handsome 12mo. vol. of 605 pp. Cloth, $2 50.

—— AN HISTORICAL SKETCH OF SACERDOTAL CELIBACY IN THE CHRISTIAN CHURCH. Second edition. In one handsome octavo volume of 684 pages. Cloth, $4 50.

LEE (HENRY) ON SYPHILIS. In one 8vo volume of 246 pages. Cloth, $2 25.

LEHMANN (C. G.) A MANUAL OF CHEMICAL PHYSIOLOGY. In one 8vo. vol. of 327 pages, with 41 woodcuts. Cloth, $2 25.

LEISHMAN (WILLIAM). A SYSTEM OF MIDWIFERY. Including the Diseases of Pregnancy and the Puerperal State. Fourth edition. In one octavo volume of about 800 pages, with about 225 illustrations.

LUCAS (CLEMENT). DISEASES OF THE URETHRA. *Preparing*. See *Series of Clinical Manuals*, p. 13.

LUDLOW (J. L.) A MANUAL OF EXAMINATIONS UPON ANATOMY, PHYSIOLOGY, SURGERY, PRACTICE OF MEDICINE, OBSTETRICS, MATERIA MEDICA, CHEMISTRY, PHARMACY AND THERAPEUTICS. To which is added a Medical Formulary. Third edition. In one royal 12mo. volume of 816 pages, with 370 woodcuts. Cloth, $3 25; leather, $3 75.

LYONS (ROBERT D.) A TREATISE ON FEVER. In one octavo volume of 362 pages. Cloth, $2 25.

CH (JOHN M.) A MANUAL OF ORGANIC MATERIA MED. New (fourth) edition. In one handsome 12mo. volume of es, with 258 beautiful illustrations. Cloth, $3.

MARSH (HOWARD). DISEASES OF THE JOINTS. In one 12mo. volume of 468 pages, with 64 illustrations and a colored plate. Cloth, $2. See *Series of Clinical Manuals*, p. 13.

MAY (C. H.) MANUAL OF THE DISEASES OF WOMEN. For the use of Students and Practitioners. New (second) edition, revised by L S. Rau, M.D. In one 12mo. volume of 360 pages, with 31 illustrations. Cloth, $1 75.

MEIGS (CHAS. D.) ON THE NATURE, SIGNS AND TREATMENT OF CHILDBED FEVER. In one 8vo. vol. of 346 pages. Cloth, $2.

MILLER (JAMES). PRINCIPLES OF SURGERY. Fourth American, from the third Edinburgh edition. In one large octavo volume of 688 pages, with 240 illustrations. Cloth, $3 75.

MILLER (JAMES). THE PRACTICE OF SURGERY. Fourth American, from the last Edinburgh edition. In one large octavo volume of 682 pages, with 364 illustrations. Cloth, $3 75.

MORRIS (HENRY). SURGICAL DISEASES OF THE KIDNEY. 12mo., 554 pages, 40 woodcuts, and 6 colored plates. Cloth, $2 25. See *Series of Clinical Manuals*, p. 13.

MÜLLER (J.) PRINCIPLES OF PHYSICS AND METEOROLOGY. In one large 8vo. vol. of 623 pages, with 538 cuts. Cloth, $4 50.

NEILL (JOHN) AND SMITH (FRANCIS G.) A COMPENDIUM OF THE VARIOUS BRANCHES OF MEDICAL SCIENCE. In one handsome 12mo. volume of 974 pages, with 374 woodcuts. Cloth, $4; leather, raised bands, $4 75.

NETTLESHIP (E.) DISEASES OF THE EYE. New (fourth) American, from fifth English edition. In one royal 12mo. volume of 500 pages, with 164 illustrations, test-types and formulæ and color blindness test. Cloth, $2.

NORRIS AND OLIVER ON THE EYE. In one 8vo. volume of about 500 pages, with illustrations. *In press.*

OWEN (EDMUND). SURGICAL DISEASES OF CHILDREN. 12mo., 525 pages, 85 woodcuts, and 4 colored plates. Cloth, $2. See *Series of Clinical Manuals*, p. 13.

PARRISH (EDWARD). A TREATISE ON PHARMACY. With many Formulæ and Prescriptions. Fifth edition, enlarged and thoroughly revised by Thomas S. Wiegand, Ph.G. In one octavo volume of 1093 pages, with 257 illustrations. Cloth, $5; leather, $6.

PARRY (JOHN S.) EXTRA-UTERINE PREGNANCY, ITS CLINICAL HISTORY, DIAGNOSIS, PROGNOSIS AND TREATMENT. In one octavo volume of 272 pages. Cloth, $2 50.

PARVIN (THEOPHILUS). THE SCIENCE AND ART OF OBSTETRICS. New (second) edition. In one handsome 8vo. volume of 701 pages, with 239 engravings and a colored plate. Cloth, $4 25; leather, $5 25.

PAVY (F. W.) A TREATISE ON THE FUNCTION OF DIGESTION, ITS DISORDERS AND THEIR TREATMENT. From the second London edition. In one octavo volume of 238 pages. Cloth, $2.

PAYNE (JOSEPH FRANK). A MANUAL OF GENERAL PATHOLOGY. Designed as an Introduction to the Practice of Medicine. Handsome octavo volume of 524 pages with 153 engravings and 1 colored plate. Cloth, $3 50.

PEPPER (A. J.) FORENSIC MEDICINE. *In press.* See *Students' Series of Manuals*, p. 14.

—— SURGICAL PATHOLOGY. In one 12mo. volume of 511 pages, with 81 illus. Cloth, $2. See *Students' Series of Manuals*, p. 14.

PICK (T. PICKERING). FRACTURES AND DISLOCATIONS. In one 12mo. volume of 530 pages, with 93 illustrations. Cloth, $2. *See Series of Clinical Manuals*, p. 13.

PIRRIE (WILLIAM). THE PRINCIPLES AND PRACTICE OF SURGERY. In one handsome octavo volume of 780 pages, with 316 illustrations. Cloth, $3 75.

PLAYFAIR (W. S.) A TREATISE ON THE SCIENCE AND PRACTICE OF MIDWIFERY. New (fifth) American from the seventh English edition. Edited, with additions, by R. P. Harris, M.D. In one octavo volume of 664 pages, with 207 woodcuts and five plates. loth, $4; leather, $5.

—— THE SYSTEMATIC TREATMENT OF NERVE PROSTRATION AND HYSTERIA. In one 12mo. vol. of 97 pages. Cloth, $1.

POWER (HENRY). HUMAN PHYSIOLOGY. Second edition. In one 12mo. volume of 396 pages, with 47 illustrations. Cloth, $1 50. See *Students' Series of Manuals*, page 14.

PURDY ON BRIGHT'S DISEASE AND ALLIED AFFECTIONS OF THE KIDNEY. Octavo, 288 pp., with 18 handsome illus. Cloth, $2.

RALFE (CHARLES H.) CLINICAL CHEMISTRY. In one 12mo. volume of 314 pages, with 16 illustrations. Cloth, $1 50. See *Students' Series of Manuals*, page 14.

RAMSBOTHAM (FRANCIS H.) THE PRINCIPLES AND PRACTICE OF OBSTETRIC MEDICINE AND SURGERY. In one imperial octavo volume of 640 pages, with 64 plates, besides numerous woodcuts in the text. Strongly bound in leather, $7.

REMSEN (IRA). THE PRINCIPLES OF THEORETICAL CHEMISTRY. New (third) edition, thoroughly revised, and much enlarged. In one 12mo. volume of 318 pages. Cloth, $2.

REYNOLDS (J. RUSSELL). A SYSTEM OF MEDICINE. Edited, with Notes and Additions, by HENRY HARTSHORNE, M.D. In three large 8vo. vols., containing 3056 closely printed double-columned pages, with 317 illustrations. Per volume, cloth, $5; leather, $6; very handsome half Russia, $6 50. *For sale by subscription only.*

RICHARDSON (BENJAMIN W.) PREVENTIVE MEDICINE. In one octavo vol., of 729 pp. Clo., $4; leather, $5.

ROBERTS (JOHN B). THE PRINCIPLES AND PRACTICE OF MODERN SURGERY. In one octavo volume of 780 pages, with 501 illustrations. Cloth, $4 50; leather, $5 50.

ROBERTS (JOHN B.) THE COMPEND OF ANATOMY. For use in the Dissecting Room and in preparing for Examinations. In one 16mo. volume of 196 pages. Limp cloth, 75 cents.

ROBERTS (SIR WILLIAM). A PRACTICAL TREATISE ON URINARY AND RENAL DISEASES, INCLUDING URINARY DEPOSITS. Fourth American, from the fourth London edition. In one very handsome 8vo. volume of 609 pages, with 81 illustrations. Cloth, $3 50.

COLLECTED CONTRIBUTIONS ON DIET AND DIGESTION. ne 12mo. volume of 270 pages. Cloth, $1 50. *Just ready.*

ROBERTSON (J. McGREGOR). PHYSIOLOGICAL PHYSICS. In one 12mo. volume of 537 pages, with 219 illustrations. Cloth, $2 00. See *Students' Series of Manuals*, p. 14.

ROSS (JAMES) A HANDBOOK OF THE DISEASES OF THE NERVOUS SYSTEM. In one handsome octavo volume of 726 pages, with 184 illustrations. Cloth, $4 50; leather, $5 50.

SAVAGE (GEORGE H.) INSANITY AND ALLIED NEUROSES, PRACTICAL AND CLINICAL. In one 12mo. volume of 551 pages, with 18 typical illustrations. Cloth, $2 00. See *Series of Clinical Manuals*, p 13.

SCHAFER (EDWARD A.) THE ESSENTIALS OF HISTOLOGY, DESCRIPTIVE AND PRACTICAL. For the use of Students. In one handsome octavo volume of 246 pages, with 281 illustrations. Cloth, $2 25.

SCHMITZ AND ZUMPT'S CLASSICAL SERIES. In royal 18mo. ADVANCED LATIN EXERCISES. Cloth, 60 cents; half bound, 70 cents.

SALLUST. Cloth, 60 cents; half bound, 70 cents.
NEPOS. Cloth, 60 cents; half bound, 70 cts.
VIRGIL. Cloth, 85 cents; half bound, $1.
CURTIUS. Cloth, 80 cents; half bound, 90 cents.

SCHREIBER (JOSEPH). A MANUAL OF TREATMENT BY MASSAGE AND METHODICAL MUSCLE EXERCISE. Translated by Walter Mendelson, M.D., of New York. In one handsome octavo volume of 274 pages, with 117 fine engravings. Cloth, $2 75.

SEILER (CARL) A HANDBOOK OF DIAGNOSIS AND TREATMENT OF DISEASES OF THE THROAT AND NASAL CAVITIES. New (3d) edition. In one very handsome 12mo. volume of 373 pages, with 101 illustrations, and 2 beautifully colored plates. Cloth, $2 25.

SENN (NICHOLAS). SURGICAL BACTERIOLOGY. New (second) edition. In one handsome octavo volume of 268 pages, with 13 plates, 9 of which are colored, and 9 engravings. Cloth, $2 00. *Just ready.*

SERIES OF CLINICAL MANUALS. A series of authoritative monographs on important clinical subjects, in 12mo. volumes of about 550 pages, well illustrated. The following volumes are now ready: Broadbent on the Pulse ($1 75); Yeo on Food in Health and Disease ($2); Ball on the Rectum and Anus ($2 25); Carter and Frost's Ophthalmic Surgery ($2 25); Hutchinson on Syphilis ($2 25); Marsh on Diseases of the Joints ($2); Morris on Surgical Diseases of the Kidney ($2 25); Owen on Surgical Diseases of Children ($2); Pick on Fractures and Dislocations ($2); Butlin on the Tongue ($3 50); Savage on Insanity and Allied Neuroses ($2); and Treves on Intestinal Obstruction, ($2). The following is in press: Lucas on Diseases of the Urethra.

For separate notices, see under various authors' names.

SIMON (W.) MANUAL OF CHEMISTRY. A Guide to Lectures and Laboratory work for Beginners in Chemistry. A Text-book specially adapted for Students of Pharmacy and Medicine. New (2d) edition. In one 8vo. volume of 480 pages, with 44 woodcuts and 7 plates showing colors of 56 tests. Cloth, $3 25.

VAUGHAN (VICTOR C.), and NOVY (FRED'K G.) PTOMAINES AND LEUCOMAINES, OR PUTREFACTIVE AND PHYSIOLOGICAL ALKALOIDS. New (second) edition. In one handsome 12mo. volume of about 325 pages. *In press.*

VISITING LIST. THE MEDICAL NEWS VISITING LIST for 1891. Four styles: Weekly (dated for 30 patients); Monthly undated, for 120 patients per month); Perpetual (undated for 30 patients each week); and Perpetual (undated for 60 patients each week). The 60 patient book consists of 256 pages of assorted blanks. The first three styles contain 32 pages of important data, thoroughly revised, and 176 pages of assorted blanks. Each in one vol., price, $1.25. With thumb-letter index for quick use, 25 cents extra. Special rates to advance-paying subscribers to The Medical News or The American Journal, or both. See p. 1.

WALSHE (W. H.) PRACTICAL TREATISE ON THE DISEASES OF THE HEART AND GREAT VESSELS. 3d American from the 3d revised London edition. In one 8vo. vol. of 420 pages Cloth, $3.

WATSON (THOMAS). LECTURES ON THE PRINCIPLES AND PRACTICE OF PHYSIC. A new American from the fifth and enlarged English edition, with additions by H. Hartshorne, M.D. In two large 8vo. vols. of 1840 pp., with 190 cuts. Clo., $9; lea., $11.

WELLS (J. SOELBERG). A TREATISE ON THE DISEASES OF THE EYE. In one large and handsome octavo volume.

WEST (CHARLES). LECTURES ON THE DISEASES PECULIAR TO WOMEN. Third American from the third English edition. In one octavo volume of 543 pages. Cloth, $3 75; leather, $4 75.

—— ON SOME DISORDERS OF THE NERVOUS SYSTEM IN CHILDHOOD. In one small 12mo. vol. of 127 pages. Cloth, $1.

WHARTON (HENRY R.). MINOR SURGERY AND BANDAGING. In one very handsome 12mo. volume of about 500 pages, with about 400 illustrations, many of which are lithographic. *In press.*

WILLIAMS (CHARLES J. B. and C. T.) PULMONARY CONSUMPTION: ITS NATURE, VARIETIES AND TREATMENT. In one octavo volume of 303 pages. Cloth, $2 50.

WILSON (ERASMUS). A SYSTEM OF HUMAN ANATOMY. A new and revised American from the last English edition. Illustrated with 397 engravings on wood. In one handsome octavo volume of 616 pages. Cloth, $4; leather, $5.

—— THE STUDENT'S BOOK OF CUTANEOUS MEDICINE. In one handsome royal 12mo. vol. Cloth, $3 50.

WINCKEL ON PATHOLOGY AND TREATMENT OF CHILDBED. With additions by the Author. Translated by James R. Chadwick, A.M., M.D. In one handsome 8vo. vol. of 484 pages. Cloth, $4.

WÖHLER'S OUTLINES OF ORGANIC CHEMISTRY. Translated from the 8th German edition, by Ira Remsen, M.D. In one 12mo. volume of 550 pages. Cloth, $3 00.

WOODHEAD (G. SIMS). PRACTICAL PATHOLOGY. A Manual for Students and Practitioners. In one beautiful octavo vol. of 497 pages, with 136 exquisitely colored illus. Cloth, $6.

YEAR-BOOK OF TREATMENT FOR 1891. A Critical Review for Practitioners of Medicine and Surgery. In contributions by 20 well-known medical writers. 12mo., of 485 pages. Cloth, $1 50. In combination with The Medical News and The American Journal of the Medical Sciences, 75 cents. See page 1.

YEAR-BOOK OF TREATMENT FOR 1886, 1887 AND 1890. Similar to above. 12mo., 320–341 pages. Limp cloth, $1 25.

YEO (I. BURNEY) ON FOOD IN HEALTH AND DISEASE. In one 12mo. volume of 590 pages. Cloth, $2. *Just ready.* See *Series of Clinical Manuals*, p. 13.

LANE MEDICAL LIBRARY
STANFORD UNIVERSITY MEDICAL CENTER
STANFORD, CALIFORNIA 94305

Ignorance of Library's rules does not exempt
violators from penalties.



25M-9-70-28042

W1
W554
1891

LANE MEDICAL LIBR
STANFORD UNIVERSI
MEDICAL CENTER
STANFORD, CALIF. 94

M111 Wharton, H.R. 77774
W554 Minor surgery and
1891 bandaging.

NAME	DATE DUE

LANE MEDICAL LIBRARY STANFORD
0940 4573 45 2

Bd. XIII

Verbandlehre

von

A. Hoffa

2. Auflage

7. 6012

Lehmann's med. Handatlanten nebst kurzgefassten Lehrbüchern.

Bisher sind erschienen:

Bd. I. **Atlas und Grundriss der Lehre vom Geburtsakt und der operativen Geburtshilfe.** In 124 farbigen Abbildungen und 16 ganzseit. Tafeln, von Dr. O. Schäffer, Privatdocent an der Universität Heidelberg V. vi[illegible]ach erwe[illegible]te Auflage. Preis [illegible]. geb. M. 8.—

Bd. II. **Anatomischer Atlas der geburtshilflic[illegible] Di[illegible]tik und Therapie.** Mit 160 farbigen Abbildungen und 3[illegible]8 S. [illegible]xt, von Dr. O. Schäffer. II. Aufl. Preis [illegible] geb. M. 12.—

LANE
MEDICAL LIBRARY
LELAND STANFORD JUNIOR UNIVERSITY
1885
Gift
San Francisco County Medical
Library

bildern. Von Prof. Dr. K. B. Lehmann und [illegible]r. K. Neumann in Würzburg. II. erweit. Aufl. Preis eleg. geb. M. 16.–

Bd. XI./XII. **Atlas und Grundriss der patholog. Anatomie.** In 120 farb. Taf. V. Prof. Dr. O. Bollinger. 2 Bde. Preis eleg. geb. je M. 12.—

Bd. XIII. **Atlas u. Grundriss der Verbandlehre** v. Professor Dr. A. Hoffa i. Würzburg. In 143 Taf. II. Aufl. Preis eleg. geb. M. 8.—

Bd. XIV. **Atlas und Grundriss der Kehlkopfkrankheiten.** In 44 farb. Tafeln. Von Dr. L. Grünwald. Preis el[illegible]. geb. M. 8.–

Bd. XV. **Atlas und Grundriss der internen Medicin u. klin. Diagnostik.** In 68 farb. Tafeln. Von Dr. Chr. Jakob. Preis eleg. gb. M. 10.—

Bd. XVI. **Atlas und Grundriss der chirurgischen Operationslehre.** Von Docent Dr. O. Zuckerkandl in [illegible]. Mit 24 farb. Taf. u. 217 Text-Abbildungen. Preis eleg. geb. M. 10.—

Bd. XVII. **Atlas der gerichtlichen Medicin** v. Hofrat Prof. Dr. E. v. Hofmann in Wien. Mit 56 farbigen Tafeln und 193 Textabbildungen. Preis eleg. geb. M. 15.—

Bd. XVIII. [illegible] **Erkrankungen des Auges.** Mit 67 fa[illegible]xtillustr. u. 16 Bog. Text v. Prof. [illegible]reis eleg. geb. Mk. 10.—

Bd. XIX. **Atlas und Grundriss der Unfallheilkunde.** Circa 40 farbige Tafeln, 142 Textabbildungen und 38 Bogen Text. Von Dr. Ed. Golebiewski in Berlin. Preis eleg. geb. Mk. 15.—.

Bd. XX/XXI. **Atlas und Grundriss der speciellen pathologischen Histologie.** Bd. 1 u. 2. Mit 120 farb. Tafeln, zahlr. Textabb., reichem Text. Von Privatdocent Dr. H. Dürck. 2 Bde. Preis eleg. geb. je Mk. 11.—.

Von **Lehmann's medicin. Handatlanten** befinden sich in Vorbereitung:

Bd. XXII. **Atlas und Grundriss der allgemeinen pathologischen Histologie** nebst einem Anhang über patholog. histolog. Technik. Ca. 80 farb. Tafeln, zahlr. Textabbildung., reicher Text. Von Privatdocent Dr. Herm. Dürck. Preis eleg. geb. Mk. 14.—

Bd. XXIII. **Atlas und Grundriss der orthopädischen Chirurgie.** Circa 100 Abbildungen. Von den Docenten Dr. Schulthess und Dr. Lüning in Zürich. Preis eleg. geb. circa Mk. 10.—

Bd. XXIV. **Atlas und Grundriss der gynaekolog. Operationslehre** mit 40 farb. Tafeln u. zahlr. schwarzen Abbild. n. Orig. v. Maler A. Schmitson von Privatdocent Dr. O. Schaeffer in Heidelberg. Preis eleg. geb. ca. Mk. 12.—

Bd. XXV. **Atlas und Grundriss der Ohrenkrankheiten.** Unter Mitwirkung von Prof. Dr. Politzer in Wien herausgegeben von Dr. G. Brühl in Berlin. Ca. 12 Bogen Text mit ca. 120 farbigen Abbildungen. Preis eleg. geb. ca. Mk. 10.—

Bd. XXVI. **Atlas und Grundriss der allgem. Chirurgie.** Unter Mitwirkung v. Geheimrat Prof. Dr. Czerny herausgeg. v. Privatdocent Dr. Marwedel. In ca. 200 Abbildungen nach Originalgemälden v. Maler A. Schmitson. Preis geb. ca. Mk. 12.—

Bd. XXVII. **Atlas und Grundriss der Psychiatrie** von Privatdocent Dr. Wilh. Weygandt in Würzburg. Circa 120 Abbildungen auf Tafeln. Preis eleg. geb. ca. M. 10.—

Bd. XXVIII. **Atlas und Grundriss der normalen Histologie.** Mit 80 farb. Tafeln und zahlr. schwarzen Abbild. von Privatdocent Dr. Johannes Sobotta in Würzburg. Preis eleg. geb. ca. Mk. 12.—

Bd. XXIX. **Atlas und Grundriss der Topographischen Anatomie** von Prof. Dr. Schultze in Würzburg. Circa 200 meist farbigen Abbildungen. Reicher Text. Preis geb. circa Mk. 14.—

Von Lehmann's medicin. Handatlanten sind Uebersetzungen in dänischer, englischer, französischer, holländischer, japanischer, italienischer, madiarischer, rumänischer, russischer, schwedischer, spanischer und tschechischer Sprache erschienen.

Verlag von J. F. LEHMANN in MÜNCHEN.

Lehmann's med. Handatlanten.

Band VIII.

Atlas und Grundriss der traumatischen

Frakturen und Luxationen

mit 200 farbigen und 110 schwarzen Abbildungen nach Originalzeichnungen von Maler Bruno Keilitz

von **Professor Dr. H. Helferich in Greifswald.**

Preis eleg. geb. **Mk. 12.—.**

Vierte vollständig umgearbeitete Auflage.

Auf 68 farbigen Tafeln werden sämtliche Frakturen und Luxationen, die für den Studierenden und Arzt von praktischer Bedeutung sind, in mustergiltiger Weise zur Darstellung gebracht. Jeder Tafel steht ein erklärender Text gegenüber, aus dem alles Nähere über die anat. Verhältnisse, Diagnose und Therapie ersichtlich ist.

Ausserdem enthält der Band ein vollständiges Compendium der Lehre von den traum. Frakturen und Luxationen. Wie bei den Bildern, so ist auch im Texte das Hauptgewicht auf die Schilderung des praktisch Wichtigen gelegt, während Seltenheiten nur ganz kurz behandelt werden.

Zur Vorbereitung für das Examen ist das Buch vorzüglich geeignet. Der Preis ist in Anbetracht der prächtigen, in Farbendruck ausgeführten Bilder ein ganz aussergewöhnlich niedriger.

Professor Dr. Klaussner schreibt: „Die Auswahl der Abbildungen ist eine vortreffliche, ihre Wiedergabe eine ausgezeichnete. Neben dem Bilde, wie es der Lebende nach der Verletzung bietet, finden sich die betreffenden Knochen- oder Gelenkpräparate, sowie eine besonders lehrreiche Darstellung der wichtigsten, jeweils zu berücksichtigenden topographisch-anatomischen Verhältnisse.

Im Texte sind die häufiger vorkommenden, den Arzt besonders interessierenden Knochenbrüche und Verrenkungen in ihrer diagnostischen und auch therapeutischen Beziehung eingehender, seltenere Formen kürzer erörtert. Die Absicht des Verfassers, *„den Studierenden die Einführung in das wichtige Gebiet der Lehre von den Frakturen und Luxationen zu erleichtern und Aerzten in der Praxis ein brauchbarer Ratgeber zu sein"*, ist als vorzüglich gelungen zu bezeichnen.

Der Verleger liess es sich angelegen sein, das Beste zu liefern; das Kolorit der Tafeln ist schön, der Druck übersichtlich, die Ausstattung hübsch, der Preis ein mässiger.

Referent zweifelt nicht, dass der Wunsch des Verfassers, es möge das ich einigen Nutzen stiften, sich vollauf erfüllen wird."

6012 DR. E. STELTZNER,

LEHMANN'S MEDICINISCHE
HANDATLANTEN.
BAND XIII.

Atlas und Grundriss
der
VERBANDLEHRE
für
Studierende und Aerzte
von

Dr. Albert Hoffa,
a. o. Professor an der Universität Würzburg.

Mit 144 Tafeln
nach Originalaquarellen von Maler Joh. Fink.

II. vermehrte und verbesserte Auflage.

München.
Verlag von J. F. Lehmann.
1900.

Das Recht der Uebersetzung bleibt vorbehalten.

YRARBL JHAJ

Kupferätzungen von *Meisenbach*, *Riffarth & Co.*,
Satz und Druck von *Kastner & Lossen*,
Papier von *O. Heck*
Einbände von *L. Beer*,
sämtlich in München.

M113
H67

Meinem verehrten Freunde

Herrn Dr. Oskar Pinner

in Frankfurt a/M.

gewidmet.

Vorrede zur zweiten Auflage.

Die neue Auflage dieser Verbandlehre ist als eine reichlich vermehrte und verbesserte zu bezeichnen. Nach dem Grundsatz: „Prüfet Alles und behaltet das Beste“ bin ich auch diesmal verfahren. Die Anzahl der Tafeln ist um 16 vermehrt worden und dem entsprechend wurde auch der Text vergrössert. Nur das wirklich Brauchbare habe ich aufgenommen und hoffe, dass das Buch auch in seiner neuen Gestalt viele Freunde gewinnen und viel Nutzen stiften möge.

Würzburg, Januar 1900.

Albert Hoffa.

Vorrede zur ersten Auflage.

Die vorliegende Verbandlehre ist auf die Anregung des Herrn Verlegers J. F. Lehmann hin entstanden. Das Bedürfnis nach einem solchen Leitfaden ist ein unter der studierenden Jugend wohl allgemein anerkanntes, und so bin ich der Anregung gern gefolgt. Ich habe die Verbandlehre so dargestellt, wie ich sie seit vielen Jahren in meinen praktischen Cursen vortrage. Nachdem ich viele Hunderte von Schülern in die Verbandtechnik eingeführt habe, habe ich die Fehler wohl kennen gelernt, die der Anfänger beim Verbinden in der Regel macht, und habe darum gleich angegeben, wie sich diese Fehler

106119

vermeiden lassen. Das Buch verfolgt überhaupt einen vorwiegend praktischen Zweck und soll auch dem jungen Arzt ein zuverlässiger Ratgeber sein in allen den einschlägigen Fragen, die ihm in seiner Praxis entgegentreten.

Ich habe deshalb ganz besonderen Wert auf die Herstellung guter Abbildungen gelegt. Die alten schematischen Zeichnungen der Verbände, die man in den früheren Lehrbüchern über die Verbandlehre findet, entsprechen nicht der Wirklichkeit. Ich habe mir Mühe gegeben, die Verbände wirklich so darzustellen, wie sie in der Praxis ausgeführt werden. Ich habe sämtliche Verbände selbst angelegt; dann sind dieselben von meinem Assistenten Herrn Dr. P. Paradies photographiert und nach den wohlgelungenen Photographien von Herrn Maler Fink gemalt worden. So haben wir Tafeln erhalten, welche wohl auch weitgehenden Ansprüchen an Deutlichkeit und Realismus entsprechen dürften.

Ganz besonders möchte ich noch das ausserordentlich grosse Entgegenkommen des Herrn Verlegers hervorheben, der selbst die grössten Kosten zur vollendeten Ausstattung des Buches nicht gescheut hat.

Würzburg, im Oktober 1896.

Albert Hoffa.

Inhalts-Verzeichnis.

Seite

A. Einfache Verbände 1
1) Bindenverbände 1
Verbände an der oberen Extremität 9
Verbände an der unteren Extremität 16
Kopfverbände 22
Verbände für die Augen 25
Verbände am Rumpf 28
2) Tücherverbände 39
Tücherverbände für die obere Extremität 40
Tücherverbände für die untere Extremität 43
Tücherverbände für den Kopf 44
Tücherverbände für die Brust und den Rumpf . . 46
B. Wundverbände 46
Grössere Wundverbände an den einzelnen Körperteilen . 56
C. Lagerungs-Verbände 62
D. Contentivverbände 67
1) Schienenverbände 68
Pappschienen 72
Blechschienen 75
Drahtschienen 81
Plastische Schienen 83
2) Erhärtende Verbände 87
Gipsverband 87
Gipsschienen 99
Wasserglasverband 104
Leimverband 106
Holzleimverband 107
Geleimter Cellulose-Verband 108
Rohrgeflecht-Leimverband 108
Kleisterverband 109
Celluloid-Mullverband 110
E. Zugverbände 111
a) Heftpflastergewichtextensionsverband 111
b) der Zug durch den Contentivverband 118
Gehverband 119
c) der Zug durch Schienen und Apparate 121
F. Druckverbände 123

Verzeichnis der Tafeln.

Taf. 1. Einköpfige Binde.
„ 2. Aufwickeln der Binde mittelst einer Bindenwickelmaschine.
„ 3. Aufwickeln einer Binde mit den Händen.
„ 4. Zweiköpfige Binde.
„ 5. a) Verbandschere, b) einfache Kompresse, c) mehrfache Kompresse, d) Longuette, e) Schleuderbinde oder Funda, f) T-Binde, g) Rollkompresse.
„ 6. Anlegen der Binde zur Ausführung der Kreistour.
„ 7. Bildung des Zipfels bei Anlegung der Kreistour.
„ 8. Dolabra serpens oder repens.
„ 9. Ausführung der Dolabra reversa I. Akt.
„ 10. Ausführung der Dolabra reversa II. Akt.
„ 11. Ausführung der Dolabra reversa III. Akt.
„ 12. Dolabra currens und Dolabra reversa.
„ 13. Beginn einer neuen Binde.
„ 14. Abwickeln der Binde.
„ 15. Spica manus descendens.
Taf. 16. Spica pollicis descendens.
„ 17. Chirotheka dimidia.
„ 18. Beginn der Chirotheka completa.
„ 19. Chirotheka completa.
„ 20. Einhüllung der Fingerkuppe.
„ 21. Involutio digitorum.
„ 22. Testudo cubiti inversa.
„ 23. Testudo cubiti reversa.
„ 24. Spica humeri ascendens.
„ 25. Spica humeri descendens.
„ 26. Einwickelung des ganzen Armes.
„ 27. Stapes.
„ 28. Spica pedis descendens.
„ 29. Testudo calcanei.
„ 30. „ „ mit Abschluss der Ferse.
„ 31. Einwickelung des Fusses von den Zehen an.
„ 32. Einwickelung des ganzen Fusses.
„ 33. Einwickelung des ganzen Fusses.
„ 34. Testudo genu inversa.
„ 35. Testudo genu reversa.
„ 36. Spica coxae ascendens.
„ 37. Spica coxae descendens.
„ 38. Spica coxae duplex.

Taf. 39. Einwickelung des ganzen Beines.
„ 40. Bier'sche Stauung.
„ 41. Capistrum simplex (gesunde Seite).
„ 42. Capistrum simplex (kranke Seite).
„ 43. Capistrum duplex.
„ 44. Mitra Hippocratis.
„ 45. Monoculus.
„ 46. Binoculus.
„ 47. Fascia nodosa.
„ 48. Funda nasi.
„ 49. Funda maxillae.
„ 50. Stella dorsi.
„ 51. Stella pectoris.
„ 52. Stella pectoris et dorsi.
„ 53. Altes, aber unpraktisches Suspensorium mammae.
„ 54. Praktisches Suspensorium mammae.
„ 55. Suspensorium mammae duplex. (Jede Brustdrüse ist für sich eingewickelt.)
„ 56. Suspensorium mammae duplex. (Beide Brustdrüsen sind abwechselnd eingewickelt worden.)
„ 57. I. Tour des Desault'schen Verbandes.
„ 58. II. Tour des Desault'schen Verbandes.
„ 59. III. Tour des Desault'schen Verbandes.
„ 60. Velpeau'scher Verband.
„ 61. Modifikation des Velpeau'schen Verbandes von Dulles.
„ 62. Sayre'scher Heftpflast.-Verband.
„ 63. Schoenborn's Modifikation des Sayre'schen Heftpflasterverbandes.
„ 64. Landerer's Modifikation des Sayre'schen Heftpflasterverbandes.
Taf. 65. Büngner's Verband zur Behandlung d. Schlüsselbeinbrüche.
„ 66. Büngner's Verband zur Behandlung d. Schlüsselbeinbrüche.
„ 67. Tuchverband für das Handgelenk.
„ 68. Tuchverband für die ganze Hand. (Anfang des Verbandes.)
„ 69. Tuchverband für die ganze Hand. (Fertiger Verband.)
„ 70. Tuchverband für das Ellbogengelenk.
„ 71. Tuchverband für das Schultergelenk.
„ 72. Mitella triangularis zur Stütze für den Oberarm.
„ 73. Mitella triangularis zur Hebung des Oberarmes.
„ 74. Moore'scher Verband.
„ 75. Mitella parva und Tuchverband für die Wange und das Ohr.
„ 76. Mitella quadrangularis,
„ 77. Tuchverband für das Fussgelenk.
„ 78. Tuchverband für den ganzen Fuss.
„ 79. Tuchverband für das Knie.
„ 80. Tuchverband für die Hüfte.
„ 81. Capitium triangulare.
„ 82. Capitium quadrangulare.
„ 83. Tuchverband für die Stirn und den Hals.
„ 84. Tuchverband für das Auge und die weibliche Brust.
„ 85. Tuchverband für den Unterleib.
„ 86. Schamtuch.
„ 87. Kleinere Wundverbände. (Neff'scher

Fingerling aus Trikotgewebe.)
Taf. 88. Wundverband am Kopf.
„ 89. Wundverband am Hals.
„ 90. Wundverband an der Brust.
„ 91. Lagerung des Patienten zur Anlegung eines Wundverbandes am Unterleib. Stützbänkchen von v. Volkmann (a) & Czerny (b).
„ 92. Lagerung des Patienten auf der Ollier'schen Stütze.
„ 93. Stützen des Beckens durch einen Assistenten von unten her.
„ 94. Hase-Beck'scher Krankenhebeapparat.
„ 95. Lagerung des Kranken in einem Stehbett.
„ 96. Wundverband am Damm.
„ 97. a) Desault'sches Kissen.
b) & c) Stromeyer'sches Kissen.
d) v. Dumreicher'sches Kissen.
„ 98. Middeldorpf'scher Triangel.
„ 99. a) Petit-Heister'sche Lade.
b) Scheuer'sche Lade.
„ 100. a) Einfach geneigte schiefe Ebene.
b) Doppelt geneigte schiefe Ebene.
c) Fialla'sche Stäbchenlade.
„ 101, Reifenbahre.
„ 102. a) Mayor'sche Drahtrinne.
b) Roser'sche Drahtrinne.
c) Bonnet'sche Drahthose.
Taf. 103. a) & b) Stromeyer'sche Handbretter. c) Nélaton'sche Pistolenschiene. d) Roser'sche Dorsalschiene für die Fractura radii loco classico. e) Schede'sche Schiene für die Fractura radii loco classico. f) Carr'sche Schiene für die Fractura radii loco classico. g) Cooper'sche Schiene f. die Fractura radii loco classico. h) Stromeyer'sche Armschiene.
„ 104. a) v. Volkmann'sche Suspensionsschiene. b) & c) Dupuytren'scher Verband bei Malleolarfraktur. d) v. Volkmann'sche Knieschiene.
„ 105. a) Watsons'sche Schiene. b) & c) v. Volkmann's Dorsalschiene für den Unterschenkel. d) König's Abduktionsschiene für das Hüftgelenk.
„ 106. a) Middeldorpf's Pressschienenverband. b) Gooch'sche Spaltschienen. c) Schnyder's Tuchschienen. d) v. Esmarch's schneidbarer Schienenstoff.
„ 107. a) v. Volkmann's Supinationsschiene.
b) & c) v. Esmarch's Resektionsschiene für die Hand.
d) Sharp'sche Beinschiene.
„ 108. Pappwatte-Verband Linhart's.

a) Verband für den gebrochenen Oberarm. b) Verband für den gebrochenen Vorderarm.
Taf. 109. Pappverband für den gebrochenen Oberarm.
„ 110. a) v. Volkmann'sche ⊤ -Schiene. b) Bruns'sche Schiene. c) Salomon'sche Schiene. d) Schön'sche Schiene für den Arm. e) Schön'sche Schiene für den Unterschenkel.
„ 111. a) Raoult-Deslongchamps'sche Schiene. b) & c) Kupferblechschienen nach Lee und Wilson. d) & e) Englische Eisenblechschienen nach Koch. f) & g) Port'scheBlechstreifenschienen.
„ 112. Steudel's Aluminiumschienen.
„ 113. a) Aluminiumblechschiene nach Hausmann. b) Lagerung derselben. c) Lagerung in Dorsalflexion. d) Lagerung in Ulnarflexion.
„ 114. a, b, c, d) Cramer'sche Schienen. e) Anwendung der Cramer'schen Schiene beim Oberarmbruch nach Helferich. f) Smith's vordere Drahtschiene.
„ 115. v. Esmarch's Drahtsiebstoff. c) Port's Drahtrollbinden. b) Heusner's Spiraldrahtschiene.
„ 116) Plastische Pappschienen nach Koch. (a, b, c, d). Filzschienen nach P. Bruns-Anders (e). f) Pappschiene für den gebrochenen Oberarm.
Taf. 117. Gipsbindenverband.
„ 118. a) Gipsbindenwickelmaschine nach Beely. b) Tricotschlauchbinde als Unterpolsterungsmaterial und zugleich als Handhabe. c) v. Esmarch'sches Gipsmesser. d) Gipsschere nach Seutin. e) Gipsschere nach Sczymanowski. f) Gipsschere nach Bruns.
„ 119. a) Zange zum Oeffnen der Gipsverbände nach J. Wolff b) Gipsschere nach Stille. c) Gipsschere nach Empfen) zeder. d) Kreis-Gipssäge. e) Blatt-Gipssäge. f) Aufsägen eines Gipsverbandes.
„ 120. Technik d. Anwendung der ersten Stille'schen Schere.
Taf. 121. a) Fenster im Gipsverband. b) Völker's Holzspan-Gipsverband c) Unterbrochener Gipsverband. d) Gipsverband mit Gelenkschienen.
„ 122. a) Abnehmbare Gipshülle für das Bein. b) Abnehmbare Gipshülle für die Hüfte. c) Beckenstütze nach v. Esmarch. d) Beckenstütze nach v. Bardeleben. e) Lagern des Patienten auf der Beckenstütze.

Taf. 123. a) Lagerung des Patienten auf dem Bruns'schen Apparat. b) Lagerung des Patienten auf den Dittel'schen Stangen. c) Albers-'sche Kragenschiene. d) Breiger'sche Gipswatteschiene.
„ 124. Extensionstisch von Schede.
„ 124a. Schede's Tisch zur Anlegung von Gipsverbänden am Rumpf.
„ 125. Verband und Extensionstisch v. Heuser.
„ 126. Beely'sche Gipshanfschienen. a, b) für die obere Extremität; c, d) für die untere Extremität (mit Suspensionsvorrichtung).
„ 127. Artikuliert-mobile Wasserglasverbände nach Kappeler und Hafter.
„ 128. a) Anlegung eines Leimverbandes. b, c, d) Verschiedene Rohrgeflechte.
„ 129. Heftpflastergewichtsextensionsverband.
„ 130. a) Anwendung der Kontraextension (mittelst einer Quehle und Hochstellung des Bettes am Fussende.) b) v. Volkmann'sches Schleifbrett.
„ 131. Rollenträger von Eberth
„ 132. a) Verwendung mehrerer Heftpflasterzüge nach Bardenheuer. b) Schede's Methode der vertikalen Extension bei Kindern.
Taf. 133. a) Schede's Methode zur Streckung von Kniegelenksverkrümmungen unter Zuhülfenahme mehrfacher Heftpflasterzüge. b) v. Volkmann's Stecknadelverband. c) Zug vermittelst des Contentivverbandes.
Taf. 134. Permanenter Extensionsverband nach Clarc.
„ 135. Permanenter Extensionsverband nach Clarc.
„ 136. Extensionsmethode nach Hennequin.
„ 137. Gehverband nach F. Krause.
„ 138. Gehverband nach F. Krause.
„ 139. a) Taylor'sche Extensionsschiene. b) Thomas'sche Schiene. c) Heusner'sche Schiene.
„ 140. a) Liermann'sche Schiene. b) Hessing'scher Schienenhülsenapparat.
„ 141. P. Bruns'sche Extensionsschienen und Lagerungsapparat.
„ 142. a) Fricke'scher Heftpflasterverband bei Orchitis und Periorchitis. b) Baynton'sche Heftpflastereinwickelung bei Ulcus cruris.
„ 143. Gibney'scher Heftpflasterverband bei Distorsio pedis.

ATLAS UND GRUNDRISS

DER

VERBANDLEHRE.

A) Einfache Verbände.

Man teilt die einfachen Verbände ein in Bindenverbände und Tücherverbände.

1) Bindenverbände.

Während man in früheren Jahren die Bindenverbände mit Leinenbinden anlegte, benutzt man heute als Stoff zu den Binden fast ausschliesslich die Baumwolle, die entweder locker gewebt als einfache Gaze- oder Mullbinde, oder fester gewebt, als Calicot- oder Cambricbinde in Verwendung kommt. Wenn die einfache Gaze- oder Mullbinde noch Stärke enthält oder mit Stärke imprägniert ist, so heisst sie appretierte oder gestärkte Gazebinde oder blaue Binde oder auch wohl Organtinbinde. Diese gestärkten Binden werden stets nass angelegt. Man taucht sie in Wasser, drückt sie gehörig aus und zieht sie beim Umwickeln recht fest an, da die Binde beim Trocknen wieder lockerer wird. Umgekehrt zieht sich die Leinenbinde trocken angelegt und dann befeuchtet, zusammen. Relativ selten kommt als Bindenstoff die reine Wolle in Verwendung. Man benutzt sie in der Form der Flanellbinde nur zu gewissen Zwecken.

Die Binden selbst verwendet man als einköpfige oder zweiköpfige Binden. Die einköpfige Binde (Tafel 1) entsteht dadurch, dass man sich die Bindenstoffe in verschieden breite Streifen schneidet und diese einzelnen Streifen nun aufwickelt. Dieses Aufwickeln kann entweder mittelst einer Bindenwickelmaschine oder mittelst der Hände geschehen.

ckelmaschinen sind in verschiedener Form angegeben worden. Sie haben alle das gemeinsam, dass man das Bindenende an einer Kurbel befes und durch Drehen dieser Kurbel den Binden- auf die Kurbelachse aufrollt. Um eine recht ige Festigkeit der Bindenrolle zu erzielen, muss vanrend man mit der einen Hand die Kurbel dreht, r andern Hand das Ende der Binde recht stramm (Tafel 2.)

nan beim gewöhnlichen Anlegen von Verbänden wickelmaschinen nicht gebrauchen kann, muss man rin üben, die Binde auch mit den Händen zuwickeln. Es geschieht dies in folgender el 3.) Man nimmt das eine Ende des Bindenlegt dasselbe mehrfach platt auf einander und o, indem man die Platte zusammendreht, Knäuel, der den Beginn der Binde dar- en Knäuel vergrössert man nun zunächst dadurch, dass man ihn zwischen Zeigefinger und Daumen beider Hände an beiden Seiten fasst und ihn zwischen diesen Fingern so um seine Achse dreht, dass immer neue Partieen des Bindenstreifens in die Rolle mit einbezogen werden. Hat die Rolle nun etwa die Dicke eines Daumens erreicht, so fasst man sie von beiden Seiten her zwischen Daumen und Mittelfinger der linken Hand. Der linke Zeigefinger kommt oben auf die Rolle selbst zu liegen. Unter den herunterhängenden Bindenstreifen legt man nun parallel mit der Rolle die rechte Hand, so dass der Radialrand des Zeigefingers direkt nach oben zu stehen kommt, während der Mittelfinger von unten her die Rolle umfasst. Indem nun die rechte Hand den Bindenstreifen zwischen Daumen und Zeigefinger fasst und ihn fest anzieht, dreht sie die Bindenrolle vermittelst des Daumens und des Mittelfingers im entgegengesetzten Sinne des Uhrzeigers so oft herum, bis der ganze Streifen aufgerollt ist. Damit die Rolle recht fest und gleichmässig aufgewickelt wird, ist es wieder notwendig, den noch nicht aufgewickelten Bindenstreifen recht fest zwischen Daumen und Zeigefinger

Tafel 1.

Einköpfige Binde.

Die linke Hand hält die Binde, wie sie beim Aufrollen gehalten werden soll.

Tafel 2.

Aufwickeln der Binde mittelst einer Bindenwickelmaschine.

Tafel 3.

Aufwickeln einer Binde mit den Händen.



Tafel 4.

Zweiköpfige Binde.

anzuspannen. Die linke Hand unterstützt die Binde beim Aufrollen. Wir teilen ihr nach dem Gesagten nur eine Nebenaufgabe zu, weil wir gefunden haben, dass das Aufrollen vermittelst der rechten Hand im allgemeinen leichter ist, als wenn der linken Hand die Hauptaufgabe des Drehens zufällt.

Ist die Binde aufgewickelt, so besteht sie aus dem Bindenkopf und dem Bindenende. Da sie nur einen Kopf hat, so nennt man sie eine einköpfige Binde.

Eine zweiköpfige Binde (Tafel 4) erhält man dann, wenn man den Bindenstreifen von beiden Enden her in der oben geschilderten Weise gleichmässig aufwickelt, so dass schliesslich nur die Mitte des Bindenstreifens übrig bleibt. Bei dem Aufwickeln der doppelköpfigen Binde muss man darauf achten, dass die beiden Bindenköpfe nach einer und derselben Seite sehen. In früheren Zeiten verwendete man wohl auch noch vielköpfige Binden; dieselben sind heut zu Tage ganz ausser Gebrauch gekommen. Dagegen wendet man noch hie und da die sog. T-Binde an (Tafel 5f), die man dadurch erhält, dass man senkrecht auf die Mitte einer doppelköpfigen Binde eine einköpfige Binde annäht.

Spaltet man eine Binde von beiden Seiten her so, dass nur ein kleines Mittelstück ungespalten bleibt, so erhält man eine sog. Schleuderbinde oder Funda. (Tafel 5e.)

Schneidet man irgend einen Verbandstoff, in der Regel die Verbandgaze, in beliebig grosse Stücke, so nennt man diese Verbandstücke Compressen. Je nach der Zahl der Verbandlagen und der Form des Umfanges unterscheidet man einfache (Tafel 5b) und mehrfache (Tafel 5c), viereckige oder Roll-Compressen. (Tafel 5g.)

Längliche, viereckige Compressen nennt man auch wohl Longuetten. (Tafel 5d.)

Als Verbandstoffe braucht man ausser den Binden in der Regel noch die Verbandgaze, d. i. Baumwollengewebe in Mullform, und die Verbandwatte, die entweder als entfettete Watte bei Wundverbänden

oder als geleimte, rohe Watte zur Unterpolsterung der Verbände in Verwendung kommt.

Das Verbinden irgend eines Körperteiles besteht nun darin, dass man die Binde in bestimmter Weise um diesen Körperteil herumführt.

Als Beispiel dafür, wie dieses Anlegen der Binde zu geschehen hat, wählen wir zunächst **das Verbinden des Vorderarmes**, da sich uns hierbei am leichtesten die Gelegenheit bietet, die Regeln zu besprechen, die man beim Anwickeln der Binde befolgen muss, um einen gutsitzenden Verband zu erhalten.

Um den Vorderarm regelrecht zu verbinden, beginnt man mit einer Kreistour oberhalb des Handgelenkes.

Diese Kreistour, fascia circularis (Tafel 6 & 7), stellt den Beginn der meisten Verbände dar. Man darf sie nicht etwa in der Weise anlegen wollen, dass man die Binde einfach kreisförmig um das untere Ende des Vorderarmes herumführt; denn dann würde beim Anziehen des Bindenkopfes das schon abgewickelte Ende der Binde vom Vorderarm einfach abgleiten. Man muss vielmehr der Binde zunächst einen Halt an dem betreffenden Körperteile geben, und dies geschieht in folgender Weise. Der Arzt stellt sich so, dass er den zu verbindenden Vorderarm zu seiner rechten Seite hat. Nun nimmt er das Bindenende, rollt es etwa handbreit vom Bindenkopf ab, legt es am unteren Ende des Vorderarmes in leicht schräger Richtung über dem Handgelenk an und fixiert es in dieser Lage mit seiner linken Hand. Der Bindenkopf sieht dabei nach oben. (Tafel 6.) Es ist ein Fehler, das Bindenende so anzulegen, dass der Bindenkopf nach unten gerichtet ist.

Die Antwort auf die Frage, auf welcher Seite man beginnen soll, resp. nach welcher Seite das Bindenende zunächst gerichtet sein soll, ergibt sich leicht aus der Regel, dass der Rechtshändige von sich aus von links nach rechts wickeln soll, d. h. von seiner eigenen linken nach seiner rechten Seite hin. Fängt er auf der eigenen rechten Seite an und wickelt nach links hinüber, so wickelt er entgegen seiner eigenen rechten Hand und begibt

Tafel 5.



a) Verbandschere, b) einfache Kompresse, c) mehrfache Ko
d) Longuette, e) Schleuderbinde od. Funda, f) T-Binde, g) Roll

Tafel 6.

Anlegen der Binde zur Ausführung der Kreistour.

Tafel 8.

Dolabra serpens oder repens.

Tafel 8.

Dolabra serpens oder repens.



sich somit des Vorteils, den beim Rechtshändigen die grössere Geschicklichkeit der rechten Hand gegenüber der linken mit sich bringt. Man legt also, wenn man den Patienten vor sich stehend denkt und annimmt, dass man den rechten Vorderarm verbinden will, die Binde in der schrägen Richtung derart an, dass das Bindenende an den ulnaren Rand des Vorderarmrückens zu liegen kommt, etwa handbreit über dem proc. styloid. ulnae, während der Bindenkopf mit seinem unteren Ende in der Höhe des proc. styloid. radii steht. Nun rollt man die Binde einmal kreisförmig um das untere Ende des Vorderarmes herum. Dadurch erhält man an dem Bindenende einen Zipfel, den man nun um die Kreistour nach unten herunterschlägt und mit einer zweiten Kreistour befestigt. (Tafel 7.) Jetzt kann man an dem Bindenkopf so viel ziehen als man will, er wird nie mehr abrutschen. So hat man die fascia circularis vollendet.

Verbindet man nun den Vorderarm in der Weise weiter, dass die einzelnen Bindentouren sich gegenseitig zum Teil decken, so erhält man die sogenannte Dolabra currens (siehe Tafel 12). Lässt man dagegen die Touren den Vorderarm spiralförmig umgehen, so entsteht die Dolabra serpens oder repens. (Tafel 8.) Letztere Art kommt im ganzen selten zur Verwendung und zwar nur dann, wenn man einen Körperteil eingewickelt hat und nun mit derselben Binde rasch an einen entfernter liegenden Körperteil gelangen will.

Um so häufiger benutzt man dagegen die Dolabra currens. Man kann dieselbe so ausführen, dass die folgende Tour die vorhergehende entweder zu zweidrittel oder zur Hälfte oder noch weniger deckt. Den Grad der Deckung wählt man nach dem Zweck, den man mit der Binde verfolgt. Will man etwa nur ein Verbandstück an dem Körperteil befestigen, so braucht die gegenseitige Deckung der Binde nur eine geringe zu sein. Verfolgt man dagegen den therapeutischen Zweck einer Compression der unterliegenden Teile, so muss man die einzelnen Bindentouren sich in stärkerem Grad decken lassen. Je mehr die folgende

Tour die vorhergehende zudeckt, desto stärker gestaltet sich der Grad der Compression der Binde auf die Unterlage.

Jede folgende Bindentour soll im allgemeinen der vorhergehenden parallel laufen, und es sollen bei einem regelrechten Verbande die Abstände zwischen den einzelnen Bindentouren möglichst gleiche sein. (siehe Taf. 12.)

Soll die Binde ihren Zweck erfüllen, so darf sie natürlich nicht zu lose am Arm angelegt werden, denn sonst verschiebt sie sich schon nach kurzer Zeit aus ihrer Lage heraus. Sie darf aber auch nicht zu fest angelegt werden, denn sonst kann sie leicht eine venöse Stauung in den peripher gelegenen Teilen, hier also in den Fingern hervorrufen. Da man den Lernenden auf die Gefahr einer zu festen Umschnürung des betreffenden Körperteiles aufmerksam machen muss, so macht man meist die Beobachtung, dass der Anfänger die Binde zu locker wickelt. Durch Uebung lernt man aber bald, die richtige Spannung der Binde herzustellen. Man erreicht diesen richtigen Spannungsgrad der Binde am besten in der Weise, dass man die Binde während des Momentes des Umwickelns selbst leicht und elastisch anzieht. Es ist das besser als die Art und Weise, die der Anfänger meistens übt, dass er nämlich die Bindentour vollständig anwickelt und dann erst am Bindenkopf zieht.

Wenn man nun vom Handgelenk an den Vorderarm nach dem Ellenbogengelenk hin in der geschilderten Weise einwickelt, so dass die einzelnen Touren sich etwa zur Hälfte oder zu Zweidrittel decken, so wird man schon nach wenigen Touren merken, dass die einzelnen Touren sich dem Gliede nicht mehr vollständig anlegen, dass sie vielmehr an ihren unteren Rändern von demselben abstehen. Je stärker der Vorderarm ist, um so eher wird sich dieses Abstehen bemerkbar machen.

Man trifft diese Erscheinung, d. h. das mangelhafte Anliegen der unteren Ränder der Binde an allen denjenigen Körperteilen, welche von der Peripherie nach dem Centrum hin kegelförmig an Dicke zunehmen, so namentlich

Tafel 9.

Ausführung der Dolabra reversa.
(I. Akt: Haltung der Binde vor dem Umschlag.)

Tafel 10.

Ausführung der Dolabra reversa.
(II. Akt: Bildung des Umschlages.)

Tafel II.

Ausführung der Dolabra reversa (III. Akt: Umschlag vollendet).

Tafel 12.



Dolabra currens und Dolabra reversa.



am Unterarm, am Unterschenkel, in geringerem Grade am Oberarm und Oberschenkel. Um auch an diesen Teilen überall ein glattes Anliegen der Binde zu erreichen, muss man sich eines Kunstgriffes bedienen: der Dolabra reversa oder des Renversé:

Die Dolabra reversa (Tafel 9, 10 u. 11) ist nichts anderes als ein Umschlagen der Binde, so dass dieselbe etwa um die Hälfte verschmälert wird. Um einen solchen Umschlag kunstgerecht zu machen, führt man, sobald man merkt, dass die betreffende Tour sich nicht glatt anlegt, die Binde leicht schräg in die Höhe, zieht sie leicht an, fixiert den unteren Rand in der Mitte des Vorderarmrückens mit dem Daumen der linken Hand (Tafel 9) und dreht nun die den Bindenkopf haltende rechte Hand, die in Mittelstellung zwischen Pronation und Supination steht, in Pronationsstellung um, indem man sie gleichzeitig dem Vorderarm nähert und etwas nach dem Handgelenk des Patienten hin führt. (Taf. 10.) Durch diese Manipulation erreicht man, dass sich die Binde gerade in ihrer Hälfte umschlägt. (Taf. 11.) Den umgeschlagenen Teil der Binde führt man nun kreisförmig um den Vorderarm herum, so dass der untere Rand der neuen Tour dem unteren Rand der vorhergehenden Tour parallel läuft. Beim Umführen der Binde um den Vorderarm muss man dieselbe wieder leicht und elastisch anspannen. Hat man den Umschlag richtig gemacht, so findet gerade in der Mitte des Vorderarms eine Kreuzung der umgeschlagenen Bindentour mit der vorhergehenden statt. Man spannt nun die Binde wieder an, fixiert sich die ebengenannte Kreuzungsstelle mit dem Daumen der linken Hand, schlägt dann die Binde mit der rechten Hand in der obengeschilderten Weise um, führt die Tour wieder parallel der vorhergehenden um den Vorderarm herum und fährt so fort bis in die Gegend des Ellenbogengelenks, d. h. bis man sieht, dass sich die Bindentouren dem wieder gleichmässig rund gewordenen Vorderarm auch ohne Umschlag glatt anschmiegen. Bei einem gut angelegten Renversé sollen sich sämtliche Kreuzungsstellen der Binde in einer geraden Linie in der Mitte des Vorderarmes b
(Taf. 12.)

Dem Anfänger will das Anlegen einer guten Dolabra reversa in der Regel nicht gelingen. Durch Uebung kommt er aber bald zum Ziel. Die Fehler, die er macht und die vermieden werden müssen, sind folgende: 1) wickelt er in der Regel den Bindenkopf zu weit ab, so dass er den langen abgewickelten Streifen nicht genügend beherrschen kann. Der Bindenkopf darf nur soweit abgewickelt werden, als etwa die Breite des betreffenden Gliedes beträgt; dann lässt er sich am bequemsten dirigieren; 2) spannt der Anfänger die Binde beim Umschlagen in der Regel zu stark an. Sobald er dies thut, legt sich die Binde in Falten und schlägt sich ungleichmässig um. Jede Faltenbildung muss aber auf das ängstlichste vermieden werden, weil an den betreffenden Stellen des Gliedes sonst leicht Druckstellen entstehen und die Kompression eine ungleichmässige wird. Die Faltenbildung wird am besten vermieden, wenn man beim Umschlagen der Binde dieselbe dem einzuwickelnden Gliede nähert; 3) führt der Anfänger die umgeschlagene Binde meist nicht parallel der vorhergehenden Tour um den Arm herum, sondern so, dass die Binde eine divergierende Richtung nimmt. Dadurch aber kommt die Kreuzung der Binde in unregelmässiger Weise zu stande.

Hat man eine Binde ganz abgewickelt und bedarf einer neuen zur weiteren Einwickelung des Gliedes, so ist ein kleiner Kunstgriff angezeigt. Um nämlich nachher beim Abwickeln der Binden nicht erst lange nach dem Ende der erst angelegten Binde suchen zu müssen, legt man die neue Binde stets so an, dass man den Anfang der neuen Binde ein kleines Stück weit unter das Ende der ersten Binde herunterführt; dann macht man eine Kreistour und erreicht so, dass einem nachher beim Abwickeln das Ende der ersten Binde von selbst in die Hand fällt. (Tafel 13.)

Soll der Verband vom Körper entfernt werden, so kann man ihn entweder abwickeln oder abschneiden.

Das Abwickeln der Binden geschieht nicht in der Weise, dass man das Ende der Binde nimmt und nun die Binde wieder in umgekehrter Richtung, wie man sie

Tafel 13.

Beginn einer neuen Binde.

Tafel 14.

Abwickeln der Binde.

angewickelt hat, wieder aufwickelt. Das würde zu viel Zeit beanspruchen und würde dem Patienten lästig sein. Man nimmt vielmehr die Binde ungeordnet ab, indem man, — möglichst ohne das verbundene Glied zu berühren oder zu bewegen — die abgewickelten Bindetouren von einer Hand in die andere gibt. (Tafel 14.)

Zum Aufschneiden der Verbände bedient man sich zweckmässig der Seutin'schen Verbandschere (Taf. 5a.) Beim Aufschneiden des Verbandes hält man sich stets über den dickeren Weichteilen und vermeidet, den Verband dort aufzuschneiden, wo er dicht über prominierenden Knochenvorsprüngen liegt. Ebenso schneidet man nicht direkt über einer Wunde auf, und schliesslich vermeidet man womöglich das Schneiden im Winkel, also z. B. auf der vorderen Seite des Fussgelenkes.

Verbände an der oberen Extremität.

Bleiben wir bei der oberen Extremität, so ist das nächste, was der Anfänger zweckmässig übt, die

Spica oder Kornähre.

Als Spica oder Kornähre bezeichnet man die Einwicklung eines Gelenkes, derart, dass man peripher oder central vom Gelenk mit einer Kreistour beginnt und das Gelenk dann mit drei aufeinander folgenden, sich etwa zu $^2/_3$ deckenden und parallel laufenden Achtertouren umgibt, um schliesslich mit einer Kreistour zu schliessen, welche die erste Kreistour deckt.

Fängt man peripher von dem Gelenk an und wickelt gegen das Centrum hin, so müssen die einzelnen Bindentouren ansteigen — es entsteht eine Spica ascendens; fängt man dagegen central vom Gelenk an und wickelt gegen die Peripherie hin, so steigen die Bindentouren ab — es entsteht eine Spica descendens.

Zwischen der Kreistour und der ersten Achtertour soll bei einer regelrecht angelegten Spica ein kleiner dreieckiger Raum frei bleiben, das sogenannte Geranium.

Unter Befolgung dieser Regeln wird die

Spica manus descendens (Tafel 15)

n der rechten Hand so angelegt: Kreistour oberhalb des ndgelenkes, Achtertour um das Handgelenk, so zwar, s die Binde von der Kreistour aus schräg über den ndrücken, zum Zeigefinger hin, um den Radialrand Zeigefingers und den Ulnarrand des kleinen Fingers rum und schräg über den Handrücken hin wieder zum lenk zurückgeführt wird. Dieser ersten Achter- gen dann noch zwei weitere gleiche Achtertouren, che der ersten parallel laufen und sich so decken, ass stets der nach dem Handgelenk hinsehende Rand der vorhergehenden Tour freigelassen wird. Hat man die drei Achtertouren gewickelt, so wird der Verband mit einer „Schluss"kreistour um das Handgelenk geschlossen, welche die erste Kreistour deckt.

Die so entstandene Spica manus ist eine descendierende; sie dient zur Feststellung des Handgelenkes.

Die **ascendierende Spica manus** würde mit einer Kreistour um die Hand herum beginnen, und die Achtertouren würden sich so decken, dass immer der nach den Fingern hinsehende Rand der vorhergehenden Tour frei bliebe. Diese Spica manus ascendens wird aber in der Praxis sehr selten verwendet.

Spica pollicis descendens. (Tafel 16.)

Kreistour um das Handgelenk, drei descendierende Achtertouren um das Metacarpophalangealgelenk des abduziert stehenden Daumens, Schlusszirkeltour um das Handgelenk herum.

Die Spica, die man an den anderen vier Fingern macht, bezeichnet man in der Regel als

Chirotheca dimidia. (Tafel 17.)

Der betreffende Verband, nehmen wir nun irgend einen Finger, z. B. den Zeigefinger, wird in der Weise elegt, dass mit einer Kreistour um das Handgelenk en wird, dass dann drei descendierende Achter- m das Metacarpophalangealgelenk des Zeige- endlich wieder eine Schlusstour um das

Tafel 15.

Spica manus descendens.

Tafel 16.

Spica pollicis descendens.

Tafel 17.

Chirotheka dimidia (χεῖρ die Hand und θήκη die Binde).

Tafel 18.

Beginn der Chirotheka completa.

Tafel 19.

Chirotheka completa.

Tafel 20.

Einhüllung der Fingerkuppe.

Handgelenk folgen. Die Achtertouren sollen nur die Grundphalanx des Fingers decken und das erste Interphalangealgelenk nicht übersteigen.

Will man den ganzen Finger einwickeln, so geschieht dies mittelst der

Chirotheca completa. (Tafel 18 u. 19.)

Die Einwickelung des ganzen Fingers ist in Anbetracht der so häufigen Panaritien und Fingerverletzungen in der Praxis sehr oft auszuführen, und ist der Verband daher gehörig einzuüben.

Nehmen wir wieder den rechten Zeigefinger als Beispiel. so beginnen wir mit einer Kreistour um das Handgelenk, führen die Binde, die nicht zu breit sein darf, von der Kreistour aus schräg über den Handrücken nach dem Zeigefinger hin, führen sie um den Radialrand desselben herum und gehen nun sofort mit Spiralwindungen, also mit einer Dolabra serpens bis an die Spitze des Fingers heran. (Taf. 18.) Von der Spitze des Fingers an wickelt man nun mit gleichmässig sich deckenden Touren einer Dolabra currens gegen die Basis des Fingers hin, indem man, wenn nötig, einen oder den anderen Umschlag hinzufügt. Ist man mit den Kreistouren bis unmittelbar an die Basis des Fingers gelangt, so führt man die Binde in Form einer halben Achtertour nach dem Radialrand des Vorderarmes hin und schliesst mit der Kreistour über der ersten Kreistour um das Handgelenk. (Tafel 19.)

Die Touren um den Finger müssen so fest angelegt sein, dass man sie von demselben nicht durch einfachen Zug entfernen kann.

Der geschilderte Verband lässt die Kuppe des Fingers frei. Will man auch die Fingerkuppe mit zudecken (Taf. 20), was beim Anlegen antiseptischer Verbände nötig ist, so legt man den Verband genau in derselben Weise an, wie er eben geschildert wurde, schliesst dann aber nicht mit der Kreistour, sondern führt die Binde in ihrer ganzen Breite erst noch einmal schräg über den Handrücken und den ganzen Finger herüber,

kt mit ihr die Fingerkuppe zu, führt sie nun an volaren Seite des Fingers und der Hand nach n Processus styloideus radii hin, wobei man ja nicht ie Binde anziehen darf, weil sonst ein unleidlicher Druck auf die Fingerspitze entsteht, führt dann die nde wieder über den Handrücken und in Spiral-ndungen bis zur Spitze des Fingers und wickelt nun hliesslich wieder wie vorher von der Spitze des Fin-ers nach seiner Basis hin, macht an letzterer ange-mmen noch eine halbe Achtertour in gleicher Weise ie vorhin und schliesst mit einer Zirkeltour um das Handgelenk.

Involutio digitorum. (Tafel 21.)

Die Einwicklung aller fünf Finger, die Involutio digitorum geschieht in der Weise, dass man mit einer Kreistour um das Handgelenk herum beginnt und dass man dann, vom Daumen oder kleinen Finger angefangen, sämtliche Finger der Reihe nach, nach Art der eben beschriebenen Chirotheca completa, einwickelt. Man gelangt von dem einen Finger zum andern so, dass man nach Ausführung der halben Achtertour, welche die Chirotheca completa beendigt, die Binde schräg über den Handrücken zu dem nächsten Finger führt, um dann mit Spiralwindungen an die Spitze desselben zu gelangen und von dieser dann wieder bis zur Basis des Fingers zurückzuwickeln. Den Schluss bildet eine Zirkeltour um das Handgelenk herum.

Die Einwicklung des Ellbogengelenkes

geschieht mittelst eines Verbandes, den man als Testudo cubiti (testudo die Schildkröte) bezeichnet.

Man unterscheidet eine testudo cubiti inversa und reversa.

Testudo cubiti inversa. (Tafel 22.)

Man legt den Verband am besten bei leicht ge-am Ellbogengelenk an. Den Beginn der Binde bildet tour, die man etwa handbreit unterhalb des

Tafel 21.

Involutio digitorum.



Tafel 22.

Testudo cubiti inversa.

Tafel 20.

Einhüllung der Fingerkuppe.

Ellbogengelenkes anlegt. Von dieser Kreistour aus führt man nun die Binde nach der Ellenbeuge hin, über dieselbe hinweg, um den unteren Teil des Oberarmes herum, wieder über die Ellenbeuge hin und zur Kreistour zurück. So hat man das Ellbogengelenk mittelst einer Achtertour umgangen. Dieser ersten Achtertour sollen nun noch zwei weitere Achtertouren in der Weise folgen, dass die einzelnen Touren sich etwa zur Hälfte decken, so zwar, dass immer der Rand der vorhergehenden Tour zugedeckt wird, der gegen das Ellbogengelenk hinschaut. Die Kreuzung der einzelnen Bindentouren findet stets in der Ellenbeuge statt. Wickelt man die Achtertouren in der angegebenen Weise an, so hat man mit der dritten Achtertour den ganzen von der ersten Achtertour freigelassenen Raum an der Hinterseite des Gelenkes zugedeckt. Hat man die dritte Achtertour vollendet, so schliesst man den Verband mit einer Kreistour, die gerade über das Gelenk hinweg verläuft.

Man hat so eine Kreistour mitten um das Gelenk erhalten, von der aus je drei Touren nach dem Oberarm hin aufsteigen, nach dem Unterarm hin absteigen.

Der Anfänger macht bei der Anlegung der testudo cubiti inversa gewöhnlich einige Fehler. Der erste Fehler ist der, dass er zu tief unterhalb des Ellbogengelenks anfängt, so dass er mit den drei vorgeschriebenen Achtertouren das Gelenk nicht einwickeln kann. Ein weiterer Fehler ist der, dass er von der Anfangskreistour aus nicht gleich über die Ellenbeuge, sondern über die Streckseite des Gelenkes hinweggeht, und schliesslich führt er die Schlusskreistour nicht um das Gelenk herum, sondern legt dieselbe unterhalb desselben an.

Testudo cubiti reversa. (Tafel 23.)

Hatte man bei der vorhergehenden Tour unterhalb des Gelenkes angefangen und gegen das Gelenk hin gewickelt, daher der Name inversa, so beginnt man bei der testudo reversa mit der Kreistour gerade über das Gelenk hinweg. Im Anschluss an diese Kreistour macht man dann, indem man die Binde zunächst wieder von

der Kreistour aus über die Ellenbeuge hinführt, drei Achtertouren um das Gelenk herum in der Weise, dass man stets denjenigen Rand der vorhergehenden Tour zuwickelt, der vom Gelenk wegschaut. Man schliesst schliesslich die Binde mittelst einer Kreistour unterhalb des Gelenkes.

Der Fehler, der bei Anlegung dieser Testudo zuerst meist gemacht wird, ist der, dass die Binde von der Anfangstour aus nicht über die Ellenbeuge, sondern über das Olecranon hinweg in die Höhe geführt wird, so dass die Kreuzung der Binden nicht in der Ellenbeuge, sondern auf der hinteren Seite des Ellbogens stattfindet.

Die Einwicklung des Schultergelenkes

geschieht mittelst der Spica humeri. Wir haben hier wieder eine Spica humeri ascendens und eine Spica humeri descendens zu unterscheiden. Die

Spica humeri ascendens (Tafel 24)

beginnt mit einer Kreistour am oberen Ende des Oberarmes etwa in der Höhe des Deltoidesansatzes. Von der Kreistour aus umgeht man nun das Schultergelenk mit einer Achtertour in der Weise, dass man die Binde über die Schulterhöhe hinweg, schräg über die Brust, unter der Achsel der gesunden Seite hin, schräg über den Rücken und über die Schulterhöhe zurück zur Kreistour führt. Dieser ersten Achtertour lässt man noch zwei weitere Achtertouren folgen, derart, dass die einzelnen Touren sich parallel laufen und sich etwa zu zwei Dritteln so decken, dass immer der obere Rand der vorhergehenden Tour zugewickelt wird, die Touren also ascendieren. Nach Beendigung der letzten Achtertour wird eine die erste Kreistour deckende Schlusszirkeltour angelegt.

Spica humeri descendens. (Tafel 25.)

Man beginnt mit einer Kreistour um die Brust herum in der Höhe der Brustwarzen, und zwar muss man, wenn man die rechte Schulter des Patienten einwickeln will, mit dem Bindenanfang an der linken Seite des vor einem stehenden Patienten beginnen, muss also entgegengesetzt

Tafel 24.

Spica humeri ascendens.

Tafel 25.

Spica humeri descendens.

Tafel 26.

Einwickelung des ganzen Armes.

Tafel 27.

Stapes.

der vorher gegebenen allgemeinen Regel jetzt von rechts nach links wickeln. Will man die linke Schulter einwickeln, so beginnt man an der rechten Seite des Patienten und wickelt dann nach der allgemein giltigen Regel. Von der Kreistour um die Brust aus führt man die Binde schräg über die Brust zur Schulterhöhe hin, um die Achsel herum, wieder über die Schulterhöhe hinweg und schräg über den Rücken zur Anfangsstelle zurück. Dieser ersten Achtertour folgen dann noch zwei weitere Achtertouren, die einander parallel laufen und sich so decken, dass immer der untere Rand der vorhergehenden Tour zugewickelt wird. Den Schluss bildet wieder eine Kreistour um die Brust herum.

Die Einwicklung der ganzen oberen Extremität

(Tafel 26)

auch wohl nach dem Chirurgen Theden Involutio brachii Thedenii genannt, wurde in früheren Jahren namentlich dann angewendet, wenn beim Aderlass die Arteria cubitalis verletzt worden war. Heutzutage bedient man sich ihrer noch zur Compression bei ödematösen oder emphysematösen Anschwellungen des Armes, namentlich aber zum Zwecke der sogenannten Autotransfusion, um bei drohender Blutleere des Gehirns den Blutzufluss zu der betreffenden Extremität möglichst einzuschränken.

Die Einwicklung geschieht kunstgerecht derart, dass man zunächst sämtliche Finger in der vorherbeschriebenen Weise mittelst der Chiroteca completa einwickelt, dass man nach vollendeter Fingereinwicklung eine Spica manus ascendens anschliesst, dass man dann den Vorderarm mit einer Dolabra currens, respektive reversa bis zum Ellbogengelenk umgibt, an diesem angekommen, eine testudo cubiti inversa macht, und darauf am Oberarm mittelst der Dolabra currens, resp. reversa bis zum Deltoidesansatz anzusteigen und schliesslich mit einer Spica humeri ascendens zu endigen.

Verbände an der unteren Extremität.

Stapes oder Steigbügel. (Tafel 27.)

Der Stapes oder Steigbügel wurde in früheren Jahren angelegt, wenn man einen Aderlass am Fusse gemacht hatte. Man beginnt mit einer Kreistour an der Basis der Zehen, macht im Anschluss an diese Kreistour drei aufsteigende Touren einer Dolabra currens und schliesst mit einer Achtertour um das Fussgelenk herum.

Spica pedis descendens. (Tafel 28.)

Die Spica pedis descendens, auch wohl Sandalium genannt, entspricht der Spica manus. Man beginnt mit einer Kreistour oberhalb der Knöchel, macht dann, — den rechten Fuss des Patienten angenommen —, drei descendierende Achtertouren um das Fussgelenk herum in der Weise, dass die Binde jeweils von der Kreistour aus schräg über den Fussrücken, nach dem medialen Rand des Fusses hin, um die Fusssohle herum und wieder schräg über den Fussrücken zurück zur Kreistour läuft, und endigt mit einer Schlusskreistour oberhalb der Knöchel.

Testudo calcanei. (Tafel 29.)

Die Testudo calcanei entspricht der Testudo cubiti inversa. Man beginnt mit einer Kreistour um den Fussrücken herum, schliesst an diese Kreistour eine Achtertour um das Fussgelenk an, mit Kreuzung der Bindentouren gerade in der Mitte des Fussrückens, schliesst dieser ersten Achtertour noch zwei weitere Achtertouren an, so dass immer derjenige Rand der vorhergehenden Tour eingewickelt wird, der nach der Ferse schaut, und endigt mit einer Kreistour gerade über die Ferse hinweg.

Testudo calcanei zum Zwecke eines antiseptischen Verbandes. (Tafel 30.)

Wenn man die testudo calcanei in der eben beschriebenen Weise anlegt, so schmiegen sich die einzelnen Touren den Umrissen des Fusses an der Fuss-

Tafel 28.

Spica pedis descendens.

Tafel 29.

Testudo calcanei.

Tafel 30.

Testudo calcanei *mit Abschluss* der Ferse.

Tafel 31.

Einwickelung des Fusses von den Zehen an.



Tafel 32.

Einwickelung des ganzen Fusses.

sohle und über der Achillessehne niemals glatt an, klaffen vielmehr an diesen Stellen mehr oder weniger erheblich. Um auch an diesen Stellen einen festen Halt für die Binden zu gewinnen und ein Abrutschen derselben zu verhüten, fügt man der eigentlichen Testudo calcanei zweckmässig einige Bindentouren hinzu, welche einen wirklichen Abschluss der Ferse herbeiführen. Man erzielt einen solchen, wenn man die Binde nach vollendeter testudo calcanei schräg über den Fussrücken, zur Fusssohle hin, um diese herum, schräg gegen die Achillessehne hin, um diese herum, wieder zum Fussrücken zurück, jetzt nach der anderen Seite hin zur Fusssohle, um dieselbe herum, wieder schräg gegen die Achillessehne hin und zum Fussrücken zurückführt.

Einwicklung des Fusses von den Zehen an. (Tafel 31.)

Will man den Fuss einwickeln, ohne die Zehen mit in den Verband hereinzunehmen, so combiniert man zweckmässig den Stapes mit der testudo calcanei. Man beginnt an der Zehenbasis und wickelt zuerst regelrecht den Stapes; die Achtertour des Stapes, die man um das Fussgelenk herumgewickelt hat, benutzt man nun gleich als erste Achtertour für die testudo calcanei, wickelt also die Ferse nur noch mit zwei weiteren Achtertouren und der Schlusskreistour ein und fügt nach Vollendung dieser noch den sit venia verbo antiseptischen Fersenabschluss hinzu.

Einwicklung des ganzen Fusses. (Tafel 32.)

Will man auch die Zehen mit in den Verband hineinnehmen, so macht man den Verband zunächst genau in der eben beschriebenen Weise, dann aber führt man die Binde in Längstouren von der Ferse über die Zehen hinweg, zur Ferse zurück, indem man an einer Seite des Fusses anfängt und die Längstouren dann nach der andern Seite hinführt, so dass sie sich leicht decken, bis die Zehen vollständig eingeschlossen sind. Jede Spannung dieser Längstouren muss auf das sorgfältigste vermieden werden, da sonst ein unleidlicher Druck oder

Zehen entsteht. Sind die Zehen vollständig verdeckt, so wickelt man nun die Längstouren mit Kreistouren ein, die man von der Spitze der Zehen nach dem Unterschenkel hinführt. Einige Achtertouren um das Fussgelenk herum, die man einschaltet, sorgen dafür, dass die Binde nicht abrutscht.

Anstatt den Fuss in dieser Weise einzuwickeln, kann man auch so vorgehen, dass man mit einer Kreistour um die Malleolen beginnt und nun so lange absteigende Achtertouren nach der Art der Achtertouren bei der Spica pedis descendens macht, bis der ganze Fuss einschliesslich der Zehen eingewickelt ist. (Taf. 33.)

Einwicklung des Unterschenkels (siehe Tafel 39).

Zur Einwicklung des Unterschenkels legt man zunächst eine Kreistour um die Knöchel herum an und geht dann mit gleichmässig sich deckenden Touren einer Dolabra currens resp. reversa bis zum Knie in die Höhe.

Der Unterschenkel ist sehr geeignet, um an ihm das Anlegen der Renversé's zu üben. Bei einem gut sitzenden Verband sollen die Renversé's in einer Linie in der Mitte des Unterschenkels liegen.

Die

Einwicklung des Knies

geschieht mittelst der Testudo genu inversa oder reversa.

Testudo genu inversa. (Tafel 34.)

Bei leicht gebeugtem Knie macht man eine Kreistour am oberen Ende des Unterschenkels, führt die Binde über die Kniekehle hinweg, mit einer Achtertour um das Gelenk herum, lässt dieser ersten Achtertour noch zwei weitere Achtertouren folgen, die sich in der Kniekehle kreuzen und sich so decken, dass immer der nach dem Kniegelenk hinsehende Rand der vorhergehenden Tour zugewickelt wird, und schliesst mit einer Kreistour, gerade über die Patella hinweg. So hat man wieder eine Kreistour über das Gelenk hinweg und je drei von dieser aufsteigende und absteigende

Tafel 33.

Einwicklung des ganzen Fusses.

Tafel 34.

Testudo genu inversa.

Tafel 35.

Testudo genu reversa.

Tafel 32.

Einwickelung des ganzen Fusses.

Testudo genu reversa. (Tafel 35.)

An eine Kreistour gerade über die Patella hinweg schliesst man drei sich in der Kniekehle kreuzende Achtertouren an, die so um das Gelenk herumgeführt werden, dass immer der vom Kniegelenk wegsehende Rand der vorhergehenden Tour zugewickelt wird.

Die

Einwicklung des Oberschenkels (siehe Tafel 39)

geschieht vom Kniegelenk aus mit gleichmässig sich deckenden Touren einer Dolabra currens resp. reversa.

Zur

Einwicklung der Hüfte

bedient man sich der Spica coxae ascendens oder Spica coxae descendens.

Spica coxae ascendens. (Tafel 36.)

Von einer Kreistour am oberen Ende des Oberschenkels aus geht die Binde nach vorn schräg über den Unterleib hinweg um den unteren Teil des Rückens herum und wieder nach vorn zum Unterschenkel an die Kreistour zurück. So ist die erste Achtertour um das Hüftgelenk entstanden, welcher man noch zwei in gleicher Weise verlaufende aufsteigende Achtertouren folgen lässt. Die Schlusskreistour deckt die Anfangskreistour.

Spica coxae descendens. (Tafel 37.)

Man beginnt mit einer Kreistour, die in der Höhe des Nabels um das Abdomen herumgeht, und wickelt von sich aus von links nach rechts, wenn man die linke Seite des Patienten bandagieren will, von rechts nach links aber, wenn die rechte Hüfte eingewickelt werden soll. Von der Kreistour aus wickelt man die erste Achtertour um das Hüftgelenk herum, indem man die Binde schräg über das Abdomen herab, um den Oberschenkel herum und nach der Anfangsstelle der Kreistour hinführt. Dieser ersten Achtertour folgen dann noch zwei weitere absteigende Achtertouren und die Schlusskreistour um das Abdomen herum.

Spica coxae duplex. (Tafel 38.)

Die Spica coxae duplex, mittelst deren man beide Hüften einwickelt, ist ein praktisch ungemein wichtiger Verband, weil er unentbehrlich ist zur exakten Anlegung antiseptischer Verbände am unteren Ende des Leibes und an den Hüften. Man beginnt, von links nach rechts wickelnd, mit einer Kreistour um das Abdomen herum in der Höhe des Nabels, führt von der Kreistour aus die Binde schräg über den Unterleib hinweg nach abwärts zum linken Oberschenkel hin, um den Oberschenkel herum, zur Anfangsstelle der Kreistour zurück, nun schräg nach vorne um den rechten Oberschenkel herum, schräg nach aufwärts über den Unterleib hin, und um den Rücken herum zur Anfangsstelle der Binde zurück; so hat man je eine Achtertour um den linken und rechten Oberschenkel gewickelt. In der gleichen Weise führt man nun noch zwei weitere Achtertouren absteigend um beide Hüften herum und schliesst mit der Kreistour ebenso, wie man angefangen hat. Ist der Verband richtig angelegt, so hat man eine Spica descendens gerade in der Linea alba etwa in der Mitte zwischen Symphyse und Nabel und je eine Spica descendens um das rechte und linke Hüftgelenk.

Einwicklung der ganzen unteren Extremität. Involutio Thedenii extremitatis inferioris. (Tafel 39.)

Die Einwicklung der ganzen unteren Extremität braucht man in der Praxis häufiger als die Einwicklung der oberen Extremität. Die Indikation zur Einwicklung geben hydropische Anschwellungen, Varicenbildungen, Verletzungen, Anwicklung von Heftpflasterstreifen zu Streckverbänden und die Autotransfusion.

Man beginnt mit einer Kreistour an der Zehenbasis, wickelt von dieser aus regelrecht den Stapes, schliesst an diesen die Testudo calcanei mit gehörigem Fersenabschluss an, geht dann am Unterschenkel mit der Dolabra currens, resp. reversa in die Höhe, macht am Kniegelenk die Testudo genu inversa, wickelt dann den Oberschenkel mit einer Dolabra currens

Tafel 37.

Spica coxae descendens.

Tafel 38.

Spica coxae duplex.

Tafel 39.

Einwickelung des ganzen Beines.

Tafel 40.

Bier'sche Stauung.
(Ellenbogengelenk und Unterschenkel.)

resp. reversa ein und schliesst mit einer Spica coxae ascendens.

Die Uebung im regelrechten Einwickeln einer Extremität in der genannten Weise, so dass kein Teil der ganzen Extremität von den Zehen bis zur Leiste unbedeckt ist, dass die einzelnen Bindentouren sich dabei gut und gleichmässig decken, dass sie ferner weder zu fest noch zu locker angelegt sind, ist dem Anfänger sehr zu empfehlen, denn er soll sich beim Anlegen eines solchen Verbandes daran gewöhnen, die Ordnung und Genauigkeit zu beobachten, die er bei der Verrichtung anderer schwierigerer Handleistungen notwendig hat.

Mit Binden wird an den Extremitäten vielfach noch der Verband zur Erzeugung der Bier'schen Stauungshyperämie und der künstlichen Steigerung des Knochenwachstums nach Helferich verwendet.

In den letzten Jahren hat Bier eine Methode zur Behandlung der Zellentuberkulosen mit Stauungshyperämie angegeben, die schon heute Erfolge aufzuweisen hat und daher weitere Verbreitung verdient. Ich will daher hier eine genaue Beschreibung der Methode folgen lassen, zumal eine längere Zeit hindurch an den Epiphysen geübte Stauungshyperämie nach den Angaben von Helferich auch anregend auf das Längenwachstum der Knochen zu wirken vermag.

Will man eine Bier'sche Stauung regelrecht ausführen, so muss man in folgender Weise vorgehen: In der Abbildung (Tafel 40) ist angenommen, dass ein krankes Ellbogengelenk der Stauung unterworfen werden soll. Der Vorderarm wird von den Fingern bis unterhalb des Ellbogengelenkes mit einer Cambricbinde, am besten doppelt eingewickelt. Diese Einwicklung hat den Zweck, dass nicht an gesunden Stellen sich das Blut unnötig staut. Oberhalb des Ellbogengelenkes wird eine Gummibinde so fest und in so vielen sich genau deckenden Gängen umgelegt, dass das Gelenk blaurot wird und anschwillt. Damit die Gummibinde nicht drückt, ist sie mit mehreren Gängen einer weichen Mullbinde unterfüttert. Man probiert in jedem einzelnen Falle aus, wie

weit man die Stauung treiben kann, denn je hochgradiger die Stauung ist, desto besser wirkt sie. Aber das soll alles ohne Schmerzen abgehen, im Gegenteil, das schmerzhafte Gelenk wird meist schon nach einigen Stunden unempfindlich. Auch sollen die gestauten Glieder sich nicht kalt anfühlen. Kann man nicht gleich die notwendigen hohen Grade der Stauung hervorrufen, so gewöhnt man die Leute allmählich daran. Die unterfütternde Mullbinde muss ziemlich stramm angelegt werden, damit sie keine Falten wirft. Ebenso muss der erste Gang der Gummibinde stärker angezogen werden als die darauffolgenden, weil sie sich sonst ebenfalls faltet. Die Binde soll bei schmerzhaften Gelenken die erste Zeit über dauernd getragen werden. Sind die Schmerzen verschwunden oder bedeutend gebessert, so wendet man die Binde nur tags- oder nur nachtsüber an. Bei dauernder Benützung der Binde muss sie, um unangenehme Druckwirkungen zu vermeiden, etwa alle 12 Stunden an eine andere Stelle gesetzt werden. Setzt man die Binde tiefer, so muss sie nach kurzer Zeit etwas strammer angezogen werden, weil sie sich durch Verdrängung des Stauungsödemes schnell lockert. Niemals darf die Einwicklung das kranke Gelenk selbst mitbetreffen, weil sonst die Stauungshyperämie nicht genügend eintritt. Stets ist eine hochgradige Stauung zu erstreben, und bei genügender Geduld auch immer zu erreichen. Auf dem beigegebenen Bilde ist ein Verband zur Erzeugung der Stauungshyperämie auch noch an dem rechten Unterschenkel angelegt abgebildet um zu zeigen, dass ein solcher im gegebenen Falle, auch unterhalb oder oberhalb des Gelenkes an einem Gliede angelegt werden kann.

Nach der Einwicklung der Extremitäten lassen wir in der Regel die

Kopfverbände

üben.

Der schwierigste der Kopfverbände ist das

Capistrum simplex. (Tafel 41 u. 42.)

Das Capistrum simplex ist ursprünglich angegeben worden, um bei Brüchen einer Unterkieferhälfte den ge-

Tafel 41.

Capistrum simplex (gesunde Seite).

Tafel 42.

Capistrum simplex. (Kranke Seite.)

brochenen Unterkiefer gegen den Oberkiefer zu schienen. Es sollen, vor dem Ohr der kranken Seite beginnend, drei vom Kinn nach dem Scheitel aufsteigende Touren den Unterkiefer gegen den Oberkiefer andrücken. Zwischen der zweiten und der dritten dieser aufsteigenden Touren soll eine über das Kinn und den Nacken hingeführte Kreistour dem Verband eine genügende Festigkeit geben, (sog. Kinntour), während der Schluss oben durch eine Kreistour um Stirn und Hinterhaupt erzielt wird. Auf der gesunden Seite sollen nicht drei, sondern nur eine aufsteigende Tour vorhanden sein.

Der Verband wird in folgender Weise angelegt: Will man die linke Gesichtsseite einwickeln, so legt man das Bindenende horizontal über dem rechten Ohr des Patienten an und macht nun von seiner eigenen linken nach der rechten Seite wickelnd, also wie wir früher ausgeführt haben, von links nach rechts zunächst eine Kreistour um Stirn und Hinterhaupt. Zur Einwicklung der rechten Seite beginnt man umgekehrt über dem linken Ohr und wickelt mit seiner eigenen linken Hand von der eigenen rechten nach der eigenen linken Seite, also von rechts nach links. Von der Kreistour aus führt man nun die Binde um den Nacken herum unter das Kinn, vor dem Ohr der kranken Seite in die Höhe, auf den Scheitel, hinter dem Ohr der gesunden Seite herab unter das Kinn, vor dem Ohr der kranken Seite in die Höhe, so dass der nach dem Auge hinsehende Rand der ersten aufsteigenden Tour zugewickelt wird, auf den Scheitel zum Nacken hin, wobei man die Binde möglichst nahe gegen das Ohr der gesunden Seite hin führt, so dass sie nicht über den Hinterkopf abrutschen kann, um den Nacken herum, von der kranken Seite her über das Kinn (Kinntour) zum Nacken zurück, um den Hals herum, vor dem Ohr der gesunden Seite in die Höhe auf den Scheitel, um den Nacken herum, wobei man sich wiederum möglichst nahe dem Ohr hält, unter das Kinn und mit der dritten Tour aufsteigend zum Scheitel, von wo aus schliesslich mit einer Kreistour um Hinterhaupt und Stirn geschlossen wird.

Ich halte das Capistrum simplex nicht nur für einen

guten Verband zur Uebung, sondern auch für recht praktisch. Wem er recht, sozusagen in Fleisch und Blut übergegangen ist, der wird niemals Schwierigkeiten bei Anlegung eines antiseptischen Verbandes am Kopfe haben. Das Einzige, was an dem typischen Verbande unpraktisch ist, ist das Herabführen der ersten Tour hinter dem Ohr der gesunden Seite, denn die so angelegte Bindentour sucht nach vorn abzurutschen und schneidet, wenn man sie straff anzieht, an der Ohrmuschel ein. Es ist daher praktisch besser, sie vor dem Ohr der gesunden Seite herabzuführen.

Capistrum duplex. (Tafel 43.)

Bei dem Capistrum duplex, das ursprünglich für doppelseitige Unterkieferfrakturen bestimmt war, sollen auf jeder Seite drei vom Kinn nach dem Scheitel aufsteigende Touren gewickelt werden; zwischen zweiter und dritter Tour soll die Kinntour eingeschaltet sein. Man beginnt, indem man das Bindenende auf die Mitte des Scheitels legt, führt die Binde am äusseren linken Augenwinkel vorbei nach dem Kinn, geht unter diesem herum zum äusseren Augenwinkel der anderen Seite und zum Scheitel zurück. So hat man gewissermassen eine Kreistour um Scheitel und Kinn gemacht. Vom Scheitel aus geht man nun, sich möglichst nahe dem linken Ohr haltend, nach dem Nacken hin, um diesen herum unter das Kinn, vor dem linken Ohr in die Höhe, so dass der nach diesem Ohr hinschauende Rand der vorhergehenden Tour zugewickelt wird, auf den Scheitel, um den Nacken herum, wieder möglichst nahe dem rechten Ohr, unter das Kinn, vor dem rechten Ohr in die Höhe auf den Scheitel, um den Nacken, von der rechten Seite her über das Kinn (Kinntour), um den Nacken, um den Hals, unter das Kinn, vor dem linken Ohr mit der dritten Tour in die Höhe auf den Scheitel, um den Nacken herum unter das Kinn und vor dem rechten Ohre mit der dritten Tour in die Höhe auf den Scheitel; schliesslich Schlusskreistour um Hinterhaupt und Stirn.

Zur Deckung der Schädelwölbung bedient man sich der

Tafel 43.

Capistrum duplex.

Tafel 44.

Mitra Hippocratis.

Tafel 45.

Monoculus.

Mitra Hippocratis. (Tafel 44.)

Die Mitra Hippocratis wird mittelst einer doppelköpfigen Binde angelegt. Man fasst die beiden Bindenköpfe, wickelt sie etwas ab und legt die Mitte der Binde auf die Stirn, so dass der untere Bindenrand die Augenbrauen berührt. Nun führt man die beiden Bindenköpfe nach dem Nacken hin, kreuzt sie hier, lässt nun den einen Bindenkopf, den wir als Bindenkopf a bezeichnen wollen, von der Mitte des Nackens über die Mitte des Scheitels hinweg bis zur Nasenwurzel, den anderen, den Bindenkopf b, aber kreisförmig um Hinterhaupt und Stirn verlaufen. Diese Kreistour deckt über der Glabella die von hinten nach vorn verlaufende, mit dem Bindenkopf a beschriebene Tour. Ist die Fixierung des Bindenkopfes a durch den Bindenkopf b erfolgt, so führt man den Bindenkopf a von der Glabella aus wieder über den Scheitel nach dem Nacken zurück, wobei diese zweite Tour die erste zu etwa einem Drittel deckt, und fixiert die Tour hinten am Nacken wieder durch eine mit dem Bindenkopf b ausgeführte Kreistour. Nun führt man den Bindenkopf a wieder nach vorn zur Glabella hin, indem man jetzt die andere Seite der ersten von hinten, nach vorn ziehenden Tour zudeckt. So wickelt man nun allmählich die ganze Schädelwölbung ein, indem der Bindenkopf a immer nur von hinten nach vorn, resp. von vorn nach hinten, und zwar stets abwechselnd nach beiden Seiten hin geführt wird, während der Bindenkopf b immer nur Kreistouren um Hinterhaupt und Stirn macht, um die Touren des Bindenkopfes b zu fixieren. Soll der Verband gut sitzen, so muss die Kreuzung der Bindetouren am Hinterkopf stets unterhalb der Protuberantia occipitalis erfolgen.

Die

Verbände für die Augen

sind der Monoculus und Binoculus.

Monoculus. (Tafel 45.)

Der Monoculus, auch wohl Druckverband für das Auge genannt, beginnt mit einer Kreistour um Stirn

und Hinterhaupt. Will man das linke Auge einwickeln, so legt man das Bindenende über dem rechten Ohr an und wickelt von sich aus von links nach rechts; um das rechte Auge einzuwickeln, legt man das Bindenende über dem linken Ohr an und wickelt von sich aus mit der linken Hand von rechts nach links. Man verhält sich also hier genau so wie beim Capistrum simplex.

Von der Kreistour aus führt man die Binde zur Glabella, über das kranke Auge hinweg, unter dem Ohr der kranken Seite vorbei, um das Hinterhaupt herum, schräg aufsteigend über dem Ohr der gesunden Seite zur Kreistour zurück, deren oberen Rand man zudeckt, und zur Glabella hin. Auf dieser kreuzt man nun die erste Tour, führt die Binde wieder über das kranke Auge hinweg, so zwar, dass man nun den unteren Rand der vorhergehenden Tour umwickelt, geht wieder unter dem Ohr der kranken Seite hin, um das Hinterhaupt herum, steigt steil, den oberen Rand der vorhergehenden Tour zuwickelnd, über die Seitenfläche des Schädels herauf, führt die Binde dann wieder zur Glabella herunter, kreuzt hier wieder, geht wieder über das Auge hinweg, wickelt jetzt wieder den unteren Rand der vorhergehenden Tour zu, geht unter dem Ohr der kranken Seite um den Hinterkopf herum zu der Stelle zurück, wo man angefangen hatte, und schliesst von hier aus mit einer Kreistour um Stirn und Hinterhaupt.

Bei dem richtig angelegten Monoculus sollen demnach die Bindentouren, die oberhalb des gesunden Ohres liegen, aufsteigend, die das Auge zudeckenden dagegen absteigend verlaufen. In der Praxis wird der Monoculus gewöhnlich mit einer Flanellbinde angelegt, weil sich diese den Contouren des Kopfes besser anschmiegt.

Binoculus. (Tafel 46.)

Soll der Verband gleichzeitig beide Augen decken, so kommt der Binoculus zur Verwendung. Dieser wird analog der Spica coxae duplex so angelegt, dass man abwechselnd das eine und das andere Auge zudeckt.

Tafel 46.

Binoculus.

Tafel 47.

Fascia nodosa.

Man beginnt mit einer Kreistour um Stirn und Hinterhaupt, führt — von sich aus — von links nach rechts wickelnd, die Binde über das linke Auge hinweg, unter dem linken Ohr vorbei, um das Hinterhaupt herum, unter dem rechten Ohr vorbei, über das rechte Auge hinweg, über die Glabella, oberhalb des linken Ohres um den Kopf herum, über dem rechten Ohr, den oberen Rand der vorhergehenden Tour zuwickelnd, wieder zur Glabella zurück. So wickelt man nun noch zwei weitere Touren, dass die Kreuzung der Bindentouren jedesmal auf der Glabella statthat und dass die oberhalb der Augenbrauen liegenden Touren aufsteigen, die die Augen deckenden Touren dagegen absteigen. Liegt der Verband gut, so strahlen die einzelnen Bindentouren sternförmig von der Glabella aus.

Fascia nodosa. (Tafel 47.)

Die Fascia nodosa oder Knotenbinde wurde in früheren Jahren angewendet zur Compression nach einem Aderlass an der Arteria temporalis. Heutzutage verwendet man sie noch gelegentlich zur Compression bei einem sich bildenden Aneurysma dieser Arterie, oder als Notverband bei Verletzungen derselben.

Der Verband wird mit einer doppelköpfigen Binde angelegt. Die Mitte dieser doppelköpfigen Binde legt man über das Ohr der gesunden Seite, führt beide Bindenköpfe über Stirn und Hinterhaupt nach der, wie wir annehmen wollen, verletzten Seite hin, kreuzt hier die Bindenköpfe, so dass der um die Stirn verlaufende Bindenkopf nach dem Scheitel zu, der vom Hinterhaupt kommende dagegen nach dem Kinne hin sieht, führt die Binden vom Scheitel resp. Kinn vor dem gesunden Ohr vorbei über den Scheitel resp. das Kinn hin zur verletzten Stelle, kreuzt hier wieder und wiederholt diese Touren mehrere Male. Durch die verschiedenen Kreuzungen der Bindentouren entstehen Knoten; soll der Verband seinen Zweck erfüllen, so müssen die einzelnen Knoten genau aufeinander zu liegen kommen; ist aber das letztere der Fall, so kann man wirklich eine starke Compression mit dem Verbande ausführen.

Funda nasi. (Tafel 48.)

Unter Funda oder Schleuder versteht man einen Bindenstreifen von etwa doppelt so grosser Breite, als sie gewöhnlich am Kopf verwendet werden, den man von beiden Seiten her bis auf ein etwa handbreites Mittelstück einreisst.

Zur Ausführung der Funda nasi legt man nun dieses Mittelstück auf die Nase, nimmt die beiden oberen Streifen, führt sie unter den beiden Ohren hinweg nach dem Hinterhaupt und knotet sie hier. Dann nimmt man die beiden unteren Streifen, führt sie in gleicher Weise über den Ohren nach dem Hinterhaupt, um sie dort gleichfalls zu knoten.

Funda maxillae. (Tafel 49.)

Zur Ausführung der Funda maxillae, auch Kinnschleuder genannt, legt man das Mittelstück auf das Kinn, so dass es mit seinem oberen Rand bis unter die Lippe, mit seinem unteren bis unter das Kinn reicht, Die beiden oberen Streifen werden nun direkt nach hinten zum Nacken geführt, hier gekreuzt und nach der Stirn geleitet, wo sie geknüpft werden. Die beiden unteren Streifen werden, der eine vor dem einen Ohr, der andere hinter dem andern Ohr nach dem Scheitel hingeführt und hier ebenfalls geknotet.

Verbände am Rumpf.

Die

Petit'sche Achtertour und die Stella dorsi. (Tafel 50.)

Wenn man den Patienten, mit dem Rücken gegen sich gekehrt, vor sich hinstellt, wenn man dann die Binde vorn an der linken Achsel anlegt, eine Kreistour um die linke Schulter und Achselhöhle macht, die Binde dann über die Schulterhöhe hinweg, schräg über den Rücken, unter die Achselhöhle deranderen Seite, über die entsprechende Schulter hin, schräg über den Rücken zurück, durch die Achselhöhle der linken Seite hindurch und

Tafel 48.

Funda nasi.

Tafel 50.

Stella dorsi.

Tafel 51.

Stella pectoris.



Tafel 52.

Stella pectoris et dorsi.

zur Höhe der linken Schulter zurück führt, so hat man die Petit'sche Achtertour ausgeführt.

Wickelt man nicht eine solche Achtertour, sondern drei derselben, die von oben nach unten hin absteigen, so hat man die Stella dorsi oder auch wohl Sternbinde oder Spica dorsi gemacht. Die Kreuzungsstellen der Bindentouren liegen gerade in der Mitte des Rückens über den Dornfortsätzen.

Stella pectoris. (Tafel 51.)

Man stellt den Patienten vor sich hin, macht unterhalb der Brustwarzen eine Kreistour, steigt von der rechten Achselhöhle aus schräg über die vordere Brustwand zur Schulterhöhe der linken Seite hin, geht an der hinteren Fläche der Schulter herab zur linken Achselhöhle hin und steigt von hier aus in gleicher Weise wie eben nach links, so jetzt schräg über die vordere Brustwand nach der rechten Schulterhöhe hin und über diese hinweg nach der Achsel der rechten Seite. So hat man die erste Achtertour vollendet. Die Kreuzungsstelle der Binden liegt gerade auf der Mitte des Brustbeins. Der ersten Achtertour folgen nun noch zwei weitere aufsteigende Achtertouren, worauf der Verband mit einer Kreistour um die Brust geschlossen wird.

Stella pectoris et dorsi. (Tafel 52.)

Man legt das Bindenende in der Axillarlinie der rechten Seite an, macht von links nach rechts wickelnd, unterhalb der Brustwarzen eine Kreistour um die Brust und den Rücken, geht dann von der rechten Achsel aus schräg über die Brust nach der linken Schulter in die Höhe, an der Hinterfläche der linken Schulter hinab, durch die linke Achselhöhle hin, schräg über die Brust zur rechten Schulter hin, hinten an der rechten Schulter herunter, durch die rechte Achselhöhle hindurch, vorn an der rechten Achsel in die Höhe zur rechten Schulterhöhe, schräg über den Rücken zur linken Achselhöhle hin, vorn an der linken Achsel in die Höhe, zur linken Schulterhöhe und schräg über den Rücken hinweg nach der rechten Achselhöhle hin. So hat man vier Kreuz-

ungen[illegible]gt: vorn über dem Brustbein, hinten über der W[illegible]ule und auf der Höhe beider Schultern.

In der gleichen Weise wie eben beschrieben, wickelt man nun noch zwei weitere aufsteigende doppelte Achtertouren über Brust und Rücken und schliesst mit einer Kreistour um Brust und Rücken.

Suspensorium mammae. (Tafel 53 u. 54.)

Das alte Suspensorium mammae ist ein schlecht sitzender Verband, der wohl nur noch zu Uebungszwecken gelehrt wird. (Tafel 53.) [illegible]r[illegible]rband soll den Zweck haben, die weibliche Bru[illegible]e Höhe zu heben und sie in der Höhe zu halten. [illegible]beginnt in der Achselhöhle der kranken Seite, [illegible] das Bindenende von einem Assistenten einstweilen [illegible]en lässt, geht mit der Binde dem unteren Rande de[illegible]ıstdrüse entlang, schräg über die Vorderfläche des Br[illegible]rbes zur Schulterhöhe der gesunden Seite und umgi[illegible]ese Schulter mit einer Achtertour, welche wieder t[illegible] der Achselhöhle der kranken Seite hinführt. Von hier aus führt man nun die Binde jetzt dem oberen Rand der Brustdrüse entlang, vorn über die Brust, geht mit einer Achtertour um die gesunde Schulter herum und wieder zur Achselhöhle der kranken Seite zurück. Man führt nun von hier aus in gleicher Weise noch je eine Bindentour an der unteren und oberen Brusthälfte, welche den nach der Brustwarze hinsehenden Rand der vorhergehenden Tour zudeckt und schliesst mit einer Kreistour, die gerade über die Mitte der kranken Brust hinwegläuft und die Brustwarze zudeckt.

Wie schon gesagt, sitzt das in dieser Weise ausgeführte Suspensorium sehr schlecht. Will man wirklich eine Brustdrüse in die Höhe heben und hoch halten, so verfährt man besser in folgender Weise. Man lässt die Brustdrüse, während man selber den Verband anlegt, von einem Assistenten von unten her mit seiner ganzen Hand in die Höhe heben. Man beginnt nun mit einer Kreistour, die man unterhalb der Brustdrüse links nach rechts wickelnd um die vordere und

Tafel 53.

Altes, aber unpraktisches Suspensorium mammae.

Tafel 54.

Praktisches Suspensorium mammae.

Tafel 55.

Suspensorium mammae duplex.
(Jede Brustdrüse ist für sich eingewickelt.)

Tafel 56.

Suspensorium mammae duplex.
(Beide Brustdrüsen sind abwechselnd eingewickelt worden.)

hintere Seite des Thorax herumführt. Nach vollendeter Kreistour geht man von der kranken Seite aus, dem unteren Rande der Brustdrüse entlang, schräg über die Vorderseite des Thorax hin, umgibt die gesunde Schulter mit einer Achtertour und kommt so wieder zu der Stelle zurück, wo man mit der Binde begonnen hatte.

Nun lässt man ganz in der gleichen Weise, wie diese eben geschilderte Tour aufsteigend noch eine Reihe gleicher Touren folgen, bis die ganze Brustdrüse vollständig zugedeckt ist. (Tafel 54.)

Suspensorium mammae duplex. (Tafel 55 u. 56.)

Soll eine Tragbinde für beide Brustdrüsen geschaffen werden, so ist es am einfachsten, jede Drüse für sich einzuwickeln, zuerst die rechte in der eben angegebenen Weise; man macht aber nach vollendeter Einwicklung der rechten Brust mit der letzten Tour keine Achtertour mehr um die linke Schulter herum, sondern führt die Tour von der Höhe der linken Schulter direkt unter der linken Brustdrüse hinweg, um nun die aufsteigenden Achtertouren um das rechte Schultergelenk anzuschliessen, bis auch die linke Brustdrüse vollständig zugedeckt ist. (Tafel 55.)

Eine andere gute Art, ein Suspensorium für beide Brustdrüsen zu schaffen, ist die, dass man in der vorher angegebenen Weise eine Stella pectoris et dorsi anlegt, jetzt aber nicht wie beim typischen Verband nur drei Achtertouren wickelt, sondern so viele Touren, dass beide Brustdrüsen vollständig zugedeckt werden. (Taf. 56.)

Compressorium mammae simplex und duplex.

Will man eine oder beide Brustdrüsen nicht nur in die Höhe heben, sondern will man zugleich einen Druckverband für die Brüste machen, so wickelt man zunächst in einer der zuletzt angegebenen Formen ein einfaches Suspensorium und schliesst dann an dieses von unten nach oben aufsteigend Zirkeltouren, welche um Brust und Rücken verlaufen.

Wir kommen jetzt zu einigen Rumpfverbänden, die

zur Behandlung von Schlüsselbeinbrüchen angegeben worden sind und sich namentlich zu Uebungszwecken gut verwerten lassen.

Desault'scher Verband. (Tafel 57, 58, 59.)

Wie alle anderen Verbände bei Clavicularfrakturen verfolgt auch der Desault'sche Verband den Zweck, die nach dem Bruch des Schlüsselbeines nach unten, innen und vorn gesunkene Schulter nach oben, aussen und hinten zu heben und in dieser Lage zu erhalten.

Um den genannten Zweck zu erreichen, wird der Desault'sche Verband in drei Touren angelegt.

Erste Tour. (Tafel 57.) Ein keilförmiges Kissen (siehe Tafel 97a) wird mit der Basis nach oben in die Achselhöhle der verletzten Seite gelegt. Dieses Kissen soll nun durch die erste Tour des Desault'schen Verbandes an den Thorax fixiert werden. Man steckt zu diesem Zweck das Bindenende mit einer Stecknadel an das Kissen an und führt nun mit dem Bindenkopf, den man schräg vom Kissen aus vorn über die Brust gehen lässt, zunächst eine Achtertour um die gesunde Schulter aus. Die Achtertour führt wieder zum Bindenanfang zurück, und nun vollführt man mit der Binde um Kissen und Thorax, von der Achselhöhle angefangen, gleichmässig sich zu etwa zwei Dritteln deckende absteigende Touren, bis das ganze Kissen von oben bis unten hin völlig eingehüllt ist.

Zweite Tour (Tafel 58.): Die zweite Tour soll den Oberarm über dem Kissen an den Thorax anbandagieren. Der untere Teil des Oberarms soll genau an die Seitenfläche des Brustkorbes zu liegen kommen, damit die Schulter wirklich über dem das Hypomochlion bildenden Kissen nach aussen abgehebelt wird.

Die Anfänger machen in der Regel den Fehler, dass sie das untere Humerusende zu weit nach vorn legen. Der Vorderarm liegt rechtwinklig gebeugt quer auf der Vorderfläche des Rumpfes auf. Hat der Arm seine richtige Stellung, so wickelt man ihn nun mit Kreistouren um den Thorax an denselben an. Man legt das Ende einer neuen Binde vorn an der gesunden Achselhöhle

Tafel 57.

I. Tour des Desault'schen Verbandes.

Tafel 58.

II. Tour des Desault'schen Verbandes.

Tafel 59.

III. Tour des *Desault*'schen Verbandes.

an, führt die Binde in einer Kreistour um den Thorax herum, so dass der obere Rand der Binde in die Höhe des Acromions der verletzten Seite zu liegen kommt, und wickelt nun die folgenden Kreistouren so, dass die absteigenden Touren die Brust und den Arm von der Höhe der Schulter bis zur Spitze des Olecranon zudecken.

Dritte Tour (Tafel 59.): Die dritte Tour des Desault soll der Wiederkehr der Dislokation der Bruchenden entgegenwirken und namentlich den Ellbogen von unten her stützen; sie soll ferner der Hand eine Stütze geben. Durch den regelrechten Verband sollen auf der Brust und dem Arm zwei Dreiecke gebildet werden, von denen das innere kleiner ist, als das äussere. Der Verband lässt sich am leichtesten anlegen, wenn man sich merkt, dass man die Bindentouren stets von der Achselhöhle der gesunden Seite zur kranken Schulter, von dieser nach dem Ellbogengelenk der kranken Seite und von diesem aus wieder zur Achselhöhle der gesunden Seite hinführt.

Dem Anfänger empfehle ich zur Unterstützung des Gedächtnisses sich ein kleines Verschen zu merken. Wenn 1 die gesunde Achselhöhle bezeichnet, 2 die Schulterhöhe der kranken Seite, und 3 das Ellbogengelenk der kranken Seite, so wird der Anfänger niemals irre werden, wenn

„Von 1 nach 2 nach 3
der innere Rand bleibt frei."

Eine andere gute Hilfe für das Gedächtnis ist das Wort „Asche", welches in der richtigen Reihenfolge die Anfangsbuchstaben der drei Stützpunkte für den Verband, Achsel, Schulter und Ellbogen, enthält.

Man legt also das Bindenende vorn an der gesunden Achselhöhle an, führt die Binde schräg über den Thorax zur Schulterhöhe der kranken Seite, geht über die kranke Schulter hinweg an der hinteren Seite des Armes herunter, um das Ellbogengelenk herum, schräg über die Brust in die Höhe zur Achselhöhle der gesunden Seite, durch dieselbe hindurch, schräg über den Rücken zur Schulterhöhe der kranken Seite und an der vorderen Seite des Armes herab zum Ellbogengelenk. So ist das erste Drei-

eck auf der Brust entstanden. Man führt nun die Binde um das Ellbogengelenk herum, schräg über den Rücken zur gesunden Achselhöhle zurück, durch dieselbe hindurch, und so ist man wieder da angekommen, wo man angefangen hat. Jetzt wickelt man das zweite grössere Dreieck, indem man die Binde in der gleichen Weise von der gesunden Achselhöhle nach der kranken Schulter und dem Ellbogengelenk der kranken Seite hin und nach der Achselhöhle der gesunden Seite zurückführt. Dadurch, dass man den inneren Rand der vorhergehenden Tour freilässt, wird das äussere Dreieck grösser als das innere.

Ist das zweite Dreieck vollendet, dadurch, dass man die Binde von der Schulterhöhe her an der Vorderseite des Armes herab zum Ellbogengelenk hingeführt hatte, so schliesst man den Verband, indem man noch ein Tragband für die Hand hinzufügt, in der Weise, dass man die Binde um das Ellbogengelenk herum, schräg über den Rücken in die Höhe, über die gesunde Schulter hinweg, vorn am Thorax herab, um das Handgelenk und zur kranken Schulter hinführt; auf dieser wird das Bindenende festgesteckt.

Während man den vollständigen Desault'schen Verband, wie er eben beschrieben wurde, mit allen seinen Touren in der Praxis wohl nur sehr selten anlegt, kommt die dritte Tour für sich allein recht häufig in Verwendung, und zwar in all' den Fällen, in denen man den Oberarm für längere Zeit an den Brustkorb zu befestigen genötigt ist.

Velpeau'scher Verband. (Tafel 60.)

Der Velpeau'sche Verband ist auch zur Behand- von Schlüsselbeinbrüchen angegeben worden. Vel- beseitigt die Dislokation des gebrochenen Schlüssel- dadurch, dass er die Hand der verletzten Seite e gesunde Schulter flach auflegen lässt, während die des Ellbogens auf die Mitte der Brust zu liegen t. In dieser Velpeau'schen Lage des Armes wird er Verband in folgender Weise befestigt. Man macht

Tafel 60.

Velpeau'scher Verband.

Tafel 61.

Modifikation des Velpeau'schen Verbandes von Dulles.

Tafel 62.

Sayre'scher Heftpflasterverband.

Tafel 63.

Schoenborn's Modifikation des Sayre'schen Heftpflasterverbandes.

zunächst eine Kreistour um Arm und Brust in der Höhe der Achselhöhe, geht dann von der Achselhöhle der gesunden Seite aus nach hinten, führt die Binde schräg über den Rücken, über die kranke Schulter hinweg, an der vorderen Seite des Oberarms der kranken Seite nach abwärts, um die Streckseite des Ellbogengelenkes herum zur gesunden Seite des Brustkorbes hin, macht von hier aus wieder eine Kreistour um Arm und Thorax, und zwar in der Höhe des Ellbogengelenkes, und wiederholt nun in gleicher Weise, jedesmal hinten von der gesunden Achselhöhle anfangend, und hinten an dem Rücken in die Höhe steigend, die von der Schulterhöhe nach dem Ellbogengelenk verlaufenden und die sich an diese anschliessenden Kreistouren in der Weise, dass die schrägen Bindengänge sich etwa zu zwei Dritteln deckend von dem Acromion gegen den Hals hin verlaufen, während die Kreistouren vom Ellbogengelenk gegen die Schulter hin aufsteigen.

Eine zweckmässige Verbesserung des Velpeau'schen Verbandes ist von Dulles angegeben worden. (Tafel 61.) Man führt die Binde von der Achselhöhe der gesunden Seite aus vorn über die Brust, über die Schulterhöhe der verletzten Seite hinweg, an der Rückseite des Armes herab zum Ellbogen, an der vorderen Seite desselben nach aufwärts zur gleichen Schulter zurück, über den Rücken zur Achselhöhle der gesunden Seite, quer über die Brust und den Arm zum Ellbogen und von hier quer über den Rücken zur Achselhöhle zurück. Diese Gänge werden nun bis zur völligen Einwicklung der Brust und des Oberarms stets in der Art wiederholt, dass die schräg über die Brust von der Achsel zur Schulter verlaufenden Touren absteigen, während die Touren um den Arm von dem Acromion gegen den Hals hin verlaufen und die Kreistouren um den Thorax und den Oberarm vom Ellbogen gegen die Schulter hin aufsteigen.

Sayre'scher Heftpflasterverband. (Tafel 62.)

Der Verband, der heute wohl am meisten zur Fixation des Armes an den Thorax bei Verletzungen des Schultergürtels angewendet wird, ist der Sayre'sche Heftpflasterverband.

Derselbe wird angelegt mit drei langen, etwa 2—3 Finger breiten Heftpflasterstreifen. Den ersten Streifen klebt man am äusseren Rand des Sulcus bicipitalis internus des zu fixierenden Oberarms an, führt ihn dann leicht spiralig nach hinten und oben über die äussere Fläche des Oberarmes, über den Rücken und unter die Achsel der gesunden Seite, und klebt ihn unter der Brustwarze fest. Diese Tour hebt die Schulter und zieht sie kräftig nach hinten.

Der zweite Streifen hat ebenfalls die Aufgabe des Schulterhebens. Er geht von der gesunden Schulter schief über die Brust nach dem rechtwinklig gebogenen Vorderarm hin, schlägt sich um den Ellenbogen herum und geht von da schief über den Rücken zurück nach der Anfangsstelle auf der gesunden Schulter.

Der dritte Streifen wird mit seinem Anfange direkt auf die Frakturstelle aufgeklebt, läuft dann an der Vorderseite des Thorax herunter, umgibt das Handgelenk und geht wieder zur Frakturstelle zurück. Er dient als Tragband für die Hand, übt aber vor allem einen direkten Druck auf die Fragmente aus.

Schoenborn's Modifikation des Sayre'schen Heftpflasterverbandes. (Tafel 63.)

Schoenborn braucht nur 2 Heftpflasterstreifen. Den ersten legt er wie Sayre an, nur klebt er den Streifen nicht an die Haut an, sondern befestigt das um den Arm in Form einer Schleife herumgeschlungene Ende mit einer Stecknadel. Der Arm erhält dann die Velpeau'sche Lage. Der zweite wird nun analog dem Velpeau'schen Verbande an dem Radialrande des Vorderarmes dicht unterhalb des Ellbogengelenks angeklebt, um den Ellbogen, wie die Velpeau'sche Schlinge, mit Vermeidung des Epicondylus internus herumgeführt und auf der gesunden Schulter angeklebt. Der Handrücken und die Vola der auf der Schulter aufruhenden Hand müssen gut mit Watte gepolstert sein.

Tafel 64.

Landerer's Modifikation des Sayre'schen Heftpflasterverbandes.

Landerer's Modifikation des Sayre'schen Heftpflasterverbandes. (Tafel 64.)

Landerer legt den Angriffspunkt des Streifens nicht an den Oberarm, sondern an die kranke Schulter selbst. Der erste Streifen besteht nach Landerer aus 3 Teilen, einem ungefähr handgrossen, vom Rande her fächerförmig gespaltenen Stück Heftpflaster, einem vorn an dieses angenähten, 20—30 cm langen Streifen elastischen Gurtes und einem wiederum an diesen angenähten, 60—80 cm langen und 5—6 cm breiten Heftpflasterstreifen. Dieser so zusammengesetzte Streifen wird nun, nachdem ein fest zusammengedrehter Wattekeil in die Achsel gelegt worden ist, mit seinem handförmigen Ende unmittelbar auf die Schlüsselbeingegend aufgeklebt, jedoch so, dass das Fragment von jeder Zugwirkung frei bleibt. Der Streifen wird dann über die Schulterhöhe weggeführt, kräftig angezogen und wie der erste Sayre'sche Streifen nach der Vorderfläche der Brust geleitet. Darüber kommt dann der zweite Sayre'sche Streifen und dann ein diesen fixierender Bindenverband.

v. Bünger's Verband für Schlüsselbeinbrüche (Taf. 65 u. 66.)

Einen recht brauchbaren Verband zur Behandlung der Schlüsselbeinbrüche hat in den letzten Jahren v. Büngner angegeben. Zu diesem Verbande bedarf man einer dreiteiligen TT-Binde und einiger Sicherheitsnadeln.

Das 60 cm lange und 4 cm breite Querstück, mit dem 120 cm lange und 10 cm breite Längsstreifen so verbunden sind, dass der mittlere rechtwinklig, die beiden anderen etwas schräg von demselben abgehen, wird um die gesunde Schulter gelegt und festgeschnallt, der mittlere und ein seitlicher Längsstreifen liegen an der Hinterseite.

Man ergreift nun zunächst den Mittelstreifen (Tafel 65), führt ihn über den Rücken weg durch die Achselhöhle um den oberen Teil des Oberarmes der kranken Seite herum und darauf zum Ausgangspunkt zurück. Hier wird das Bindenende angesteckt. Auf diese Weise wird der kranke Arm direkt nach hinten gezogen. Darauf

wird der untere Streifen ebenfalls über den Rücken weg und in gleicher Richtung um den Oberarm der kranken Seite herumgeführt, aber so, dass er den untersten Teil desselben umgreift und um die Ellbogen gewunden wird, um dann ebenfalls zum Ausgangspunkt zurückzukehren. Auf diese Weise wird der kranke Arm teils, wie vorher, nach hinten gezogen, teils angehoben. (Der zweite Streifen b wird also genau wie der erste Streifen a angelegt, nur dass er statt um den obersten Teil des Oberarmes um den untersten Teil desselben und um den Ellbogen zu liegen kommt.)

Der obere und letzte Streifen endlich (Tafel 66) geht als Mitellatour nach vorn, unterstützt das Handgelenk und geht darauf, die Bruchfragmente niederdrückend, über die Frakturstelle und die verletzte Schulter hinweg an die Rückseite der letzteren, um hier an den beiden ersten Streifen a und b befestigt zu werden, wodurch diese in ihrer Lage so sicher fixiert werden, dass sie sich in keiner Weise verschieben können.

Der Verband zieht die Schulter der verletzten Seite nach hinten, oben und aussen; er übt diese Wirkung in jeder Stellung des Körpers, er gestattet die ambulante Behandlung, denn er kann sowohl über als unter den Kleidern getragen werden; er lässt den Thorax frei und befestigt den Arm der kranken Seite nicht an denselben. Man legt den Verband bei der Stellung des betreffenden Armes an, bei welcher die Dislokation am Schlüsselbein und an der Schulter vollständig ausgeglichen ist. Der erste Gang des Verbandes muss so stark angezogen werden, dass der mediale Rand des Schulterblattes so weit der Wirbelsäule genähert werde, dass der Abstand beider Schulterblätter von dieser wiederum der gleiche ist. Auch die scheinbare Verlängerung des Armes der verletzten Seite muss nach Anlegung des zweiten Ganges vollständig beseitigt sein. Unmittelbar nach der Verletzung wird der Verband auf blosser Haut angelegt und bleibt durchschnittlich bis zum zehnten Tage, d. h. bis zur beginnenden Consolidation Tag und Nacht liegen.

Tafel 65.

Verband zur Behandlung der Schlüsselbeinbrüche
nach v. Büngner.

Tafel 66.

Verband nach v. Büngner.

Es ist zweckmässig, an den Stellen des Armes der verletzten Seite, wo der Verband anliegt, eine Wattebinde unterzulegen und ebenso die Bruchstelle mit Verbandwatte zu polstern. Ist der provisorische Callus gebildet und eine Verschiebung der Bruchenden nicht mehr zu fürchten, dann wird der Verband für die Nacht abgenommen und nur noch am Tage, und zwar über den Kleidern, getragen, so dass dem Kranken das Verlassen der Wohnung ermöglicht ist. Nach Bildung des knorpeligen Callus sichert der über der Kleidung getragene Verband ausreichend die Feststellung der Bruchenden; für die Nacht wird er mit einer einfachen Mitella vertauscht und der Rücken des Kranken noch eine Zeit lang durch ein Rollkissen unterstützt, damit die verletzte Schulter möglichst nach hinten sinke.

Ein und derselbe Verband kann sowohl an der rechten wie an der linken Seite des Kiefers verwandt werden; die Schnalle des Quergurtes der TT-Binde kommt nur auf der einen Seite vorn, auf der anderen hinten zu liegen.

2. Tücherverbände.

Tücherverbände wurden in früheren Jahren besonders von Gerdy und Mayor angewendet. Neuerdings sind sie wieder von v. Esmarch als Notverbände bei Verletzungen im Kriege empfohlen worden. In der Praxis werden sie gewöhnlich als Notverbände benützt, denn reine Taschentücher, Servietten u. s. w. finden sich ja wohl überall.

Zur Anlegung des Tücherverbandes benützt man entweder ein dreieckiges oder ein viereckiges Tuch, oder man legt das Tuch in Form einer Kravatte zusammen.

Das dreieckige Tuch hat die Form eines rechtwinkligen Dreieckes. Den rechten Winkel des dreieckigen Tuches bezeichnet man wohl auch als Spitze des Tuches; die beiden spitzen Winkel heissen die Zipfel.

Das viereckige Tuch ist gewöhnlich quadratisch von

1 Meter Seitenlänge. Halbiert man ein viereckiges Tuch von dieser Grösse in der Diagonale, so erhält man die übliche Grösse des dreieckigen Tuches.

Als Tücherstoff kommt die Leinwand, Baumwolle oder wohl auch schwarze Seide in Verwendung. Das von v. Esmarch empfohlene dreieckige Tuch, auf welchem die wichtigsten Tücherverbände, die mit demselben hergestellt werden, abgebildet sind, besteht aus Kalikot.

Tücherverbände für die obere Extremität.

Tuchverband für das Handgelenk. (Tafel 67.)

Das dreieckige Tuch wird zu einer Kravatte zusammengefaltet. Die Mitte der Kravatte wird quer über die Hohlhand an der Fingerbasis angelegt. Die beiden Enden werden dann über dem Handrücken gekreuzt, nach dem Handgelenk hin, und um dieses herumgeführt. Auf der Dorsalfläche des Handgelenkes werden die Enden geknotet.

Tuchverband für die ganze Hand. (Tafel 68 & 69.)

Das dreieckige Tuch wird auf einer horizontalen Unterlage flach ausgebreitet. Nun legt man die Hand mit der Volarseite so auf das Tuch, dass das untere Ende des Vorderarmes auf der Mitte der langen Seite des Tuches ruht, während die Fingerspitzen nach dem rechten Winkel hinausschauen. Dieser wird nun über die Rückenfläche der Hand bis zum Vorderarm zurückgeschlagen; nun werden die beiden langen Zipfel über dem Handrücken gekreuzt nach dem Handgelenk hin und mit einer Kreistour um dieses herumgeführt und auf der Dorsalseite des Handgelenks geknotet. (Tafel 69.)

Tuchverband für das Ellbogengelenk. (Tafel 70.)

Man legt die Mitte des kravattenförmig zusammengefalteten dreieckigen Tuches auf die Streckseite des Ellbogengelenkes, führt die beiden Zipfel analog den Touren einer Testudo cubiti reversa um den Vorderarm herum und knotet sie schliesslich in der Ellenbeuge.

Tafel 67.

Tuchverband für das Handgelenk.

Tafel 68.

Tuchverband für die ganze Hand. (Anfang des Verbandes.)

Tafel 69.

Tuchverband für die ganze Hand.
(Fertiger Verband.)

Tafel 70.

Tuchverband für das Ellenbogengelenk.

Tafel 71.

Tuchverband für das Schultergelenk.

Tafel 72.

Mitella triangularis als Stütze für den Oberarm.



Tuchverband für das Schultergelenk. (Tafel 71.)

Zur Deckung des Schultergürtels verwendet man am besten zwei dreieckige Tücher. Das erste dreieckige Tuch legt man so auf die Schulter, dass sein rechter Winkel gegen den Hals hinschaut, während die lange Seite auf die Aussenseite des Oberarmes zu liegen kommt. Den Rand dieser langen Seite faltet man nun ein- bis zweimal ein, bis man ungefähr in der Höhe des Deltoidesansatzes angekommen ist. Dann schlingt man die beiden Zipfel kreisförmig um den Arm herum und knotet sie auf der Aussenseite desselben.

Das zweite Tuch wird in Kravattenform zusammengelegt mit seiner Mitte in die gesunde Achselhöhle gelegt. Die beiden Zipfel werden dann auf die kranke Schulter geführt und hier über dem rechten Winkel des ersten Tuches geknotet. Schliesslich wird der rechte Winkel des ersten Tuches über den Knoten umgeschlagen und mit einer Sicherheitsnadel fixiert.

Mitella triangularis als Stütze des Oberarmes. (Taf. 72.)

Der Tuchverband, den man wohl überhaupt am häufigsten anwendet, ist das dreieckige Stütztuch für den Oberarm, die Mitella triangularis. Man fasst, um die Mitella für den rechten Arm anzulegen, den einen Zipfel des dreieckigen Tuches in die rechte Hand, den rechten Winkel desselben in die linke Hand und legt alsdann das Tuch so an den Thorax an, dass der Zipfel auf die rechte Schulter, der rechte Winkel in die rechte Achselhöhle zu liegen kommt. Nun legt man den rechtwinklig gebogenen Arm so auf das Tuch auf, dass die Hand auf die Mitte der Basis, der Ellbogen unterhalb des rechten Winkels zu liegen kommt. Jetzt wird der am Leib herabhängende Zipfel zur linken Schulter geführt und im Nacken mit dem über die Schulter geführten Zipfel geknotet.

Liegt der Vorderarm gut im Tuche, so wird jetzt der hinter dem Ellbogen vorstehende rechte Winkel des Tuches nach Glättung der Falten nach vorn umgeschlagen und hier mit einer Sicherheitsnadel befestigt.

Mitella triangularis zur Hebung des Oberarmes. (Taf. 73.)

Soll die Mitella nicht nur die Schwere des Armes übernehmen, soll sie vielmehr zugleich den Ellbogen heben, so gibt man dem Arm zweckmässig die Velpeau'sche Lage, lässt also den Patienten die Hand der kranken Seite auf die gesunde Schulter legen.

Der Verband wird dann in folgender Weise angelegt: Man legt das dreieckige Tuch so an der Vorderseite des Thorax an, dass die lange Seite an die Aussenfläche des Brustkorbes auf der kranken Seite, die Mitte der langen Seite etwa in die Höhe der Achselhöhle und der obere Zipfel auf die kranke Schulter zu liegen kommt. Der rechte Winkel schaut nach der gesunden Seite hin. Nun gibt man dem Arm über dem Tuche die Velpeau'sche Lage, schlägt den am Leib herunterhängenden Zipfel über das Ellbogengelenk in die Höhe, führt es über die gesunde Schulter hinweg und knotet ihn im Nacken mit dem über die kranke Schulter geführten Zipfel. Jetzt schlägt man den rechten Winkel um den Vorderarm in die Höhe und fixiert ihn mit der Sicherheitsnadel an dem um die kranke Schulter herumgeführten Zipfel.

Moore'scher Verband. (Tafel 74.)

Ein sehr zweckmässiger Ersatz der Mitella, namentlich für Schlüsselbeinbrüche ist von Moore angegeben worden. Ein entsprechend langes und etwa handbreites, kravattenförmiges Tuch wird gerade auf seiner Hälfte unter dem Ellbogen angelegt und von da über dessen Rückseite nach vorn über die Schulter geführt. Das andere Ende verläuft über die Vorderfläche des Ellbogens und über den Rücken, um hier mit dem von der anderen Schulter zurückkommenden Anteil vereinigt zu werden. Eine Mitella parva hält den Vorderarm leicht in die Höhe.

Mitella parva. (Tafel 75.)

Unter Mitella parva, oder auch Tragbinde für den kranken Arm genannt, versteht man ein kravattenförmig zusammengefaltetes Tuch, das, im Nacken geknotet,

Tafel 73.

Mitella triangularis zur Hebung des Oberarmes.

Tafel 74.

Moore'scher Verband.

Tafel 75.

Mitella parva und Tuchverband für die Wange und das Ohr.

Tafel 76.

Mitella quadrangularis.



Tafel 77.

Tuchverband für das Fussgelenk.

Tafel 78.

Tuchverband für den ganzen Fuss.

Tafel 79.

Tuchverband für das Knie.



den rechtwinklig gebeugten und am Leib anliegenden Vorderarm an seinem unteren Ende schlingenförmig umgreift.

Mitella quadrangularis. (Tafel 76.)

Mit dem viereckigen Tuch wird die Mitella in der Weise angelegt, dass man bei herabhängendem Tuch die beiden oberen Zipfel desselben in beide Hände fasst, sich an die kranke Seite des Patienten stellt, die Mitte des zu obigen Zipfeln gehörigen Randes in die Achselhöhle der kranken Seite legt und die Zipfel dann auf der gesunden Schulter knotet. Nun legt man den rechtwinklig gebeugten Vorderarm an den Leib heran, ergreift die beiden herabhängenden Tuchzipfel und führt sie ebenfalls nach der gesunden Seite hin, wo auch sie geknotet werden. Auf dem Rücken des Patienten ist dann eine leere Tuchtasche entstanden; diese führt man um die kranke Schulter herum nach vorn auf die Brust und steckt sie hier fest.

Tücherverbände für die untere Extremität.

Tuchverband für das Fussgelenk. (Tafel 77.)

Die Mitte des kravattenförmig zusammengelegten Tuches wird quer über die Mitte der Fusssohle gelegt, die beiden Enden werden nun sich kreuzend über den Fussrücken nach dem unteren Ende des Unterschenkels hingeführt, wo sie nach einer Kreistour über den Knöcheln auf der Vorderfläche des Beines geknotet werden.

Tuchverband für den ganzen Fuss. (Tafel 78.)

Man stellt den Patienten so auf das dreieckige Tuch, dass der rechte Winkel des Tuches nach vorne und die Basis des Tuches hinten an die Achillessehne zu liegen kommt. Nun schlägt man den rechten Winkel über die Zehen hinweg über den Fussrücken bis zum Unterschenkel zurück, fasst die beiden Zipfel, kreuzt sie über dem Fussrücken, führt sie zum Unterschenkel hin, bildet über den Malleolen eine Kreistour und knotet auf der Vorderfläche des Beines.

Tuchverband für das Knie. (Tafel 79.)

Man legt die Mitte des kravattenförmig zusammengelegten Tuches auf die Patella, führt die beiden Zipfel entsprechend den Touren einer Testudo genu reversa um Oberschenkel und Unterschenkel herum und endet in der Kniekehle.

Tuchverband für die Hüfte. (Tafel 80.)

Der Tuchverband für die Hüfte wird am besten mit zwei Tüchern hergestellt. Das erste Tuch legt man so auf die Aussenseite des Oberschenkels, dass der rechte Winkel nach dem Rumpf, die lange Seite nach dem Fuss hin schaut. Nun faltet man zwei- bis dreimal diesen unteren Rand, führt die beiden Zipfel kreisförmig um den Oberschenkel herum, um sie an der Aussenseite desselben zu knoten.

Das zweite kravattenförmig zusammengelegte Tuch wird etwas unterhalb der Höhe des Nabels kreisförmig so um den Unterleib herumgeführt, dass es den rechten Winkel des ersten Tuches kreuzt. Nun wird dieser rechte Winkel nach unten heruntergeschlagen und mit einer Sicherheitsnadel befestigt.

Tücherverbände für den Kopf.

Capitium triangulare. (Tafel 81.)

Das Capitium triangulare dient zur Deckung der Schädelwölbung. Man legt die Mitte der langen Seite des dreieckigen Tuches auf der Stirn dicht über den Augenbrauen an, während der rechte Winkel am Nacken herunterhängt. Nun fasst man die beiden Zipfel der langen Seite und führt sie dicht über den Ohren vorbei über den Hinterkopf herum, wo sie den rechten Winkel decken, und zur Stirn hin, auf deren Mitte sie geknotet werden. Nun schlägt man den rechten Winkel gegen den Scheitel hin in die Höhe und befestigt ihn hier mit einer Nadel an die Haube.

Man kann das Capitium triangulare auch umgekehrt herstellen, indem man die Mitte der langen Seite des

Tafel 80.

Tuchverband für die Hüfte.

Tafel 81.

Capitium triangulare.

Tafel 82.

Capitium quadrangulare.

Tafel 83.

Tuchverband für die Stirn und den Hals.

Tuches am Hinterhaupt unter der Protuberantia occipitalis, während man den rechten Winkel über die Stirne hinaufschlägt und am Scheitel feststeckt.

Capitium quadrangulare. (Tafel 82.)

Das Capitium quadrangulare deckt ausser der Schädelwölbung auch noch die Ohren und einen Teil der Wangen. Man schlägt sich das viereckige Tuch derart um, dass ein grösseres und ein kleineres Viereck entsteht und der Rand des grösseren Vierecks den des kleineren um etwa Handbreite überragt. Das so zusammengelegte Tuch wird nun derart auf die Wölbung des Schädels gelegt, dass der obere, dem kleineren Viereck angehörige Rand gerade an die Augenbrauen heranreicht, während der untere, dem grösseren Viereck entsprechende, über das Gesicht herabhängt. Man fasst nun die beiden oberen Zipfel, welche dem kleineren Viereck angehören, und knotet die Zipfel über dem Kinn. Nun fasst man die beiden unteren, dem grösseren Viereck angehörigen Zipfel, zieht dieselben hervor, bis die Haube dem Kopf glatt anliegt und führt sie nach hinten zum Nacken hin, wo sie geknotet werden.

Tuchverband für die Stirn, Stirnband. (Tafel 83.)

Die Mitte der Kravatte wird auf die Stirn aufgelegt, die beiden Zipfel werden nach hinten gegen den Nacken, unterhalb der Protuberanz herum und nach der Stirn zurückgeführt, wo sie geknotet werden.

Tuchverband für das Auge. (Tafel 84.)

Man legt die Mitte des kravattenförmig zusammengelegten Tuches schräg auf die Glabella auf und führt den einen Zipfel über das zuzudeckende Auge, den anderen über die entgegengesetzte Kopfseite hin nach dem Nacken, wo man sie knotet.

Tuchverband für die Wange und das Ohr. (siehe Taf. 75.)

Die Mitte des zusammengefalteten Tuches wird unter das Kinn gelegt; die beiden Zipfel werden, der eine

vor dem Ohre, resp. über dem Ohre, der andere hinter dem Ohre nach dem Scheitel in die Höhe geführt und auf demselben geknotet.

Tuchverband für den Hals. (siehe Tafel 83.)

Man legt die Mitte des kravattenförmig zusammengelegten Tuches vorn auf die Mittellinie des Halses, führt die Zipfel kreisförmig um den Hals herum und knotet sie in der Mittellinie.

Tücherverbände für die Brust und den Rumpf.

Tuchverband für die Mamma. (siehe Tafel 84.)

Das kravattenförmig breit zusammengelegte Tuch wird mit seiner Mitte von unten her so unter die weibliche Brust gelegt, dass dieselbe in die Höhe gehoben wird. Die beiden Zipfel werden nun schräg nach der entgegengesetzten Schulter hin und mit einer Achtertour um dieselbe herumgeführt. Die Knotung geschieht auf der Höhe der Schulter.

Schamtuch. (Tafel 85 u. 86.)

Am Unterleib wird das dreieckige Tuch als Schamtuch verwendet. Die lange Seite umgreift das Becken von vorn her etwa in Nabelhöhe. Der rechte Winkel wird zwischen den Schenkeln durchgezogen: die drei Enden werden schliesslich an der Rückseite des Rumpfes geknotet. (Tafel 85.)

Eine andere Art des Schamverbandes wird mit zwei Tüchern ausgeführt. Das erste Tuch wird in Nabelhöhe nach Art eines Gürtels kreisförmig um den Leib gebunden. Das zweite Tuch wird breit kravattenförmig zusammengefaltet, zwischen den Beinen durchgezogen und vorne und rückwärts an der Kreistour geknotet oder mit Sicherheitsnadeln befestigt. (Tafel 86.)

B) Wundverbände.

1. Kleinere Wundverbände.

Wunden, bei denen man keinen regelrechten anlegen will, namentlich bei den alltäglich vor-

Tafel 84.

Tuchverband für das Auge und die weibliche Brust.

Tafel 85.

Tuchverband für den Unterleib.

Tafel 86.

Schamtuch.

Tafel 87.



Kleinere Wundverbände. (Neff'scher Fingerling aus Trikotgewebe.)

kommenden kleinen Verletzungen, kann man einen Schutz gegen Verunreinigung von aussen dadurch erzielen, dass man auf die Wunde englisches Pflaster oder Pflanzenpapier oder ein Stück Heftpflaster oder eine der von Unna angegebenen, mit Sublimat, Zinkoxyd, Jodoform, Salicylsäure präparierten Salben oder Pflastermulle auflegt.

Sollen diese Stoffe halten, so darf die Wunde nicht mehr bluten, und ihre Umgebung muss ganz trocken sein.

Trockenheit der Wundumgebung ist auch notwendig, wenn man die kleine Wunde mit Collodium decken will, welches man mit einem Pinsel oder etwas Watte aufträgt. Benützt man das Collodium an den Fingern, so darf man ja keine zirkuläre Bepinselung des Fingers vornehmen, da sonst leicht schwere Ernährungsstörungen infolge der Druckanämie entstehen können.

Der Jodoform-Gaze-Collodium-Verband wird in der Weise angelegt, dass man die Wunde mit einem entsprechend grossen Stück Jodoformgaze deckt, das man an den Rändern mit Collodium befestigt. Auf dieses erste Stückchen Jodoformgaze legt man dann noch eine Reihe anderer gleicher Schichten auf; nur die Ränder dürfen jeweils mit Collodium bestrichen werden. Die Mitte des Verbandes muss frei bleiben, um etwaige Sekrete durchzulassen.

Will man das Collodium von der Haut entfernen, so benützt man am besten dazu den Aether aceticus. Rückständige Heftpflastermassen entfernt man am besten durch Waschen mit Aether oder Terpentin.

Die Art der Anlegung der kleineren Verbände erhellt aus Tafel 87.

Wir sehen da die Fingerspitzen gedeckt durch einen Heftpflasterverband, einen Collodiumverband und einen Fingerling. An dem kleinen Finger sehen wir den Neff'schen Fingerling aus Trikotgewebe; auf dem Handrücken sehen wir das Heftpflaster in Sternform oder nach Art einer radiär eingeschnittenen Scheibe oder in Form des

sogen. Maltheserkreuzes, d. h. als ein Viereck mit eingekerbten Ecken aufgelegt.

Die Pasta cerata von Schleich wird hergestellt aus geschmolzenem, gelben Bienenwachs, das mit Ammoniak behandelt wird, um die Verbindung des Wachses mit $1^1/_2$—2 Volumen Wasser zu ermöglichen. Dadurch entsteht ein crêmeartiger, neutraler Salbenkörper, der kein Ammoniak mehr enthalten darf. Diese Paste dient zur Wundbedeckung namentlich bei Verbrennungen. Verbindet man die Paste mit Pepton, Zinkoxyd, Gummi- und Amylum, so gibt sie ein ausgezeichnetes Verbandmittel, mit dem Gaze auf die Wunde festgeklebt werden kann, da die Paste in einigen Minuten fest antrocknet.

Ebenfalls von vorzüglicher Wirkung für gewisse Formen von Wunden ist der Unna'sche Zinkleimverband. Wir beschreiben die Anlegung des Verbandes zur Behandlung eines Unterschenkelgeschwürs. Nach gründlicher Abwaschung des ganzen Gliedes mit Seifenwasser wird dasselbe bis auf das Geschwür dick mit erwärmtem Zinkleim

Rp. Zinci oxydati
Gelatinae purissimae aa. 10.0
Glycerin
Aqu. destillatt. aa. 40.0
S. Erwärmt aufzupinseln.

bestrichen. Auf das Geschwür selbst wird Jodoform aufgepudert oder eine Lage von Watte oder mit Jodoform oder Sublimat imprägnierter Gaze gelegt. Hierauf wird eine gewöhnliche appretierte Mull- (Organtin-) Binde („blaue Binde") von zwei Seiten gleichmässig aufgerollt, in Wasser getaucht und in folgender Weise an dem eingeleimten und noch nicht ganz trockenen Unterschenkel angelegt: Der Patient sitzt mit erhobenem Bein dem Arzt gegenüber. Dieser fasst die beiden Köpfe der Binde, so dass die sie verbindende Brücke dem Geschwür gegenüber an die Hinterseite des Unterschenkels zu liegen kommt, wo sie sofort anklebt. Die beiden Köpfe werden so nach vorn geführt, dass sie sich über dem Ge-

schwüre und der dasselbe bedeckenden Watte kreuzen; die dabei aneinander vorüberstreichenden Hände wechseln die beiden Bindenköpfe unter sich aus, ziehen sie fest an, so dass der Unterschenkel hier an der Stelle des Geschwürs in seinem Umfange merklich verkleinert wird, und führen die Köpfe zu einer neuen Kreuzung nach hinten, die entweder ober- oder unterhalb des Geschwürs, jedenfalls aber diesem gegenüber stattfindet. Dabei tauschen beide Hände die Köpfe wieder aus, ziehen sie fest an und ziehen sie nach vorn, wo sie sich wieder an einer anderen Stelle kreuzen, und so fort, bis der ganze Unterschenkel, so weit er erkrankt und eingepinselt war, wenigstens aber vom Mittelfuss bis zur Wade, mit solchen gekreuzten Touren bedeckt ist. Sodann wird sofort eine zweite, feuchte appretierte Mullbinde über die erste gelegt, die den Zweck hat, etwaige dünn bedeckte Stellen der ersten Bindenlage auszufüllen und durch weitere spiralig auseinander gelegte Touren den ganzen Verband zu befestigen. Diese zweite Binde kann, ebenso wie die erste, zweiköpfig, oder auch, wie gewöhnlich, einköpfig angelegt werden. Unna legt sie meistens anfangs zweiköpfig an und wickelt allerdings häufig jeden Kopf für sich zu Ende. Nach kurzer Zeit ist der Verband vollständig erhärtet und bleibt 2—4 Tage, je nach der Menge des gelieferten Sekretes, liegen. Die Schmerzen beim Gehen und Niedersetzen des Fusses mindern sich unter ihm bald, so dass der Patient in seiner Beschäftigung nie unterbrochen wird. Je fester der Verband angelegt wird, um so besser erweist sich der Erfolg beim Abnehmen desselben; man legt den Verband immer so fest an, als der Patient es momentan eben gut ertragen kann. Wenn die Menge des Sekretes sich mindert, kann dieser Dauerverband später auch 8 Tage liegen. Die lege artis angelegte Binde, welche wegen der vorausgeschickten Einleimung des Unterschenkels überall fest anliegt, bedingt eine Dehnung der gesunden Haut nach der hiedurch verkleinerten Geschwürsfläche und verhindert deren Zurückziehen.

Der Zinkleimverband gestattet vollständig die Haut-

ausdünstung; ebenso ist er durchlässig für Wundsekrete. Er lässt sich durch einfaches Befeuchten mit warmem Wasser leicht entfernen.

Grössere Wundverbände.

Die heutige Generation von Studierenden sieht in den Kliniken in der Regel nur die aseptische Wundbehandlungsmethode und weiss nicht viel von den Wandlungen, welche die Lister'sche Methode in den letzten drei Dezennien durchgemacht hat. Es ist hier nicht der Platz, die Geschichte der gesamten Lister'schen Wundbehandlung in allen ihren Details zu entwickeln. Für uns handelt es sich nur darum, die Verbandmethoden zu beschreiben. Hier sind wir aber der Ansicht, dass der junge Arzt durchaus mit den Modifikationen vertraut sein soll, welche das Lister'sche Verbandverfahren seit seiner Einführung Ende der Sechziger Jahre erfahren hat.

Wir schildern daher zunächst den

Typischen Listerverband,

um dann die allmähliche Umgestaltung dieses Verbandverfahrens und die Gründe für die Umgestaltung desselben in seiner heutigen Form zu besprechen.

Bei dem typischen Listerverbande kam auf die Wunde selbst der Protektiv-Silk (Wachstaffet überzogen mit einem Gemisch von 1 Teil Dextrin, 2 Teilen Amylum, 16 Teilen 5% Carbollösung), auf diesen die feuchte, oder verlorene Compresse (feuchte Carbolgaze, 6 blättrig, etwas grösser als die Wunde); hierauf folgte die antiseptische Gaze in 7facher Schicht, die hergestellt wurde aus gebleichter oder ungebleichter Baumwollengaze in 6 Meter langen und 1 Meter breiten Stücken, die im kochenden Wasser 2—3 Stunden durchwärmt, ausgebreitet und mit folgender Mischung versetzt wurde: 1 Teil crystallisierte Carbolsäure, 5 Teile Harz, 7 Teile festes Paraffin; als vierte Schichte wurde der Makintosh, d. i. Baumwollstoff, der durch Kautschuk [illegible]durchlässig gemacht ist, (anstatt des Makintosh ist jetzt [illegible]h der Billrothbattist oder das Guttaperchapapier),

als fünfte Schicht die achte Lage Gaze aufgelegt. Auf diese Schicht kamen dann Binden aus Cambric, Callicot und Gaze, die mit 5% Carbolsäure getränkt waren, eventuell elastische Binden.

Der Lister'sche Verband wurde angelegt, während ein Hand- oder Dampfspray mit $2^1/_2$% Karbollösung über das ganze Verbandsgebiet hinweggeleitet wurde.

Diesen typischen Listerverband modificierte zunächst v. Volkmann in der Weise, dass er an den Rändern des Verbandes noch eine Schicht entfetteter Watte anlegte, um dem Eindringen von Mikroorganismen unter den Verband von den Rändern her entgegenzutreten. Lister selbst hatte diesen Abschluss des Verbandrandes dadurch herbeizuführen gesucht, dass er nach Abschluss des Verbandes noch eine elastische Binde um den Verbandrand herumführte.

Der weiterhin zunächst typische Lister'sche Verband wurde nun in der Weise geändert, dass man Ersatzmittel für die giftige, zur Imprägnation der Gaze dienende Karbolsäure suchte. So führte Ranke das Thymol, Maas die essigsaure Thonerde, von Mosetig das Jodoform, Kocher das Wismut, v. Bergmann das Sublimat, Maas das Sublimat-Kochsalz-Glyzerin, Sozin das Zinkoxyd, Lister selbst das Eucalyptusöl und das sog. Alembrotsalz (Chlorammonium 1.0, Sublimat 2.5) als Ersatz für die Karbolsäure ein, indem sie also z. B. Thymolgaze, Sublimatgaze, Jodoformgaze etc. empfahlen.

Praktische Bedeutung für die Wundbehandlung im allgemeinen haben nur die Sublimatgaze und Jodoformgaze behalten, die andern Gazesorten werden in der Regel nur in Ausnahmefällen verwertet.

In den allerletzten Jahren sind wieder eine ganze Reihe antiseptischer Mittel und bezüglicher Gazen empfohlen worden, ohne jedoch eine allgemeine Verbreitung gefunden zu haben, so die Kreolingaze, die Lysolgaze, die Dermatolgaze, Europhengaze, Loretingaze, Nosophengaze.

Eine weitere Modifikation des Listerverbandes bezog sich dann auf den Ersatz der Gaze selbst. So führte Lister selbst an Stelle der Gaze für gewisse Zwecke den Borlint ein, d. h. gewöhnlichen Lint, der in heisse, gesättigte Borsäurelösung getaucht und dann an der Luft getrocknet war. v. Bruns empfahl zuerst die entfettete Watte, Thiersch die Salicylwatte, v. Bergmann die Sublimatwatte, v. Mosengeil-Bardeleben Karboljute, Bardeleben Chlorzinkjute, Thiersch Salicyljute, Juillard desinfizierte mit 3% Karbollösung getränkte Badeschwämme, Morosow den Hanf oder Werg oder auch Oacum genannt, Weljaminoff den Theerwerg, Arton den Perubalsamwerg, Makuschina den Flachs, Kirchner Sublimatschnüre, Lukas-Championnière die Torfwatte, Esmarch-Neuber den Torfmull in Polstern, Schede die Glaswolle, Kümmell Steinkohlenasche in Kissen, Leisrink-Hagedorn das Waldmoos, Hagedorn Moospappe und Moosfilzplatten, P. Bruns Sublimatholzwolle, Romberg Cellulosenwatte, Kümmell Waldwolle und Holzcharpie, Escher Sägespähne, Port frische Holzspähne, Beduin ungeleimtes Filtrierpapier, Gedeke Sublimatpapier.

Der Verbandwechsel beim Listerverband wurde in der Regel schon am Tage nach der Operation nötig und geschah dann weiterhin alle 2—3 Tage.

Dieser häufige Verbandswechsel war nun ein entschiedener Nachteil des Lister'schen Verfahrens. Er widersprach vor allem dem auch von Lister mit Nachdruck betonten Grundsatz, dass die Wunde zu ihrer Heilung unbedingt Ruhe nötig habe. Diese Ruhe der Wunde zu beschaffen, war nun das Bestreben einer ganzen Reihe von Chirurgen, von den besonders Esmarch, Neuber und Maas genannt sein mögen. Diese Bestrebungen hatten die Einführung des Dauerverbandes im Gefolge.

Das Prinzip dieses Dauerverbandes ist das, dass der erste unmittelbar nach der Operation oder nach der Verletzung angelegte Verband bis zu vollendeten Wundheilung liegen bleiben soll. Die Wunde wurde mit Pro-

tectiv bedeckt, darüber der Verbandstoff gelegt und das Operationsgebiet möglichst fixiert und ruhig gelagert.

Als die Bedingungen für das Gelingen des Dauerverbandes waren tadellose Antiseptik, exakte Blutstillung, resorbierbares Ligatur- und Nähmaterial, Vermeidung von Höhlenbildungen in der Wunde, Sekretableitung ohne Drainageröhren und schliesslich Verbandstoffe notwendig, welche neben einem genügenden antiseptischen Depot eine möglichst hohe Aufsaugungsfähigkeit besassen.

Die ersten Dauerverbände, die man in dieser Weise ausführte, hatten einen grossen Erfolg. Es heilten nicht nur Operationswunden, sondern auch schwere Verletzungen, wie komplizierte Frakturen, unter einem einzigen Verband. Diese Art der Dauerverbände hatte aber noch einen grossen Nachteil. Man deckte sie nämlich noch mit einer Schicht von Guttaperchapapier und machte so eine Verdunstung des Wundsekretes unmöglich. Es entstanden daher unter den Verbänden sehr leicht Ekzeme, und ausserdem nahmen die lange liegenden Verbände, namentlich an den unteren Extremitäten, von den macerierten Hornschichten der Haut aus, bald einen penetranten Geruch an.

Diese beiden Nachteile vermied man nun bald, indem man das deckende Gummipapier, also den undurchlässigen Stoff, wegliess. Namentlich Neuber und Bruns lehrten, dass der Dauerverband zugleich ein Trockenverband sein müsse.

Ein weiterer Fortschritt, der erst in den letzten Jahren gemacht wurde, war der, dass man von dem antiseptischen Verfahren zu dem aseptischen überging.

Nachdem man gelernt hatte, dass die Wundheilung am besten dann vor sich geht, wenn man unter möglichster Ausschaltung der Luft- und namentlich der Kontaktinfektion operiert, sah man auch sofort ein, dass es unnötig ist, einen Verbandstoff mit einem antiseptischen Depot zu versehen.

Hat man wirklich aseptisch operiert, so braucht man die Wunde auch nicht mit antiseptischem Verband-

material zu decken, es genügt vielmehr auch eine aseptische Bedeckung, und diese hat dabei den Vorzug, dass sie jede Reizung der Wunde vermeidet.

Die Asepsis des Verbandmateriales kann man auf zweierlei Weise erreichen. Einmal kann man das Verbandmaterial kochen in einem gewöhnlichen Dampfkochtopf; dann hat man aber den Nachteil des schweren Trocknens; oder man benutzt zur Sterilisierung den strömenden Wasserdampf.

Um letztere Methode zu verwerten, braucht man eigene Apparate, sogenannte Sterilisatoren. Solcher Sterilisatoren sind eine ganze Anzahl konstruiert worden. Für die Praxis eignen sich ziemlich in gleicher Weise die Apparate von Schimmelbusch, v. Braatz, v. Kronacher, v. Rotter, Körte und anderen.

Diese Apparate gestatten, den Verbandstoff jedesmal vor der Operation oder vor der Anlegung des Verbandes keimfrei zu machen.

Um für die praktischen Aerzte keimfreies Verbandmaterial vorrätig zu haben, wird jetzt die Sterilisation von den Verbandstofffabriken besorgt. Die Verbandstoffe müssen zuerst in der Weise eingepackt werden, wie sie verschickt werden sollen, und müssen dann sterilisiert werden. Um dies erreichen zu können, hatte zuerst Gleich an der Billroth'schen Klinik empfohlen, die Verbandstoffe innerhalb verschlossener Pappschachteln in trockener Hitze zu sterilisieren. Turner und Crupin haben dann einen Apparat konstruiert, in dem es gelingt, die in den verschlossenen Pappschachteln befindlichen Verbandstoffe auch durch Wasserdampf zu sterilisieren. In neuerer Zeit hat endlich Dührssen „sterilisierte, antiseptische und aseptische Einzelverbände" in verschieden grossen Blechkapseln eingeführt; in diesen Blechkapseln befinden sich sterilisierte Jodoformkompressen, Watte und Binden. (Bezugsquelle Hirschapotheke in Würzburg.)

Das unstreitig beste Verbandmaterial zu aseptischen Zwecken ist die einfache Gaze (Mull).

Für grössere Betriebe ist die Gaze allein aber zu teuer; man legt dann auf die Wunde selbst nur eine dünnere Schicht Gaze auf und legt dann auf diese ein billigeres, aber auch gut aufsaugendes und leicht austrocknendes Verbandmaterial; namentlich haben sich die Moosverbände (Torfmoos, Mooswatte, Moospappe), ebenso auch die Holzwolle einer grösseren Verbreitung zu erfreuen.

Eine besondere Modifikation des aseptischen Verbandes erfordert die Heilung der Wunden unter dem feuchten Blutschorf nach Schede. Nach streng aseptischer Operation lässt man die Wundhöhle sich mit Blut anfüllen, legt auf dieselbe Protectiv und darüber einen aseptischen Verband. Der Protectiv soll eine Verdunstung des Blutes in der Wunde verhindern, dagegen soll der Verband nach Möglichkeit die Verdunstung und Eintrocknung des überflüssigen, in den Verband gedrungenen Blutes befördern. Die Methode eignet sich besonders für Operationen an den Knochen, namentlich für Necrotomieen.

Eine andere Modifikation des aseptischen Verbandverfahrens besteht in der von Kocher und v. Bergmann hauptsächlich empfohlenen Jodoformgazetamponade. Dieselbe wird in der Weise ausgeführt, dass man die Wunde mit Jodoformgaze ausstopft und diese Jodoformgaze dann nach mehr oder weniger langer Zeit, nach 3—8 Tagen, entfernt, um dann die Wunde etweder sekundär zu nähen oder von der Tiefe granulieren zu lassen. Die Methode kommt namentlich bei der Operation tuberkulöser Affektionen der Haut, Drüsen, Gelenke und Knochen, bei allen Operationen, bei denen keine exakte Blutstillung möglich ist, und bei den Operationen am Mund, an der Nase, am Mastdarm und Urogenitaltractus in Anwendung.

Anstatt der Jodoformgaze verwendet man heutzutage in all den Fällen, in denen man nicht gerade die Wirkung des Jodoform betonen will, zur Tamponade zweckmässiger sterile Gaze. Man hat dann nicht die Gefahr der Jodoformintoxikation.

Grössere Wundverbände an den einzelnen Körperteilen.

Nachdem wir nunmehr die Prinzipien kennen gelernt haben, nach denen die antiseptischen resp. aseptischen Verbände angelegt werden sollen, wollen wir beschreiben, in welcher Weise die Anlegung grösserer aseptischer Wundverbände an den einzelnen Körperteilen geschehen soll. Ein bestimmtes Schema kann man natürlich für diese Verbände nicht aufstellen; wer sich die einzelnen typischen Verbände gut zu eigen gemacht hat, wird auch diese grösseren Verbände ohne jede Schwierigkeit anlegen können, indem er sich die für den betreffenden Körperteil früher angegebenen Verbandtypen von Fall zu Fall variiert.

Allgemein gilt die Regel, auf die Wunde zunächst eine mehrfache Schicht steriler Gaze zu legen; behält man nun die Gaze als Verbandmaterial bei, so kommt auf diese erste Lage geschichteter Gaze, eine solche ungeschichteter, sogenannter gekrüllter Gaze. Auf diese Schicht kommt dann sterilisierte entfettete Watte, und das Ganze wird mit Mullbinden oder Cambricbinden eingewickelt. Befürchtet man ein leichtes Abrutschen dieser Binden, so kann man über diese noch eine gestärkte Binde, sog. blaue Binde herumwickeln.

Nimmt man statt der Gaze als weiteres Verbandmaterial z. B. Torfmoos, so wird über die Gazelage das Torfmooskissen gelegt und dieses dann eingewickelt. Diese Torfmooskissen haben den Vorteil, dass sie dem betreffenden Körperteil gleich eine gute Stütze gewähren.

Sehr wichtig ist für die Anlegung der grösseren Wundverbände die Regel, dass man den Patienten beim Anlegen des Verbandes die Lage einnehmen lässt, welche er nachher im Bette einnehmen soll. Befolgt man diese Regel nicht, so wird man sehr oft z. B. am Hals, an der Brust und am Bauch und ebenso an den Extremitäten Misserfolge erleben.

Tafel 88.

Wundverband am Kopf.

Tafel 89.

Wundverband am Hals.

Wundverband am Kopfe. (Tafel 88.)

Der Wundverband am Kopfe muss in der Weise angelegt werden, dass die ganze Schädelwölbung mit dem Verbandstoff gedeckt wird. Man macht einen solchen „Wickelverband“ für den Kopf in der Weise, dass man mit einer Zirkeltour um Hinterhaupt und Stirn beginnt und die einzelnen Bindentouren so verlaufen lässt, wie die Touren des Capistrum simplex oder duplex. Nach Fertigstellung des Verbandes bedecken die Bindengänge nicht nur die Schädelwölbung, sie gehen vielmehr auch vor beiden Ohren vorbei, unter dem Kinne her und um den Hals herum.

Wundverband am Hals. (Tafel 89.)

Nach Operationen oder Verletzungen am Hals polstert man den Verband gut, die Polsterung geht auch über die Brust herüber und unter die Achselhöhlen. Die den Verbandstoff befestigenden Bindentouren umfassen nicht nur den Hals, sondern umgeben auch nach unten hin, nach Art der Stella pectoris et dorsi beide Schultern, die Brust und den Rücken; nach oben hin umfassen sie, um ein Abrutschen des Verbandes nach unten hin zu verhüten, in einigen Kreistouren Stirn und Hinterhaupt. Zu diesen Kreistouren gelangt man, indem man die Binde vom Kinn aus vor dem Ohr in die Höhe auf den Scheitel und von hier aus um den Nacken unter die Protuberantia occipitalis externa hinführt.

Ehe man den Verband am Hals abschliesst, legt man vom Kinn an ein Stückchen Gummipapier auf die obersten Bindentouren auf und wickelt dasselbe mit der letzten Bindentour fest. Oben am Hals und am Kinn schlägt man dieses Gummipapier nach innen um und erreicht so einen Schutz des Verbandes gegen Beschmutzen beim Essen und Trinken.

Wundverband an der Brust. (Tafel 90.)

Nehmen wir als Beispiel die Anlegung eines Verbandes nach Mammaamputation, so deckt man die Wunde mit dem Verbandstoff, polstert die gesunde Brust und

Achselhöhle, die Ränder des Verbandes nach oben und unten hin ordentlich mit Watte, polstert ebenso die Achselhöhle der operierten Seite mit Krüllgaze oder Watte aus und führt ebenfalls etwas Watte um den oberen Teil des Oberarms dieser Seite herum. Nun wickelt man zunächst mit Kreistouren den Verband an den oberen Teil des Leibes und an die Brust an und wickelt dann mit Touren einer Spica humeri descendens die Schulter und den oberen Teil des Oberarms ein. Ist die Schulter zugedeckt, so macht man noch einige Kreistouren um die Brust und am unteren Ende des Verbandes. Um hier wirklich einen guten Abschluss zu erzielen, müssen die letzten Bindentouren angelegt werden, während die Patientin in horizontale Lage gebracht wird. Hat man die letzten Touren im Sitzen der Patientin angelegt, so werden dieselben unfehlbar vom Leibe abstehen, sobald die Patientin ins Bett gelegt wird.

Wundverbände am Unterleib. (Tafel 91 u. 92.)

Bis vor wenigen Jahren wurden nach Laparatomien ausserordentlich grosse Verbände um den Leib gewickelt. In neuerer Zeit ist dies nicht mehr üblich. Man legt eine dünne Schicht steriler Gaze auf die genähte Wunde auf und befestigt die Gaze an den Rändern entweder mit Heftpflasterstreifen oder durch Aufpinseln von Collodium. Will man aber eine Compression durch den Verband erzielen, so muss man den Verbandstoff mit Kreistouren befestigen und muss, damit diese nicht abrutschen, die Binden auch mehrmals nach Art einer Spica coxae descendens um beide Hüften und Oberschenkel herumführen.

Man erleichtert sich das Anlegen der Bindentouren wesentlich dadurch, dass man unter das Becken ein Stützbänkchen unterschiebt. Die am meisten zu empfehlenden Bänkchen sind die von v. Volkmann und Czerny. Das letztere hat den Vorteil, dass es sich leicht höher und tiefer stellen lässt. (Tafel 91.) Im Notfall können auch zwei kräftige Assistentenhände das Becken von unten her heben. Welches von diesen beiden Bänkchen man auch wählen mag, so muss man immer dafür sorgen,

Tafel 90.

Wundverband an der Brust.

Tafel 91.

a

b

Lagerung des Patienten zur Anlegung eines Wundverbandes am Unterleib.
Stützbänkchen von v. Volkmann (a) & Czerny (b).



Tafel 92.



Lagerung des Patienten auf der Ollier'schen Stütze.

Tafel 93.

Stützen des Beckens durch einen Assistenten von unten her.

dass die Schulterblattgegend und der Kopf durch entsprechend hohe Kissen ebenfalls gestützt werden. Um sowohl das Becken als den Rumpf gleichmässig stützen zu können, hat Ollier eine recht praktische Vorrichtung angegeben, die sich für jede Körpergrösse benützen lässt. Ihre Gestalt und Anwendungsweise erhellt wohl ohne weiteres aus der Abbildung. (Tafel 92.)

Wundverbände am Becken. (Tafel 93, 94 u. 95.)

Wundverbände am Becken sind sehr häufig anzulegen. Wir brauchen sie nach den Operationen am Hüftgelenk, nach Operationen in der Leistengegend, so nach Drüsenexstirpationen und Bruchoperationen, nach Operationen am Kreuzbein, an den Hoden u. s. w.

Um diese Verbände richtig anlegen zu können, ist zunächst eine richtige Lagerung der Patienten nötig. Man kann sich da auf verschiedene Weise helfen. Entweder hebt man das Becken in die Höhe, indem man, wie wir das eben beschrieben haben, ein v. Volkmann-sches oder Czerny'sches Bänkchen oder die Ollier'sche Stütze unterschiebt, oder man legt den Patienten quer so auf den Operationstisch, dass er auf diesem nur mit seinem Oberkörper aufliegt. Damit er dann nicht abrutscht, muss ihn ein Assistent halten und zwar in der Weise, dass er die Brust unterhalb der Achselhöhlen von beiden Seiten her mit den flach aufgelegten Händen und abducierten Daumen umgreift; ein zweiter Assistent muss die beiden abducirten Beine halten; schliesslich braucht man noch einen dritten Assistenten, der das Becken von unten her in die Höhe drücken muss. (Tafel 93.) Diese Lagerung ist namentlich bei erwachsenen Patienten unbequem und ermüdend für die Assistenten. Man hat deshalb besondere Lagerungsvorichtungen konstruiert, die es gestatten, den Patienten vom Operationstisch direkt ohne Assistentenhände in die Höhe zu heben und dies zugleich zu thun, ohne das Verbandgebiet selbst zu verdecken. Der bekannteste und empfehlenswerteste dieser Apparate ist der von Hase-Beck konstruierte Krankenheber. (Tafel 94.)

Wie die Abbildung zeigt, geschieht mittelst dieses

Apparates die Hebung des Patienten in der Weise, dass mehrere Zangen den Oberkörper des Patienten umgreifen, während das Becken selbst und die Extremitäten durch T förmige Unterstützungsvorrichtungen unterstützt werden. Mittelst einer Kurbel wird dann der Patient eleviert. Der Apparat leistet ganz ausgezeichnete Dienste, er belästigt den Patienten absolut nicht und gestattet dem Arzt ein ausserordentlich bequemes Hantieren.

Hat man nun auf die eine oder andere Weise das Becken hochgelagert, so dass man bequem herankommen kann, so werden in jedem Falle die beiden Spinae ilei anteriores superiores durch Auflegen einer dickeren Schicht Watte geschützt.

Ueber jeder Spina macht man in die Watteschicht ein Loch, so dass sich die Watte gut um die Spina herumlegen kann. Dann deckt man die Wunde mit der einfachen Gazeschicht, legt auf diese Krüllgaze auf und polstert schliesslich von der Mitte beider Oberschenkel an bis zum Nabel herauf die Weichteile mit Watte. Das Ganze wickelt man ein, indem man die Touren der Spica coxae duplex als Grundlage benutzt. Besondere Sorgfalt muss man darauf verwenden, dass der untere Rand des Verbandes an beiden Oberschenkeln, ferner der Rand des Verbandes an der Leiste gegen die Labien resp. das Scrotum und schliesslich der obere Rand des Verbandes am Leib gut anliegt. Gerade am Leib aber dürfen die Bindentouren nicht zu fest sitzen, sonst schneidet der Verband ein.

Besondere Sorgfalt muss man bei Kindern darauf verwenden, dass der Verband nicht alsbald von Urin und Kot durchtränkt wird. Den Abschluss gegen Urin erreicht man noch am besten in der Weise, dass man nach völliger Fertigstellung des Verbandes zwei breite Streifen Billrothbattist, entsprechend der Leistengegend [illegible]er inneren Seite des Oberschenkels mit Stärke[illegible]wickelt; der Billrothbattist muss beiderseits [illegible]e herunterreichen. Um den Kot vom Ver[illegible]n, lagere ich kleinere an der Hüfte oder

Hase-Beck'scher Krankenhebeapparat.

Tafel 95.

Lagerung des Kranken in einem Stehbett.



Tafel 96.

Wundverband am Damm.



an der Leiste operierte, mit dem eben geschilderten Wundverband versehene Kinder stets in ein Phelps'sches Stehbett. (Tafel 95.) Unter ein solches Bett kann die Bettschüssel einfach untergestellt werden, und der Kot vermag den hinteren Teil des Verbandes nicht zu beschmutzen.

Wundverband am Damm. (Tafel 96.)

Nach Operationen am Damm beim männlichen oder weiblichen Geschlecht liegt der Patient zur Anlegung des Verbandes gewöhnlich in Steinschnittlage, d. h. ein unter das Kreuzbein geschobenes Bänkchen erhöht das Becken, während gleichzeitig die beiden Beine von Assistenten im Hüftgelenk gebeugt gehalten werden. In der geschilderten Stellung wird nun zunächst auf die Wunde am Damm die Verbandgaze aufgelegt, beim männlichen Geschlecht muss man hierzu das Scrotum in die Höhe halten lassen. Auf die einfache Verbandschicht wird Krüllgaze aufgelegt und dann ordentlich mit Watte nachgepolstert. Die Wattepolsterung umfasst noch die oberen Drittel beider Oberschenkel und den unteren Teil des Rumpfes; eine besondere Polsterung umfasst ferner den Penis und das Scrotum, und zwar so, dass man die Watteschicht in der Mitte durchbohrt und den Penis hindurchsteckt. Nun wickelt man zuerst eine regelrechte Spica coxae duplex an, dann führt man die Binden in Achtertouren um beide Hüftgelenke herum, so dass eine Spica gerade in der Mitte des Dammes entsteht. Die Bindentouren verlaufen dabei von der Spina ilei ant. sup. der einen Seite unter dem Scrotum resp. der Vulva vorbei, um die Rückseite des anderen Schenkels herum zur Spina ilei ant. sup. dieser Seite hin und von da in gleicher Weise über den Damm und den anderen Oberschenkel herum, zu der Spina, von der man ausgegangen ist. Beim Anwickeln dieser letzteren Bindentouren muss man darauf achten, dass die Peniswurzel nicht eingeschnürt wird. Ist der Verband fertig, so sichert man die richtige Lage der Binden am Damm dadurch, dass man sie noch mit

einer Sicherheitsnadel aneinanderheftet. Endlich muss man darauf achten, dass die Analöffnung völlig frei ist; um eine Beschmutzung des Verbandes mit Kot zu verhüten, schiebt man unter die hinteren, sich kreuzenden Touren ein Stückchen Billrothbattist oder Guttaperchapapier ein und schlägt dieses nach oben um. Einen ebensolchen Schutz fügt man beim weiblichen Geschlecht am oberen Rande des Dammverbandes ein zur Verhütung einer Beschmutzung von der Scheide aus; bei Männern legt man den Billrothbattist so an, dass man ein Loch in den Battist schneidet und den Penis hindurchsteckt.

Wundverbände an den Extremitäten.

Das Anlegen von Wundverbänden an den Extremitäten bietet in der Regel keine Schwierigkeiten. Man gibt dem betreffenden Gliede die erwünschte Stellung, legt dann den Verband an und fixiert in der Regel das Glied in dieser Stellung noch durch einen Contentivverband.

c) Lagerungs-Verbände.

Die Lagerungsverbände gebraucht man teils zur Behandlung entzündeter, teils zur Behandlung verletzter Glieder; sie spielen namentlich in der Frakturbehandlung eine Rolle und haben hier im wesentlichen den Zweck, durch eine passende Lagerung des Gliedes der Zunahme der Dislokation der Frakturenden entgegenzuwirken, bis der definitive Frakturverband angelegt werden kann.

Man kann die einzelnen Lagerungsverbände in verschiedene Gruppen unterbringen.

Kissen.

Kissen sind in verschiedener Form in der alltäglichen Praxis in Gebrauch. Da haben wir zunächst die Spreukissen, die mit Haferspreu oder Hirse-
eu gefüllt werden, die Häcksel- und Sand-
n. Man füllt alle Kissen nur bis zur Hälfte an,

Tafel 97.

a) Desault'sches Kissen.
b & c) Strohmeyer'sches Kissen.
d) v. Dumreicher'sches Kissen.



Tafel 98.

Middeldorpf'scher Triangel.

so dass sie sich dem betreffenden Glied gut anschmiegen. Eine Abart der Sandkissen sind die Sandsäcke, die in der Regel in Wurstform vollgefüllt werden und als Stütze für eine operierte oder verletzte Extremität dienen.

Andere Kissen werden von vornherein in einer bestimmten Form angefertigt und mit Rosshaar oder Seegras gepolstert. Repräsentanten dieser Art von Kissen sind für die obere Extremität das Desault'sche Kissen, das Stromeyer'sche Kissen und der Middeldorpf'sche Triangel, für die untere Extremität das v. Dumreicher'sche Kissen.

Das Desault'sche Kissen haben wir bereits kennen gelernt anlässlich des Desault'schen Verbandes. (Taf. 97a.)

Das Stromeyer'sche Kissen (Tafel 97b) kommt zur Verwendung bei Verletzungen des Schultergürtels und des Oberarmes. Es ist ein dreieckiges, an den Spitzen abgestumpftes Kissen, das von der schmalen Basis an gegen die rechtwinklige Ecke an Dicke zunimmt. Die obere stumpfe Spitze kommt in die Achselhöhle: die untere an das Handgelenk, der rechte Winkel unter den kranken Ellbogen. Mittelst einer über die gesunde Schulter geführten Binde wird zunächst das Kissen an den Thorax befestigt, dann wird der rechtwinklig gebeugte Arm über dem Kissen an den Leib gelegt und über dem Kissen an den Thorax mittelst einer Binde angewickelt. Ueber das Ganze kommt dann noch eine gewöhnliche Mitella triangularis, und so hat der Arm wirklich eine gute Stütze.

Der Middeldorpf'sche Triangel (Tafel 98) dient zur Fixation des gebrochenen Oberarms in Abduktionsstellung. Ein dreieckiges Rosshaarkissen wird mit seiner Basis längs der Seitenfläche des Rumpfes angelegt; mittelst Gurten, die an den beiden Enden der Basis angenäht sind, wird das Kissen an den Thorax befestigt. Ueber die beiden kurzen Seiten des Dreiecks wird nun der Ober- und Unterarm so herübergelegt, dass das Ellbogengelenk gerade an den rechten Winkel zu liegen kommt. In dieser Stellung wird dann der Arm mit Binden an das Kissen

anbandagiert. Um ein Oedem des Armes zu ver
muss derselbe vor dem Anbandagieren auf das
von den Fingerspitzen bis zu der Schulter gut
wickelt werden. In neuerer Zeit benutzt man sta
Kissens lieber einen dreieckigen Rahmen. Dieser R
besteht aus drei, entweder aus Holz oder Draht
fertigten Hohlrinnen, die einerseits zur Aufnahm
Ober- und Vorderarm, andererseits zum Anlegen
Seitenfläche des Thorax bestimmt sind. Die Anwe
ist die gleiche wie beim Kissen. Der Rahmen wir
weder mit Gurten an den Thorax befestigt oder n
Binden angewickelt, der Arm in gleicher Weise
oben beschrieben, am Triangel anbandagiert.

Das v. Dumreicher'sche Kissen für die
Extremität ist ein keilförmiges Polsterkissen, bes
zur Aufnahme des Beines bei Oberschenkelbr
(Tafel 97 d.)

Laden. (Tafel 99.)

Unter Laden in ihrer einfachsten und ältesten
versteht man längliche, viereckige, oben offene K
welche die ganze Länge des Unterschenkels aufne
und gepolstert, dem Bein von der Ferse bis zum
eine bequeme Lagerung ermöglichen sollen. Fu
und Seitenteile der Laden lassen sich auf- und n
klappen. Die bekanntesten dieser Laden sind die
Heister'sche Lade und die Scheuer'sche Lade.

An der Petit-Heister'schen Lade ist
der eigentlichen Lade noch ein Gestell zum Hoch
Niederstellen derselben angebracht und ebenso noc
Unterlage für den Oberschenkel, die mit dem
der Lade einen stumpfen Winkel bildet. (Tafel 9

Recht einfach ist die Scheuer'sche Lade
ausser dem Fussbrett einfach aus drei Holzlatten b
die durch einfache Coulissenvorrichtungen befestig
mit Binden oder Riemen an den Unterschenkel
bunden werden. (Tafel 90 b.)

Tafel 99.



a) Petit-Heister'sche Lade.
b) Scheuer'sche Lade.

Tafel 100.



a) Einfach geneigte schiefe Ebene.
b) *Doppelt* geneigte schiefe Ebene.
c) *Fialla'sche* Stäbchenbeinlade.

Die einfach und doppelt geneigte schiefe Ebene (Taf. 100).

Die einfach geneigte schiefe Ebene (Planum inclinatum simplex) dient einfach zur Hochlagerung des Beines. Man stellt die erhöhte Lage am einfachsten durch Unterlegen von Kissen oder durch Unterschieben eines umgelegten Stuhles oder schiefes Einlegen eines Brettes her. (Tafel 100a.)

Die doppelt geneigte schiefe Ebene (Planum inclinatum duplex) hält zur Erschlaffung der Muskeln das Bein im Hüft- und Kniegelenk leicht gebeugt. Wenn der Oberschenkelteil des Apparates länger ist als der Oberschenkel des Patienten, so kann man mit dem Apparat gleichzeitig noch eine Extension am Oberschenkel ausüben, indem man die Schwere des Rumpfes nach unten zieht, während der Gegenzug am oberen Ende des Unterschenkels statt hat. Eine zweckmässige Ergänzung brachte v. Esmarch an der doppelt geneigten schiefen Ebene an, nämlich eine Reihe etwa fingerdicker Stäbe, die in Löcher des Apparatrandes in bestimmten Abständen eingesteckt werden. Dieselben dienen als gute Stütze für die Polsterung und das Glied und können nach Bedarf entfernt und wieder eingesetzt werden. (Taf. 100b.)

Aehnlich dieser v. Esmarch'schen Vorrichtung ist die Fialla'sche Stäbchen-Beinlade, die aus einer grösseren Anzahl etwa 30 cm langer Holzstäbchen und einer Eisenstange mit Schraubenmutter besteht. (Tafel 100c & d.) Die an einer durchbohrten Scheibe festsitzenden Stäbchen werden über den am unteren Ende mit einer Platte versehenen Eisenstab geschoben und können an demselben in jeder beliebigen Stellung durch die Schraube befestigt werden.

Reifenbahren. (Tafel 101 d.)

Reifenbahren sind aus Holz oder Metall oder aus einer Verbindung beider hergestellte Lagerungsapparate, welche dazu dienen, irgend einen Körperteil, in der Regel den Fuss, vor dem Drucke der Bettdecke zu bewahren. Sie können ferner Verwendung finden, um Eisbeutel an

ihnen anzubringen oder auch um das betreffende Glied an ihnen zu suspendieren. Die gewöhnliche Reifenbahre besteht aus Holzstäben, an denen in bestimmten Abständen Drahtreifen angebracht sind.

Stehbetten (siehe Tafel 95).

Die alte v. Renz'sche Spreizlade, welche das gebrochene Bein in Abduktionsstellung fixieren sollte, ist in neuerer Zeit zum Phelps'schen Stehbett ergänzt worden. Dieses Phelps'sche Stehbett, das in der Orthopädie häufig angewendet wird, benützen wir, wie schon früher erwähnt, zur Lagerung kleinerer, an der Hüfte operierter Kinder, ferner bei Oberschenkelbrüchen kleinerer Kinder. Man lässt nach der Form und Grösse des Körpers eine Holzlade anfertigen, deren Seitenhöhe etwas grösser ist als der sagittale Durchmesser des Rumpfes. Die beiden Beinladen befinden sich in mässiger Spreizstellung. An den Seitenwänden sind an den betreffenden Stellen Armausschnitte angebracht. Der Analgegend entsprechend befindet sich ein ovaler Ausschnitt an dem Rückenbrett. Die Polsterung geschieht mittelst eines gleichgeformten, in die Lade passenden Rosshaarkissens, dessen Analausschnitt man zum Schutze gegen Durchnässung und Beschmutzung mit Wachstaffet überzieht. Auf dieses Kissen legen wir noch in ganzer Ausdehnung der Lade eine dünne Schicht entfetteter Watte. Ist das Bett präpariert, so wird das Kind hineingelegt und mit Binden an dasselbe angewickelt. Um die Oberschenkel befestigt man zum Schutze gegen den Urin eine Schicht wasserdichten Stoffes (siehe Tafel 95).

Das Stehbett hat ausserordentlich grosse Vorteile. Man kann die Kinder in den Betten überall herumtragen; ausserdem ist die Defäkation und Urinentleerung ausserordentlich erleichtert, da man die Kinder im Verband nicht jedesmal in die Höhe zu heben und auf das Stechbecken zu setzen braucht, indem man dies letztere vielmehr einfach unter den Analausschnitt einschiebt.

Tafel 101.

Reifenbahre.

[redacted]

[redacted]

Tafel 102.

a



b



c

a) Mayor'sche Drahtrinne.
b) Roser'sche Drahtrinne.
c) Bonnet'sche Drahthose.

Drahtrinnen und Drahthosen (siehe Tafel 102).

Die Drahtrinnen wurden zuerst von Mayor empfohlen, dann namentlich von Roser und Bonnet vervollkommnet.

Die einfachen Mayor'schen Drahtrinnen bestehen aus einem Rahmen von stärkerem Drahte, welcher von einem Geflecht dünneren Drahtes übersponnen ist. Die Polsterung geschieht entweder mittelst einer dünnen Rosshaarmatratze oder mittelst Watte. (Tafel 102a.)

Roser stellte die Drahtrinnen für die untere Extremität aus drei Stücken her, um die Rinnen leichter jeder Extremitätengrösse anpassen zu können. (Taf. 102b.)

Durch Verbindung zweier Drahtrinnen und Hinzufügung eines Beckenteiles erzeugte Bonnet seine bekannte Drahthose zur bequemen Lagerung der Beine und des Beckens. (Tafel 102c.)

d) Contentivverbände.

Die Contentivverbände dienen zur Feststellung der verschiedensten Körperteile und finden eine ausserordentlich mannigfaltige Verwendung, da sie sowohl die Ruhigstellung entzündeter und operierter, als die Fixation gebrochener Glieder erstreben.

Man teilt die Contentivverbände ein in Schienenverbände, oder auch amovible Verbände, und in erhärtende oder auch inamovible Verbände genannt.

Für die Anwendung der Contentivverbände gelten folgende allgemeine Regeln:

1. Man soll in der Regel schon vor Anlegung des Verbandes dem Glied diejenige Stellung geben, in der man später den betreffenden Körperteil fixiert haben will.

2. Die Contentivverbände sind gut zu unterpolstern. Dabei sind besonders hervorragende Knochenpunkte zu berücksichtigen, damit über denselben kein Decubitus entsteht.

3. Der Verband darf nicht zu locker angelegt wer
weil er sonst seinen Zweck nicht erfüllt.

4. Der Verband darf aber auch nicht zu fest ang
werden. Namentlich darf an keiner Stelle eine zirk
Einschnürung des Gliedes durch eine zu fest angezo
Bindentour statthaben, weil es sonst leicht zu ven
Stauung, ja zu einer Gangrän des abgeschnürten T
kommen kann.

5. Der Contentivverband soll in der Regel
beiden Gelenke oberhalb und unterhalb des zu fixie
den Körperteils einschliessen, da bei Einbeziehung
eines Gelenkes eine völlige Ruhigstellung nicht zu
zielen ist.

6. Der Contentivverband muss sorgfältig überw
werden, damit er beim Eintreten ungünstiger Ersch
ungen sofort gewechselt werden kann. Solche
günstige Erscheinungen sind: Heftige und anhalt
Schmerzen, sowie Anschwellung und blaurote
färbung der freigelassenen peripheren Teile des Gli
Finger und Zehen.

A. Schienenverbände.

Schienen sind verschieden geformte, feste Verb
geräte, welche zur Stütze irgend eines Körperteiles die
und sich daneben noch zu mannigfachen anderen
richtungen verwenden lassen, wie zur passenden Lage
zur Suspension, zur Extension und Distraktion.

Wir besprechen die Schienen nach dem vers
denen Material, aus dem man sie herstellt.

Holzschienen.

Die Holzschienen werden aus den mannigfach
Holzarten hergestellt, namentlich aber aus dem
der Birke, Eiche, Linde, Tanne, des Nussbaum
des Ahorn.

Die gebräuchlichsten Holzschienen sind folge
ür die obere Extremität haben wir zunächst
denen Hand- und Vorderarmschienen; sie

Tafel 103.



a & b) Strohmeyer'sche Handbretter. c) Nélaton'sche Pistolenschiene.
d) Roser'sche Dorsalschiene für die Fractura radii loco classico.
e) Schede'sche Schiene
f) Carr'sche Schiene
g) Cooper'sche Schiene
h) Strohmeyer'sche Armschiene.

Tafel 104.



a) v. Volkmann'sche Suspensionsschiene.
b & c) Dupuytren'scher Verband bei Malleolarfraktur.
d) v. Volkmann'sche Knieschiene.

stehen aus leichtem Holz und werden mit Watte gepolstert.

Zur Ruhigstellung der Hand dienen die einfachen Handbretter (Stromeyer), die mit oder ohne besondere Schiene für den Daumen in Gebrauch sind. (Tafel 103a, b.)

Die Nélaton'sche Pistolenschiene dient zur Fixierung des gebrochenen Radius. (Tafel 103c.) Sie wurde verbessert von Schede in der Weise, dass sich die Hand auf der Schiene nicht nur in ulnarer Abduktion, sondern auch in volarer Flexion befindet. (Tafel 103e.)

Ebenfalls zur Behandlung der Radiusfraktur dienen: die Roser'sche Dorsalschiene, die Carr'sche Schiene und die Cooper'sche Schiene.

Zur Herstellung der Roser'schen Dorsalschiene wird eine gut gepolsterte, von dem Ellbogen bis zu den Fingerspitzen reichende Holzschiene von der Breite des Armes so auf die Dorsalfläche des Vorderarmes gelegt, dass die Hand in volarer Flexionsstellung herabhängt. Zwischen den Handrücken und die vorstehende Schiene wird nun Watte keilförmig eingesteckt. Dann wird die Schiene mittelst einer Binde am Arm befestigt, während die Finger frei bleiben. (Tafel 103d.)

Die Carr'sche Schiene ist eine leichte Holzschiene, auf die ein konvexes hölzernes Lager aufgeschraubt wird, das genau der konkaven Radiusfläche entsprechend, auf der radialen Seite etwas über 1 cm dick ist, nach der ulnaren Seite hin sich abflacht und vor dem Handgelenk ganz niedrig wird. Vorn ist schräg an die Schiene ein rundliches Querstück befestigt, das der Hand als Stützpunkt dient. Der Patient umgreift dieses Querstück, so dass der Daumen radial nach unten zu liegen kommt; dann wird die Schiene an den Vorderarm mit Binden angewickelt. (Tafel 103f.)

Die Cooper'schen Radiusschienen sind wellenförmig gestaltet. Sie besitzen eine Höhlung für das Handgelenk und einen leicht winklig abgebogenen Handteil.

Die Stromeyer'sche Armschiene dient zur Aufnahme des ganzen Armes. Sie ist flach ausgehöhlt, stumpfwinklig abgebogen und mit einem Loch für die Condylen des Oberarms versehen. (Tafel 103h.)

Die v. Volkmann'sche Flügel- oder Suspensionschiene besitzt am Oberarmteil einen Flügel, d. h. ein senkrecht stehendes ausgehöhltes Brett, das in einer Rinne verschiebbar ist und beliebig für beide Arme eingestellt werden kann; ausserdem ist am unteren Ende der Schiene ein Ring angeschraubt, mittelst dessen man die Schiene mitsamt dem anbandagierten Arm an einem über dem Bett angebrachten Galgen aufhängen kann. (Tafel 104 a.)

Für die untere Extremität stehen zur Zeit noch folgende Holzschienen in Gebrauch: Die Dupuytrensche Schiene dient zur Fixation des Fusses in Adduktionsstellung beim Knöchelbruch; eine flache, vom Knie bis zu den ausgestreckten Zehen reichende Schiene wird auf der Innenfläche des Unterschenkels auf ein dickes Kissen gelegt; das Kissen reicht nach oben bis zum oberen Ende des Schienbeins, nach unten bis zum inneren Knöchel. Es wird nun zunächst der Unterschenkel über dem Kissen an die Schiene anbandagiert, dann wird der Fuss durch mehrere Achtertouren in starker Adduktionsstellung gegen die Schiene hin befestigt. (Tafel 104 b & c.)

Die v. Volkmann'sche Knieschiene ist eine kurze, ausgehöhlte Schiene, die bei Einwicklung des Kniegelenkes in die Kniekehle gelegt wird, um einen Druck der Bindentouren auf die Gefässe und Nerven der Kniekehle zu verhindern. (Tafel 104 d.)

Die Watson'sche Schiene, die zur Fixation des resecierten Kniegelenkes angegeben wurde, hat an ihrem unteren Ende die Form eines Stiefelknechtes; über die beiden Gabeln wird ein Fussbrett gestellt. Der Teil der Schiene, der unter das Kniegelenk zu liegen kommt, ist abgerundet. (Tafel 105 a.)

Die v. Volkmann'sche Dorsalschiene für den Unterschenkel ist eine flache Hohlrinne, die an ihrem unteren Ende stumpfwinklig abgebogen ist. Die

Tafel 105.



a) Watson'sche Schiene.
b & c) v. Volkmann's Dorsalschiene für den Unterschenkel.
d) König's Abduktionsschiene für das Hüftgelenk.

Tafel 106.



a) Middeldopf's Pressschienenverband. b) Gooch'sche Spaltschienen. c) Schnyder's Tuchschienen. d) v. Esmarch's schneidbarer Schienenstoff.



Schiene wird auf den Fussrücken und die Vorderfläche des Unterschenkels aufgelegt; sie trägt an ihrer oberen Fläche mehrere Ringe, mittelst deren man den Unterschenkel bequem suspendieren kann. (Tafel 105 b & c.)

Die König'sche Abduktionsschiene für das Hüftgelenk ist eine entsprechend dem Hüftgelenk stumpfwinklig abgebogene, leicht ausgehöhlte Holzschiene mit einem kleineren aufsteigenden Teile für die Brust und einem grösseren absteigenden Teile für den Ober- und Unterschenkel. (Tafel 105 d.)

Die bisher genannten Holzschienen bestehen aus starrem Material. Es gibt nun auch eine ganze Reihe formbarer, biegsamer Holzschienen.

Die besten derselben, die eine weitere Verbreitung verdienen, sind die Mitteldorpf'schen Ahornfourniere. Es sind das breite, aus Ahornholz ganz dünn ausgeschnittene Schienen, die, in warmem Wasser angefeuchtet, sich den Körperformen leicht anschmiegen. Sehr gut eignen sich die Ahornfourniere zum Verband des gebrochenen Vorderarms. Je ein Fournierblatt, etwas breiter als der betreffende Vorderarm, wird in heisses Wasser getaucht, wohlgepolstert auf die Volar- und Dorsalseite des Vorderarms aufgelegt und mittelst Heftpflasterstreifen an diesem befestigt. Sie lassen die Frakturstelle dem Auge frei zugänglich und können leicht loser und fester angezogen werden. (Tafel 106 a.)

Weniger Eingang in die Praxis haben die Gooch'schen Spaltschienen, die Schnyder'schen Tuchschienen und der v. Esmarch'sche schneidbare Schienenstoff gefunden.

Die Gooch'schen Spaltschienen bestehen aus dünnen Fichtenbrettern, welche durch seichte, nicht ganz durchdringende Einschnitte in 1 cm breite, parallele Streifen geschnitten und auf Leder oder Leinwand geklebt werden. Sie lassen sich einem Arm oder Bein leicht anschmiegen und sind dabei doch völlig fest. (Tafel 106 b.)

Die Schnyder'schen Tuchschienen bestehen aus Nussbaumfournieren, welche dicht neben einander zwischen zwei Stücken Leinwand eingenäht sind. (Tafel 106 c.)

Der v. Esmarch'sche schneidbare Schienenstoff besteht aus zwei Lagen Leinwand, zwischen denen Tapetenspanstreifen in Abständen von je 5 mm neben einander gelegt und mit Wasserglas, Kleister oder Leim fest verklebt sind. Der Schienenstoff lässt sich leicht herstellen und bequem mit der Schere schneiden. (Tafel 106 d.)

Versieht man einfache Holzschienen an einem Ende mit Blechhülsen, so kann man sich durch Zusammenstecken solcher Schienen beliebig lange Schienen z. B. für eine Extremität herstellen (v. Esmarch).

Die Holzschienen lassen sich bequem durch eiserne Bügel mit einander vereinigen. Die bekannteste der so entstehenden Schienen ist die v. Volkmann'sche Supinationsschiene. An derselben ist der Handteil rechtwinklig zur übrigen Schiene angebracht, so dass die Hand mit dem Vorderarm in Supinationsstellung zu stehen kommt. (Tafel 107 a.)

Recht praktisch sind ferner die v. Esmarch'schen Resektionsschienen für die Hand. (Tafel 107 b & c.)

Auch durch Scharniere lassen sich die einzelnen Holzschienen mit einander verbinden. Als Beispiel diene die Sharp'sche Beinschiene, die an der Seite des Beines angelegt, vom Becken bis sum Mittelfuss reicht, am Knie das Charnier trägt und für den Knöchel einen Ausschnitt hat. (Tafel 107 d.)

Pappschienen.

Die Pappschienen werden am besten aus der gewöhnlichen grauen, überall käuflichen Buchbinderpappe hergestellt. Man taucht die Pappe vor dem Gebrauch in warmes Wasser; dann schmiegt sie sich den Körperkontouren sehr gut an und behält nach 12—24 Stunden, trocken geworden, ihre Form bei.

Zur Herstellung der Pappschienen schneidet man sich mit dem Messer die gewünschte Breite und annähernde Form her, schneidet aber die Pappe nicht ganz durch, sondern reisst sie in dem vorgezeichneten Schnitt

Tafel 107.



a) v. Volkmann's Supinationsschiene.
b & c) v. Esmarch's Resektionsschiene für die Hand.
d) Sharp'sche Beinschiene.

Tafel 108.



a'

b'

Pappwatte-Verband Linhart's.
a) Verband für den gebrochenen Oberarm.
b) „ „ „ „ Vorderarm.

ab. So erhält man einen dünnen, sich dem Körper leicht anschmiegenden Rand der Schiene. Schneidet man dagegen die Pappe sofort ganz durch, so werden die Ränder hart und scharf und erzeugen leicht Decubitus

Pappschienen kann man an allen Körperteilen anlegen. Sehr gut eignet sich z. B. ein Pappverband zur Fixation des in der Mitte gebrochenen Oberarms. Man schneidet, resp. reisst sich einen breiten Pappstreifen von doppelter Breite des Armes aus gewöhnlicher Pappe. der von der Wurzel des Halses über die Schulter hinweg bis zur Hand reicht. Am oberen Ende dieses Streifens macht man in gleichen Abständen vier Längseinschnitte, dem Ellbogen entsprechend aber einen runden Ausschnitt für den Epicondylus externus. Nun taucht man den Streifen in heisses Wasser, legt ihn an der Aussenseite des rechtwinklig gebeugten Armes an, biegt die oberen Längsstreifen kappenförmig um die Schulter herum und wickelt nun die Pappe mit einer Binde, von dem Handgelenk an, am ganzen Arm an. Spicatouren um die Schulter geben der Schiene oben deren richtigen Halt. Schliesslich wickelt man dann noch den ganzen Arm an den Thorax an.

Eine grosse Rolle spielte in früheren Jahren der auch heutzutage noch praktisch wohl verwertbare Pappwatteverband Linharts. Aus gewöhnlicher Pappe schneidet man sich für die verschiedenen Körperteile passende Modelle, welche gut mit Watte gepolstert und dann an dem betreffenden Gliede angewickelt werden. Wir bilden die betreffenden Modelle für die Frakturen des Oberarmes und des unteren Radiusendes ab. Die Flügelfortsätze an dem letzteren Modell werden so aufgebogen, dass ein Teil an die Volar- und ein Teil an die Dorsalseite der Hand zu liegen kommt. (Tafel 108.)

Recht zweckmässig ist die Modifikation des Pappverbandes für den Bruch des Oberarms, wie ihn Urban angegeben hat und beschreibt. Der Verband ist ein Pappschienenverband mit verstärkenden Schusterlatten. Sämtliche Pappstücke werden vor der Anlegung in warmem Wasser gut aufgeweicht. Der Verband wird

in folgender Weise ausgeführt: Man bindet ein Stück dicke Pappe (Tafel 109 a), welches von der Achselhöhle bis annähernd zur Darmbeingrube reicht, nachdem man es gut gepolstert hat, fest an dem seitlichen Teile des Rumpfes an. Die Breite entspricht mindestens der Entfernung der vorderen von der hinteren Axillarlinie. Diese Tafel soll als Stütze dienen für den Arm, der später, wenn er für sich geschient ist, noch an den Thorax befestigt wird, so dass dieser in letzter Linie als Schiene betrachtet werden kann. Die Schienung des Oberarmes selbst vollzieht sich in folgender Weise: Man braucht vier Pappschienen, eine für die Innenfläche, eine für die Aussenfläche, eine für die Vorderfläche und eine für die Hinterfläche. Die für die Aussenfläche berechnete reicht vom seitlichen Drittel des Schlüsselbeines bis zur Hand; es ist eine rechtwinklige Schiene. (Tafel 109 b.) Der rechte Winkel entspricht dem Ellbogengelenk. Diese Schiene hat neben der Fixierung der Bruchenden den Zweck, das Ellbogengelenk im rechten Winkel und den Vorderarm ruhig zu stellen. Die vorderen, inneren und hinteren Pappschienen entsprechen der Länge der Vorder-, Innen- und Hinterfläche des Oberarmes (Tafel 109 c). Auf diese Pappschienen kommen je nach ihrer Dicke vier bis acht verstärkende Schusterlatten. Die Pappschienen sind selbstverständlich sorgfältig gepolstert. Die Bruchstücke werden durch Extension und Gegenextension in ihrer normalen Lage erhalten und nun der ganze Schienenapparat von der Hand bis über das Schultergelenk hinauf durch eine sorgfältige Bindeneinwickelung befestigt. Man hat besonders darauf zu achten, dass sich die Pappschienen überall genau anschmiegen, damit man eine Art Kapselverband erhält. In dieser Verfassung (Tafel 109 d) wird der Arm an den Thorax angelegt und durch Bindengänge nach Art der zweiten und dritten Tour der Bindengänge des Desault'schen Verbandes an den Brustkorb befestigt (Tafel 109 e). Bei der Anlegung der Pappschienen ist im onderen darauf zu achten, dass sie weder in der Ellnbeuge noch in der Achselhöhle einen Druck aus-

Tafel 109.



Pappverband für den gebrochenen Oberarm nach Urban.

üben; man mache sie an diesen Stellen lieber 1 cm zu kurz, weil ihr Druck auf die Nerven und Gefässe einen so unheilvollen Einfluss ausüben kann, dass sich die Entfernung und Erneuerung des Verbandes nötig macht. Ist der Verband sorgfältig angelegt, so kann er ohne weiteres vier Wochen liegen bleiben, andernfalls wird er nach 14 Tagen gewechselt und erneuert. (Tafel 109.)

Neuerdings hat Solowjow angelegentlichst Pappe-Gipsschienen empfohlen. Die Schienen werden aus Pappe in beliebiger Form ausgeschnitten und mit Gipsbinden umwickelt. So lassen sich auch gut Lagerungsapparate für den Fuss und Unterschenkel nach Art der v. Volkmann'schen Schiene herstellen. Die Schienen können im Dampf sterilisiert werden und werden danach wieder hart.

Blechschienen.

Blechschienen kommen teils als im voraus geformte, teils als biegsame, formbare Schienen in Verwendung.

Die geformten Schienen aus verzinntem Eisenblech oder Zinkplatte haben meist die Form von Rinnen oder Kapseln für die obere und untere Extremität.

Die weiteste Verbreitung hat wohl die v. Volkmann'sche T-Schiene gefunden, die zur ruhigen Lagerung des Unterschenkels dient. Sie besteht aus der Rinne für den Unterschenkel, dem Fussbrett und dem T-Eisen. Entsprechend der Ferse ist in der Rinne eine Oeffnung angebracht. Das T-Eisen verhütet das Umfallen des Fusses nach der einen oder anderen Seite. Beim Anlegen der Schiene muss man darauf achten, dass der Fuss genau rechtwinklig steht, dass er dem Fussbrett durchaus anliegt und dass er fest an dieses anbandagiert wird. Die Ferse muss vor jedem Druck bewahrt sein, da sonst leicht Decubitus an derselben entsteht. (Tafel 110 a.)

Eine sehr zweckmässige Modifikation der v. Volkmann'schen Schiene ist neuerdings von P. Bruns an-

gegeben werden. Die Bruns'sche Schiene ist eine flache, rinnenförmige Schiene aus verzinntem Eisenblech, welches dem Rosten widersteht. Die Rinne ist aus zwei Hälften zusammengesetzt, welche sich übereinanderschieben lassen, um sie beliebig vergrössern resp. verkleinern zu können. Damit kein Druck der Ferse entstehen kann, ist an der Schiene über den Zehen ein einfacher Querbalken angebracht, an welchem sich der Fuss mittels einer Heftpflasterschlinge leicht suspendieren lässt. Anstatt des v. Volkmann'schen T sind an der Schiene zwei zusammenklappbare, unten halbkreisförmig abgebogene Arme aus Rundeisen angebracht, welche auf der Unterlage keine so starke Reibung verursachen. (Tafel 110 b.)

Formbare Blechschienen stellt man meist aus Weissblech her. Am bekanntesten sind die formbaren Schienen von Salomon und Schoen.

Die Salomon'schen Schienen sind platte Schienen aus dünnem Weissblech, welche je 35 cm lang und 10 cm breit, an einem Ende mit zwei dreigeteilten Fortsätzen und an dem andern mit zwei Spalten versehen sind, durch welche jene Fortsätze gesteckt und durch Umbiegen festgemacht werden können, so dass sich auf diese Weise leicht und schnell Schienen von beliebiger Länge herstellen lassen. (Tafel 110 c.)

Schoen schneidet aus dünnem Zinkblech mit der Schere Schienen, die beliebig formbar sind. Mehrere solche Schienen können in beliebigem Winkel zu einander gebogen, durch Zinkbrücken in Verbindung erhalten, durch Scharniergelenke untereinander beweglich verbunden werden. Die Zinkplatten können, um die Ausdünstung der Hand besser zu erlauben, mit Löchern versehen werden. Wir bilden ein Schoen'sches Modell zum Ausschneiden einer Armrinne und einer Unterschenkelrinne ab. Die Armrinne wird so geformt, dass man sie in der Längsachse zur Halbrinne biegt und in der Querachse an den eingekerbten Stellen senkrecht aufstellt. Durch die an den Kerben befindlichen Löcher werden Bindfaden gezogen, durch deren Zusammen-

Tafel 110.



a) v. Volkmann'sche ⊤-Schiene. b) Bruns'sche Schienenfläche.
c) Salomon'sche Schiene. d) Schön'sche Schiene für den Arm.
e) *Schön'sche Schiene* für den Unterschenkel.

Tafel III.



a) Raoult-Deslongchamps'sche Schiene. b & c) Kupferblechschienen nach Lee und Wilson. d & e) Englische Eisenblechschienen nach Koch. f & g) Port'sche Blechstreifenschienen.

knüpfen die gewünschte Winkelstellung erzielt wird. (Tafel 110 d.)

Nach Ausschneiden des Unterschenkelmodelles wird wieder aus dem Längsteil eine Rinne gebildet, während der unten zweifach eingeschnittene Teil senkrecht aufgestellt wird zur Bildung eines Fussbrettes. (Taf. 110 e.)

Besondere Modelle für die Anfertigung von Zinkschienen hat Raoult-Deslongchamps angegeben. Wir bilden das Modell der Schiene für den Bruch des Unterschenkels ab und zeigen, wie die nach dem Modell ausgeschnittene Schiene den Unterschenkel gut fixiert. (Tafel 111 a.) Sehr elegant, brauchbar, aber auch sehr teuer sind die aus vernickeltem Kupferblech hergestellten Schienen von Lee und Wilson. Die Figuren (Taf. 111 b & c) zeigen die Schienen für die Hand und den Vorderarm.

Ausserordentlich praktisch sind die englischen Eisenblechschienen für die Extremitäten. Dieselben stellen verschieden lange, etwa 2 Finger breite Schienen dar, die mit Watte gepolstert und dann mit Leinwand überzogen werden. Sie lassen sich leicht biegen, besitzen aber doch eine ziemlich grosse Festigkeit. Ich benutze sie fast täglich, ebenso werden sie von Helferich sehr empfohlen. (Fabrikant: Instrumentenmacher Stöpler in Greifswald.) Dr. Koch in Neuffen hat sie in letzter Zeit elastisch aus Nickelblech herzustellen gelehrt. Auch hat er mehrere Schienen durch Scharniere verbunden, so dass man dadurch eine Biegung der Schienen über die Kante erzielen kann. (Tafel 111 d & e.)

Durch Verbindung mehrerer Blechstreifen unter einander erhält man die namentlich im Kriege sehr verwendbaren Port'schen Blechstreifenschienen, die in grosser Anzahl leicht transportiert werden können Für alle Extremitätenfrakturen reichen drei dieser Schienen. aus, eine für die obere Extremität und je eine für den Ober- und Unterschenkel. Die Abbildungen zeigen die Schienen abgeflacht, wie sie transportiert werden, und zurecht gebogen, wie sie in Verwendung kommen. (Tafel 111 f & g.)

Bleischienen.

Verbandschienen aus Walzblei, welches mit Stoff überzogen wird, sind von Kohlmetz empfohlen worden. Die Schienen eignen sich für Schutzverbände aller Art, besonders aber für Notverbände. Ihre Handhabung ist eine sehr einfache. Beim Gebrauch schneidet man sich mit einer Schere oder mit einem Messer ein passendes Stück zurecht, drückt dasselbe dem verletzten Körperteil an und hält es durch ein oder zwei Tücher in seiner Lage fest. Die Schiene schmiegt sich fest an und kann auch wohl in geeigneten Fällen als Dauerverband verwertet werden. Man muss dann das Glied gut polstern und die Schiene mit Binden gut anwickeln. Es können auch gut Fenster in die Schiene geschnitten werden, so dass sie sich auch bei komplizierten Brüchen verwerten lässt.

Aluminiumschienen.

Die in der letzten Zeit von Steudel empfohlenen Aluminiumschienen eignen sich in erster Linie zu unterbrochenen Verbänden bei komplizierten Knochenbrüchen, besonders wenn dabei grössere äussere Wunden vorhanden sind, oder die Wunden bereits verunreinigt sind, und zur Nachbehandlung von Operationen an Gelenken. Auch zur Verstärkung erstarrender Verbände und zu Improvisationen verschiedener Art sind sie mit Vorteil zu verwenden. (Tafel 112.)

Vermittelst einer Universalzange können die Aluminiumschienen mit Leichtigkeit zu dem gewünschten Zweck vorbereitet werden. Es gehört zur Handhabung dieser Zange wenig mechanisches Geschick, wohl aber etwas rasch zu erlernende Uebung. Mit den unterhalb des Drehpunktes der Zange angebrachten messerförmigen Schneiden werden die Schienen in beliebiger Länge abgeschnitten. Bei den schmäleren gelingt dies ohne weiteres, bei den breiteren genügt es, in derselben Weise eine Einkerbung zu machen, und bricht man alsdann, durch einige kurze Biegungen, die Schiene an dieser Stelle ab.

Tafel 112.

Steudel's Aluminiumschienen.



In der eigentlichen Zange sind kleine Haken angebracht, diese pressen sich in die flach eingelegten Schienen derart ein, dass 4 Zähnepaare am Rande der Schienen sich aufstellen. Die oberen Zacken sind für die breiteren, die unteren für die schmalen Schienen. Die Zähne greifen in die umgelegten Binden so fest ein, dass ein Verschieben und Loswerden der Schienen verhindert wird, sie machen es möglich, mit den so vorbereiteten Aluminiumschienen ohne Gips mit einfachen steifen Gazebinden unterbrochene Verbände anzulegen; solche Verbände haben den Vorzug grosser Leichtigkeit.

Die an der Zange angebrachten schlitzförmigen Oeffnungen dienen dazu, die Schienen zu biegen und zwar die horizontal stehende Oeffnung zum Biegen über die Fläche, die senkrecht stehenden Oeffnungen zum Biegen der Schienen über die hohe Kante. Letztere Biegung, welche wichtig ist, um die Schienen an rechtwinklig gestellten Gliedmassen, z. B. dem Ellbogen nicht nur an der Beuge- und Streckseite, sondern auch an der Aussen- und Innenseite anlegen zu können, gelingt bei einiger Uebung an den schmalen Schienen leicht bis zu einem rechten, ja spitzen Winkel, bei den breiten ist etwas mehr Kraft und Geschicklichkeit dazu notwendig. Es ist bei diesen zweckmässig, die Biegung auf mehrere Stellen einer kurzen Strecke zu verteilen, so dass nicht eine scharf rechtwinkelige, sondern eine viertelkreisförmige Biegung resultiert. Endlich können mit der in 2 Hälften auseinander genommenen Zange die Schienen spiralförmig gebogen werden; eine Biegung, welche dann notwendig wird, wenn die Schienen nicht genau in der Längsaxe eines Gliedes, sondern zu dieser etwas schräg verlaufend angelegt werden sollen. Sollten bei den Biegungen Ausbiegungen nach der Seite zu stande kommen, wie solche bei den mit der Zange nicht Geübten zuweilen entstehen, so lassen sich diese leicht mit der als Hammer benutzten Zange wieder gerade klopfen.

Es ist zweckmässig, mehr als 2 Schienen an einem Gliede zu verwenden, da bei 3 oder 4 Schienen der Verbiegung nach irgend einer Richtung stets mindestens

eine Schiene mit ihrem starken Breitendurchmesser widerstrebt. Es wird dadurch trotz der Biegsamkeit der Aluminiumschienen eine grosse Festigkeit erreicht und die Uebersichtlichkeit über den in den Fenstern freiliegenden Körperteil bei der Schmalheit der Schienen in keiner Weise gestört. Bügel anzubringen ist bei Aluminiumschienen überflüssig, da Aluminium ein ungiftiges und leicht aseptisch zu machendes Material ist und deshalb ohne Bedenken in den Wundverband mit eingeschlossen werden kann.

Die Aluminiumschienen und die dazu gehörige Universalzange werden von der „Deutschen Metallpatronenfabrik in Karlsruhe-Baden" hergestellt.

Unbrauchbar gewordene Schienen und Abfall werden von der Fabrik zurückgenommen und dafür dem Aluminiumwerte entsprechend neue Schienen geliefert. Der Wert des alten Aluminiums entspricht etwa $^1/_3$ des Preises für neue Schienen.

Aluminiumblechschienen.

Eine aus Aluminiumblech hergestellte praktische Schiene hat Hansmann in der Form einer Universal-Vorderarm-Extensions- und Lagerungsschiene angegeben. Dieselbe kann bestens empfohlen werden. Diese Schiene besteht, wie auf Figur 1 (Tafel 113) dargestellt ist, aus zwei von Aluminiumblech hergestellten, bei a a mit excentrisch gerichteter Axe gelenkig verbundenen Lagerungsteilen für Vorder- und Oberarm, denen durch Feststellen dieses Gelenkes jede beliebige Stellung gegeben werden kann. An dem Vorderarmteil ist bei b ein abnehmbarer Bügel angebracht, welcher wiederum bei c c um eine horizontale Axe drehbar und feststellbar ist. An dem vorderen Teil des Bügels befindet sich ein bei d um seine Mitte drehbarer kleiner Stab, der fixiert ist, und an dessen beiden Enden Haken mit Flügelschrauben durchgeführt werden können.

Ausserdem ist dem Apparat ein Handbrett beigegeben, welches statt des Extensionsbügels eingefügt wird, wenn die Schiene nur als Lagerungsschiene dienen soll.

Tafel 113.



Fig. 1. Aluminiumblechschiene nach Hausmann.
„ 2. Lagerung derselben.
„ 3. „ „ in Dorsalflexionsstellung.
„ 4. „ „ in Ulnarflexionsstellung.

Die Lagerung der Schienen gestaltet sich nun äusserst einfach.

Zunächst wird ein mit 2 Haken (siehe Figur 2, Tafel 113) versehenes Handbrettchen, welches von den Metacarpophalangealgelenken bis zum Handgelenk reicht, mit Heftpflasterstreifen und einer darüber gelegten Mullbinde in der Vola der Hand befestigt und dann die Schiene nach genügender Unterpolsterung angelegt.

Bei der für sehr viele Fälle anwendbaren, auf Fig. 2 (Tafel 113) zur Anschauung gebrachten Lagerung in halber Supinationsstellung ist die Wirkung der Schiene ohne weiteres ersichtlich.

Während der Vorder- und Oberarm in beiden Lagerungsteilen durch Binden fixiert sind, wird die Extension an dem Handbrettchen bewirkt, von welchem aus zwei Gummizüge in die Haken der Schrauben eingehängt sind. Durch Andrehen der Schrauben kann nun eine beliebig starke Extension bewirkt werden und es ist durch diese Einrichtung eine stete Regulierbarkeit, die man einigermassen verständigen Patienten nach genügender Anleitung selbst überlassen kann, gegeben. Ausser dieser Extension in halber Supinationsstellung ist natürlich die volle Supinations- und Pronationsextension und durch Heben oder Senken des Extensionsbügels die Ab- resp. die Aduktionsstellung in ebenso einfacher Weise auszuführen. Ein Bild von der besonders zur Mobilisierung des Handgelenkes bei Steifigkeiten, wie sie nicht selten nach Frakturen und Gelenkerkrankungen vorkommen, sehr schnell zum erwünschten Ziele führenden Anwendungsweise gibt die Fig. 3 (Tafel 113), die die Wirkung der Schiene z. B. in starker Dorsalflexionsstellung darstellt.

Fig. 4 (Tafel 113) veranschaulicht den Gebrauch bei gewünschter Ulnarflexion in halber Supinationsstellung.

Drahtschienen.

Wir haben schon früher die Mayor-Bonnet-Roser'schen Drahtschienen kennen gelernt. Den Vorteil der Leichtigkeit und Reinlichkeit, der diesen

Lagerungsapparaten zukommt, zeigen auch alle anderen Drahtschienen. Die unzweifelhaft besten Schienen dieser Art, die sich einer weiten Verbreitung erfreuen; sind die Cramer'schen Drahtschienen.

Diese Cramer'schen Drahtschienen stellen flache, aus verzinntem Eisendraht hergestellte Hohlrinnen dar. Zwischen zwei festeren, längsverlaufenden Drähten sind jeweils dünnere, parallellaufende feinere Drähte wie Leitersprossen ausgespannt. Die Schienen lassen sich leicht über die Fläche und Kante biegen und daher bequem allen Körperteilen anpassen. Sie lassen sich ferner leicht verlängern, indem man einfach zwei Stücke dadurch verbindet, dass man einige Querdrähte ausschneidet und diese zum Zusammenbinden benutzt. Indem man ferner mehrere Querstäbe ausschneidet, lassen sich leicht gefensterte Schienen jeglicher Art darstellen. (Tafel 114 a, b, c, d.)

Als Beispiel der grossen Verwendbarkeit der Schienen wollen wir den Helferich'schen Verband für die Oberarmfraktur beschreiben. Helferich erreicht eine permanente Extension in der Längsrichtung des Oberarms dadurch, dass er die Drahtschiene dem rechtwinklig gebeugten Arm entsprechend biegt und sie dann am Arm fixiert. Das obere Ende der Schiene ist so gebogen, dass es der Schulter nicht dicht anliegt. Wird nun ein durch Watteeinlage gut präparierter Bindenzügel in die Achselhöhle gelegt, und an dem vorragenden Schienenende unter mässigem Zug befestigt, so ist ein permanenter Zug geschaffen, der durch erneutes Knüpfen der Achselbinde leicht reguliert werden kann. (Tafel 114 e.)

Von den sonstigen Drahtschienen hat sich die Smith'sche vordere Drahtschiene in den letzten Kriegen vielfach bewährt. Sie wird aus Telegraphendraht hergestellt, entsprechend dem Fuss-, Knie- und Hüftgelenk gebogen und angewickelt. (Tafel 114 f.)

Mittelst zweier Drähte, die man an den Halbringen anbringt, lässt sich auch leicht eine Suspension der Extremität ausüben.

Recht brauchbar sind ferner die Schienen aus Drahtsiebstoff (v. Esmarch). Mehrere Stücke des Draht-

Tafel 114.



a, b, c, d) Cramer'sche Schienen. e) Anwendung der Cramer'schen Schiene beim Oberarmbruch nach Helferich. f) Smith's vordere Drahtschiene.

gewebes werden, durch Schnüre mit einander verbunden, an die Extremität angewickelt. (Tafel 115 a.)

Port hat gelehrt, diesen Drahtsiebstoff in schmale Streifen zu zerschneiden und die einzelnen Streifen wie Binden aufzurollen. So entstehen die Port'schen Drahtrollbinden, die dann am besten halten, wenn sie spiralig um das Glied herumgewickelt werden, während man zwischen die einzelnen Touren noch etwas Gipsbrei einstreicht. (Tafel 115 c.)

Für eine Reihe von Verletzungen und Gelenkerkrankungen hat in neuester Zeit Heusner Spiraldrahtschienen empfohlen. Starker englischer Draht verläuft bei diesen Schienen in stumpfen Zickzackwindungen. Der Draht wird auf einer Filzplatte angenäht. Die Schienen lassen sich in jeder beliebigen Form über die Fläche und Kante biegen und streben gebogen kräftig in die alte Form zurück. Sie verbinden also Stützkraft mit Federkraft. Wir bilden die für die Radiusfraktur angegebene Schiene für sich und angelegt ab. (Tafel 115 b.)

Plastische Schienen.

Als Material zur Herstellung plastischer Schienen dienen: die Pappe, der Filz und die Guttapercha.

Plastische Pappschienen erhält man nach P. Bruns, wenn man gewöhnliche Pappe mit einer starken Schellacklösung durchtränkt. Die so hergestellte Pappe erweicht beim Erwärmen, wird dann am Körper geformt und behält nach dem Erkalten die neue Form bei. Diese plastischen Pappschienen werden in neuester Zeit als besondere Spezialität von Dr. Koch in Neuffen (Württemberg) ausgeführt. Koch hat die meisten der gebräuchlichen Schienen aus plastischer Pappe herzustellen gelehrt. Er hat die Schienen ausserdem, wenn nötig auch noch durch Aufnieten von Blechstreifen haltbar gemacht und hat schliesslich durch Anbringen von Scharnier- und Kugelgelenken sogenannte Universalschienen konstruiert. Die Figuren zeigen die Mannigfaltigkeit, mit der sich diese Schienen verwenden lassen (Tafel 116.) Wir bilden ab: die Schienen für den gebrochenen Vorder-

arm von Cooper und Schede, die Lagerungsvorrichtung für den Vorderarm und den Ersatz der v. Volkmann'schen Blechschiene. Die Koch'schen Schienen haben den Vorteil, dass sie bequem anzulegen und dass sie sehr leicht und gut zu reinigen und zu sterilisieren und dabei sehr billig sind.

Filzschienen.

Zur Herstellung von Filzschienen benützt man nach Angabe von Bruns gewöhnlichen Sohlen- oder Einlagefilz von 6—8 mm Dicke, den man mit einer konzentrierten alkoholischen Schellacklösung (660.0:1 Liter) in der Weise tränkt, dass man auf beiden Seiten der Platte portionsweise von der Lösung aufgiesst und dieselbe mit einem groben Pinsel verstreicht, bis sich die Poren des Filzes gleichmässig vollgesaugt haben. Kleinere Filzstücke taucht man einfach in die Lösung ein. Soll der Filz eine aussergewöhnliche Stärke haben, so wird er, nachdem sich ein Teil des Alkohols verflüchtigt hat, noch einmal getränkt.

Der auf diese Weise plastisch gemachte Filz trocknet bei höherer Temperatur innerhalb einiger Stunden. Ehe die Platte völlig erstarrt ist, glättet man sie durch Ueberfahren mit einem heissen Bügeleisen. Die Platte ist nach dem Erstarren bretthart. Erwärmt man sie aber auf 70° R., so wird sie weich und biegsam, lässt sich dann in jede beliebige Form bringen und behält diese Form auch nach dem Wiedererstarren bei. Das Erwärmen geschieht auf trockenem oder feuchtem Wege dadurch, dass man die Platte in einen warmen Ofen hält, mit einem heissen Eisen überstreicht oder in heisses Wasser taucht. Beim Gebrauch wird die erweichte Schiene auf die durch eine Unterlage gegen die Einwirkung der Hitze geschützte Haut gelegt, dem Gliede genau angepasst und durch eine Binde befestigt. Das Erstarren erfolgt innerhalb weniger Minuten; man muss sich daher mit dem Anlegen und Formen der Schiene beeilen.

Diese Filzschienen hatten den Nachteil, dass sie am Körper warm wurden und dann nachgaben. Anders

Tafel 115.

a) v. Esmarch's Drahtsiebstoff. c) Port's Drahtrollbinden.
b) Heusner's Spiraldrahtschiene.

Tafel 116.



Plastische Pappschienen nach Koch (a, b, c, d). Filzschienen nach P. Bruns-Anders (e). f) Pappschiene für den gebrochenen Oberarm.

hat neuerdings gezeigt, dass dies lediglich an der Art des verwendeten Filzes liegt. Er empfiehlt, einen Filz zu gebrauchen, der aus Hasenfellen mit Zusatz von Kaninchenhaar fabriziert wird, und diesem Filz durch Walken ein recht starkes Gefüge zu verleihen. Die Plastizität erhält der Filz durch Tränkung mit einer syrupdicken Lösung von Gummi lacca in tabulis mit Spirit. vini rectificat. Diese Lösung wird erst mit etwas Spiritus verdünnt, dann reichlich auf die Innenfläche des Filzobjektes gegossen und mit der Hand verstrichen, bis sie auf der anderen Seite gleichmässig durchgedrungen ist.

Die in Vorrat gehaltene dicke Lacklösung muss vor dem Tränken des Filzes einmal erwärmt gewesen sein, dann wird der Filz nachher haltbarer.

Ist der Filz bis zum Ueberfliessen getränkt, so wird er auf das Gipsmodell gespannt, dem er sich gut anschmiegt, wenn man ihn mit Binden gleichmässig anwickelt. Das Trocknen muss bei gewöhnlicher Zimmertemperatur geschehen. Nach 48 Stunden lässt sich der geformte Filz in der Regel leicht vom Modell abnehmen und entsprechend herrichten. (Tafel 116.)

Guttaperchaschienen.

Die Guttapercha ist ein eingedickter Pflanzensaft, der bei gewöhnlicher Temperatur fest ist, im Wasser von 50—60 Grad Wärme jedoch weich und biegsam wird, so dass man ihm jede beliebige Form geben kann. Diese Form behält er dann nach dem Erkalten bei.

Zum Gebrauch beim Verbande schneidet man aus den Guttaperchatafeln ein ausreichend grosses Stück heraus, wobei man berücksichtigen muss, dass das Material in heissem Wasser in der Länge und Breite ab-, in der Dicke aber zunimmt. Man bringt dann das ausgeschnittene Stück in warmes Wasser, legt es aus diesem, sobald es genügend erweicht ist, unmittelbar auf die vorher mit Oel bestrichene Haut auf und passt es durch Drücken und Streichen überall genau den Körperformen an. Die so gewonnene Form erhält man bis zum Erhärten der Schiene, das nach kurzer Zeit erfolgt, dadurch, dass man

die fertig angepresste Schiene mit einer feuchten Binde an das Glied anwickelt. Das Erweichen der Guttapercha erfordert eine gewisse Uebung. Erweicht man sie zu wenig, so formt sie sich nicht gut, erweicht man sie aber zu sehr, so klebt sie überall fest.

Fiberschienen.

Als ein „ideales" Schienenmaterial hat kürzlich Wiener das Fiber empfohlen. Fiber wird von den Elektrikern schon lange zum Isolieren benutzt und findet, wasserdicht gemacht, immer mehr Verwendung im täglichen Leben als Tischplatten, Waschgefässe, Eimer etc. Die Herstellung ist ein Geheimnis; es kommt in verschieden mächtigen Platten, von der Dicke des Pergamentpapiers bis zu 1 cm Dicke in den Handel. Die Farbe ist meist indianerrot, doch wird es auf besondere Bestellung auch schwarz und grau geliefert. Das beste und für chirurgische Zwecke einzig brauchbare Produkt liefert die Kartavest Fibra Company unter dem Namen „Flexible Sheel Kartavest". Wiener empfiehlt als zur Herstellung von Schienen geeignetstes Material die 1,5 mm dicken Platten.

Das Fiber, das im gewöhnlichen Zustand schwer zu schneiden und elastisch biegsam ist, wird in heissem Wasser plastisch und weich, wobei es etwas aufquillt und sich wie ein Stück weiches Leder handhaben lässt. Um rasch zum Ziele zu kommen, oder um es aseptisch zu machen, kann es ohne Schaden in siedendes Wasser gebracht werden. In diesem schmiegsamen Zustand um eine Extremität, z. B. den Oberschenkel gelegt und mit Binden befestigt, gibt es getrocknet und dann abgenommen, die Konfiguration des Gliedes wieder. Eine Fiberhülse kann mit der Schere leicht verengt werden, oder wenn das nicht angeht, so kann man eine den veränderten Verhältnissen angepasste neue Kapsel mit wenig Mühe in den Stahlrahmen einpassen.

Das Trocknen der Kapsel am Körper hatte viel Unbequemes für den Patienten und hat sich auch in der Folge als unnötig herausgestellt. Ein genaues Papiermuster wird hergestellt, danach wird die Fiberkapsel

Tafel 117.

Gipsbindenverband.

aus den Platten herausgeschnitten, aufgeweicht und dann so gerollt, dass sich die beiden freien Längskanten berühren. So wird die Hülse getrocknet, nachdem noch der obere und untere Rand nach aussen umgebogen ist, um ein Scheuern der Haut zu verhindern. Dann wird die Hülse in den Stahlrahmen mit Kupferstiften eingenietet. Die Bohrlöcher im Fiber reissen niemals aus. Polsterung mit weichem Flanell, ein Anstrich mit Schellack und das Anbringen von Schnürhaken vollenden das Werk.

Hornhautschienen.

Als gutes Material zur Herstellung von Hülsen und Schienen empfiehlt Vulpius die Hornhaut. Dieselbe besteht aus vorher enthaarter und getrockneter Tierhaut, die mit einem Lack imprägniert und gebrauchsfertig von der Firma Wagner in Düsseldorf bezogen wird. Sie wird nach einem Papiermuster zurechtgeschnitten, 12—15 Stunden in kaltes Wasser gelegt, dann auf dem Modell gewalkt. Das völlige Trocknen erfordert etwa 6 Tage.

B) Erhärtende Verbände.

Die erhärtenden Verbände teilt man in rasch erhärtende und in langsam erhärtende Verbände ein.

1. Rasch erhärtende Verbände.

Der beste der rasch erhärtenden Verbände ist der 1852 von Mathysen erfundene

Gipsverband.

Gips ist schwefelsaurer Kalk $CaSO_4 + 2H_2O$. Der Gips, wie wir ihn zur Verbandtechnik brauchen, wird in grossen Gipslagern gefunden und ist im Naturzustand nicht zu verwenden. Er muss erst, um für uns dienlich zu sein, gebrannt werden. Durch dieses Brennen verliert er sein Krystallwasser und bekommt die Eigenschaft, mit Wasser zu einem Brei angerührt, sein Krystallwasser wieder aufzunehmen und zu erhärten. Der Gips wird also dadurch hart, dass er sein beim

Brennen verlorenes Krystallwasser wieder aufnimmt.

Nimmt der Gips dagegen Wasser aus der Luft auf, so wird er unbrauchbar, indem er nicht wieder erhärtet. Man muss den Verbandgips daher an völlig trockenen Orten aufbewahren. Will man grössere Quantitäten Gips vorrätig halten, so schafft man sich am besten grosse Thonkruken an, die mit einem für Wasser undurchlässigen Stoff, wie Billrothbattist oder Guttaperchapapier überzogen werden. Auch in gut verschliessbaren Blechkisten hält sich der Gips ganz gut, wenn diese Kisten an trockenen Orten stehen. Einzelne Gipsbinden werden in der Regel in Blech- oder Papphülsen luftdicht verschlossen gehalten und verkauft.

Ist der Gips durch Aufnahme von Wasser aus der Luft verdorben, so kann man ihn durch vorsichtiges Erhitzen, das man so lange fortsetzt, bis keine Wasserdämpfe mehr aufsteigen, wieder brauchbar machen. Durch zu starkes Erhitzen wird er, wie man sagt, „totgebrannt“ d. h. unfähig, Wasser aufzunehmen und zu erhärten.

Man wählt zur Verbandtechnik stets nur feinsten Alabastergips, der zu einem feinen weissen Mehl gemahlen ist und absolut keine grobkörnigen Beimengungen enthalten darf.

Guter Gips wird schon nach wenigen Minuten hart. Man kann das Erhärten des Gipses aber dadurch noch beschleunigen, dass man zu einer Waschschüssel voll Wasser etwa eine Hand voll Alaun hinzusetzt.

Wir brauchen den Gips zum Verband in verschiedener Form.

1. Gipsbindenverband.

Weitaus in der grössten Mehrzahl der Fälle bedient man sich heutzutage des Gipsbindenverbandes. (Taf. 117.)

Die zum Gipsbindenverband notwendigen Gipsbinden müssen mit grosser Sorgfalt hergestellt werden, denn von ihrer Güte ist man völlig abhängig.

Als Material zu Gipsbinden braucht man neben dem Gips entweder die gestärkten blauen Gazebinden oder gewöhnliche Gazebinden. In beiden Fällen muss die Gaze

recht engmaschig sein, und das Gipspulver muss fest in die Maschen eingerieben werden.

Dieses Einreiben des Gipspulvers geschieht entweder mit der Hand oder mittelst eigener Gipsbindenmaschinen.

Am rationellsten ist das Einreiben des Gipspulvers mit der Hand. Man schüttet einen grösseren Haufen guten Gipses auf einen breiten Tisch, legt zur linken Seite dieses Gipshaufens den Anfang der mit dem Gips zu imprägnierenden Binde hin und bringt nun mit der rechten Hand von dem Gipshaufen eine gute Handvoll des Gipses auf das Bindenende, zieht dieses etwas hervor, streicht nun mit der ulnaren Kante der rechten Hand den Gips nach dem Bindenende hin und wickelt den so behandelten Teil der Binde, d. h. den Bindenanfang, wie eine gewöhnliche Binde, aber recht lose, auf. So fährt man mit dem Einstreichen des Gipses in die Binde fort und hat dann nach Aufwickelung des jeweils imprägnierten Teiles schliesslich eine Gipsbinde hergestellt, auf die man sich wirklich verlassen kann. Braucht man sie nicht sofort, so empfiehlt es sich, sie in Papier einzuwickeln und in einer verschliessbaren Blechschachtel aufzuheben.

Der Fehler, der bei der Herstellung der Gipsbinden meist begangen wird, ist der, dass man das Gipspulver in zu dicker Schicht auf die Binde einstreut, so dass die Binde nicht wirklich imprägniert wird, sondern der Gips, beim späteren Einlegen der Binde in Wasser, einfach herausfällt.

Gestärkte, mit Gips imprägnierte Binden erhärten nicht so schnell, als einfache Gaze-Gipsbinden. Man wählt daher überall da, wo man ein rasches Erhärten des Verbandes wünscht, lieber die einfachen Gazebinden.

Von den zahlreich angegebenen Gipsbindenmaschinen ist die einfachste und praktischste die von Beely. Der Gips wird in einen viereckigen Kasten gethan, der 2 Spalten hat, eine breitere in der halben Höhe der einen Querwand und eine schmälere am Grund der entgegengesetzten Querwand. Die letztere Spalte

lässt sich durch eine Schiebevorrichtung zur Regulierung der Gipsmenge beliebig grösser und kleiner machen. Der Anfang der Binde wird durch die breitere Spalte in den Kasten, durch den Gips hindurch und durch die schmälere Spalte aus dem Kasten herausgeführt. Der überflüssige Gips wird durch die Schiebevorrichtung abgestrichen und die Binde nun recht lose aufgewickelt. (Tafel 118a.)

Zum Anlegen des Gipsverbandes braucht man ausser den Gipsbinden noch ein Gefäss mit warmem Wasser und event. noch ein Gefäss zur Herstellung eines Gipsbreies. Dieser Gipsbrei wird so gemacht, dass man in das Gefäss eine Handvoll Gips thut und auf diese etwa die gleiche Menge warmen Wassers giesst und nun Gips und Wasser innig vermischt, wobei man mit den Fingern sofort alle entstehenden Knollen sorgfältig verreibt, so dass ein absolut gleichmässiger, gar keine Bröckel mehr enthaltender Brei entsteht.

Die Technik des Gipsverbandes gestaltet sich nun folgendermassen: Erstlich gibt man dem Gliede die Stellung, die es nach dem Verbande haben soll. Man wird aber den Gipsverband nur selten auf die blosse Haut anlegen; will man dies thun, so muss man die Haut gut einölen, damit der Gips nicht mit den Härchen verklebt und der Patient nachher beim Abnehmen des Verbandes Schmerzen hat.

In der Regel soll man den Gipsverband unterpolstern. Diese Unterpolsterung macht man entweder durch vorherige sorgfältige Einwickelung des Gliedes mit einer Flanellbinde oder durch Ueberziehen eines Trikotschlauches oder durch Umwickeln von Watte.

Der neuerdings vielfach gebrauchte Trikotschlauchüberzug ist recht praktisch. Man hält sich für die verschiedenen Körpergrössen verschieden breite Trikotschlauchbinden (Firma Römpler in Zeulenroda) vorrätig. Diese Schlauchbinden werden wie Strümpfe über die Extremität herübergezogen, wobei jede Faltenbildung vermieden werden muss. Die Schläuche haben den

Tafel 118.

a) Gipsbindenwickelmaschine nach Beely. b) Tricotschlauchbinde als Unterpolsterungsmaterial und zugleich als Handhabe. c) v. Esmarch'sches Gipsmesser. *d) Gipsschere nach Seutin.* e) *Gipsschere nach Szymanowski.* f) Gipsschere nach Bruns.

Vorteil, dass man sie etwa handbreit über das Ende einer Extremität, die Hand oder den Fuss, frei hervorragen lassen kann, so dass man eine gute Handhabe erhält, um das Glied in der gewünschten Stellung zu halten. (Tafel 118b.)

Zur Polsterung mit Watte wickelt man sich rohe, gereinigte Baumwolle in Form von Binden auf und wickelt diese Wattebinden dann um das Glied herum. Diese rohe Watte eignet sich zur Polsterung besser, als die entfettete Watte, da diese den Schweiss aufsaugt und sich deshalb leichter zusammenballt.

Besonders hervorragende Knochenteile, wie die Spinae ilei ant. sup., die Knöchel, die Tibiakante, polstert man, zumal bei mageren Patienten noch besonders, indem man einen Kranz von Watte oder Filzstücke auf sie legt.

Auf die gepolsterte Extremität werden nun die Gipsbinden aufgewickelt. Zu dem Zweck taucht man die Gipsbinde zunächst in warmes Wasser, dem man, wie oben erwähnt, zum schnelleren Erhärten des Gipses in der Regel noch eine Handvoll Alaun zugesetzt hat. Das Eintauchen der Gipsbinden in das Wasser hat nicht so zu geschehen, dass man sie ohne weiteres ins Wasser hinein steckt. Man hat vielmehr folgendes zu beachten. Erstlich soll das Gefäss so viel Wasser enthalten, dass die Gipsbinden völlig untertauchen. Zweitens soll man vor dem Einlegen das Bindenende etwas abrollen und es auf den Rand des Gefässes legen; sonst verkleben Bindenende und Kopf häufig so, dass man nur mit Mühe das Bindenende findet. Die Gipsbinde bleibt nun so lange im Wasser, bis keine Luftblasen mehr aufsteigen; dann nimmt man sie vorsichtig heraus und drückt sie nun aus, aber nicht etwa in der Weise, dass man sie in die Hand nimmt und diese nun zur Faust ballt, sondern man legt die Binde zwischen beide Hohlhände und drückt die beiden Hände sanft und gleichmässig gegen einander. So entfernt man das überflüssige Wasser und etwa überflüssigen Gips. Bevor man nun diese Binde umwickelt, legt man stets gleich noch eine neue Binde in derselben Weise in das Wasser

ein, die sich dann während des Anlegens der ersten sicher mit Wasser vollsaugt. Je loser und besser die Gipsbinden gewickelt sind, um so leichter gestaltet sich die ganze Arbeit.

Beim Anwickeln der Gipsbinden verfährt man wie beim Anwickeln jeder anderen Binde. Man vermeidet dabei aber sorgfältig jedes Anziehen der Binde, lässt diese vielmehr nur so um das Glied herum abrollen, dass die Binde von einer Hand in die andere gewissermassen fällt. Hat man mehrere Touren abgewickelt, so streicht und glättet man mit der einen Hand kreisförmig über sie herüber, damit sie sich dem Körperteil gut anschmiegen und gut an die vorhergehende Tour anlegen. Umschläge mit der Binde braucht man nicht zu machen. Die beim Wickeln abstehenden Teile der Binde streicht man vielmehr einfach mit der Hand fest. Wo es, wie beim Gehverband, darauf ankommt, den Gipsverband den Körperkontouren auf das innigste anzuschmiegen, da schneidet man, anstatt Umschläge zu machen, die Binde lieber mit der Schere ab, um sie dann von neuem wieder anzulegen. Wenn man stets so wickelt, dass die folgende Tour die vorhergehende etwa zu zwei Dritteln deckt, so braucht man zu einem zweckentsprechenden Gipsverband durchschnittlich vier aufeinanderliegende Lagen von Gypsbinden.

Gewöhnlich ist das Anlegen des Verbandes mit dem Umlegen der Gipsbinden beendigt; in besonderen Fällen kann man nun noch etwas Gipsbrei über den fertigen Verband verstreichen. Man gleicht damit alle Unebenheiten des Verbandes aus und gibt ihm dadurch ein gefälligeres, schöneres Aussehen.

Ist man fertig mit Wickeln, so muss man eine besondere Sorgfalt stets noch dem obern und untern Rande des Verbandes schenken; man muss darauf sehen, dass diese Ränder nicht einschneiden und muss zu dem Zwecke, namentlich in der Leistenbeuge, öfters noch etwas mit der Schere oder dem Messer abtragen, wenn man zu hoch hinaufgewickelt hatte. Auch einige Längsschnitte an den Rändern und Umkrämpeln der eingeschnittenen

Tafel 119.



a) Zange zum Öffnen der Gipsverbände nach J. Wolff (Wolffsmaul). b) Gipsschere nach Stille. c) Gipsschere nach Empfenzeder. d) Kreis-Gipssäge. e) Blatt-Gipssäge. f) Aufsägen eines Gipsverbandes.



Randteile schützen gut vor einem zu starken Druck des Verbandrandes. Ist der Verband an den Rändern noch weich genug, so kann man auch recht gut mit dem Zeigefinger unter den Verbandrand eindringen und ihn ringsherum nach aussen umlegen, so dass auch so jedes Einschnüren vermieden wird. Hat man mit Flanell- oder Gazebinden unterpolstert, so kann man an beiden Enden des Verbandes vorstehende Teile dieser Binden nach dem Trocknen des Verbandes wie eine Manschette umschlagen und mittelst Heftpflaster befestigen.

Guter Gips trocknet in der Regel nach wenigen Minuten, so dass der Verband hart ist, sobald man fertig ist. Will man das Trocknen beschleunigen, so bringt man die eingegipsten Glieder in die Nähe eines Ofens und lässt den Verband unbedeckt.

Unter allen Umständen müssen die Zehen- und Fingerspitzen frei bleiben; sie dienen, wie oben angegeben, zur Kontrolle des guten Sitzens des Verbandes.

Um Gipsverbände gegen Feuchtigkeit zu schützen, hat man verschiedene Mittel. Dieffenbach tränkte den fertigen Gipsverband mit einer Lösung von Kolophonium in Alkohol (1:12); Mitscherlich benutzte eine spirituöse Schellacklösung (3:50) oder eine Lösung von Damarharz in Aether (1:4). Herrgott bestrich den Gipsverband einfach mit Wagenlack.

Das Abnehmen des Gipsverbandes vom Körper ist in der Regel schwieriger als das Anlegen desselben. In jedem Falle kann man sich das Abnehmen des Verbandes dadurch erleichtern, dass man den Gips mit einer konzentrierten Kochsalzlösung durchtränkt. Man macht dies am besten so, dass man sich eine solche Lösung herstellt, dann etwas Watte oder einen Schwamm in die Lösung eintaucht und die Watte oder den Schwamm auf den Gips aufdrückt. Der Gips wird durch die Berührung mit dem Kochsalz weich und locker und lässt sich so unendlich viel leichter aufschneiden.

Zum Aufschneiden des Gipsverbandes dienen die Gipsmesser, Gipsscheren und Gipssägen.

Das Gipsmesser nach v. Esmarch ist ein

kräftiges, kurzes Messer mit bogenförmiger Schneide. (Tafel 118c.)

Gipsscheren sind in mannigfachster Form angegeben worden von Seutin, Sczymanowski, Bruns, Empfenzeder u. a. Unbedingt die beste ist die neue Gipsschere von Stille in Kopenhagen, mit welcher das Aufschneiden ausserordentlich leicht zu bewerkstelligen ist. (Tafel 118 d, e, f, und Tafel 119 b u. c.) Tafel 120 zeigt die Technik der Anwendung der ersten Stille'schen Schere. Man führt die eine Branche unter den Verband und führt nun die untere Branche der oberen in einem Bogen entgegen. So drückt die Schere nicht nur, sondern sie schneidet und so lassen sich mit der Schere selbst die stärksten Verbände spielend leicht eröffnen.

Gipssägen sind entweder als Kreissägen (Leiter) oder als Blattsägen (v. Bergmann) in Verwendung. (Tafel 119 d u. e.) Neuerdings hat man wohl auch empfohlen, Kettensägen (Schinzinger) oder andere dünne Sägen (Gigli'sche Sägen) mit einzugipsen und mit diesen den Verband von innen nach aussen zu durchsägen (Schinzinger).

Um beim Aufschneiden des Verbandes nicht die unterliegende Haut zu verletzen, empfiehlt es sich oft, einen schmalen Streifen angefeuchteter Pappe oder einen schmalen Blechstreifen oder ein Stück Schnur direkt auf die Haut zu legen, in der Linie, in der man später den Verband aufzuschneiden beabsichtigt. Das Instrument läuft dann beim Aufschneiden auf diesem Streifen und die Haut ist sicher geschützt.

Hat man den Verband der Länge nach aufgeschnitten, so bricht man ihn nach beiden Seiten hin ein, indem man die Finger beider Hände in den klaffenden Spalt des Verbandes einzwängt. Sehr praktisch ist zu diesem Zweck auch eine von Julius Wolff angegebene Zange, mit der man die Ränder sehr bequem nach der Seite einlegen kann. (Tafel 119 a.)

Wir haben bisher nur von dem einfachen Gipsbindenverband gesprochen. Mit demselben lassen sich in unzählige Modifikationen eingehen.

Tafel 121.

a) Fenster im Gipsverband. b) Völker's Holzspahn-Gipsverband.
c) Unterbrochener Gipsverband. d) Gipsverband mit Gelenkschienen.

Zunächst ist es des öfteren angezeigt, ein sogen. Fenster in den Verband zu schneiden, um z. B. eine in der Tiefe gelegene Wunde verbinden zu können. Man schneidet dann an der betreffenden Stelle mit einem scharfen Messer ein viereckiges Stück des Gipsverbandes aus und umklebt die Ränder nachträglich mit amerikanischem Heftpflaster, damit sie nicht bröckeln oder einschneiden. (Tafel 121.)

Weiterhin ist es nicht selten wünschenswert, dem Verbande eine besondere Festigkeit zu geben. Man fügt ihm zu diesem Zwecke stützende Einlagen, sogen. Verstärkungsschienen hinzu. So hat man Tapetenspan, Fournierholz, Pappe, Zinkblech oder Eisenblech, Telegraphendraht, Guttapercha, Filz und noch viele andere Materialien zwischen die einzelnen Schichten der Gipsbinden eingeschaltet. Wir bilden als Beispiel den Völker'schen Holzspan-Gipsverband ab. (Tafel 121 b.)

Man kann ferner recht gut unterbrochene Gipsverbände anlegen, indem man die Kontinuität des Verbandes durch je nach Bedürfnis gebogene Schienen aus Bandeisen oder Telegraphendraht herstellt. Solche unterbrochenen Gipsverbände kommen zumeist an den Gelenken in Anwendung. (Tafel 121 c.)

Will man dabei die Gelenke beweglich haben, so gebraucht man zum Eingipsen sogen. Gelenkschienen, d. h. Schienen, welche mit einem Scharnier oder mit Sektoren versehen sind. (Tafel 121 d.)

Sehr oft verwertet man die Elastizität des Gipsverbandes, um ihn als abnehmbare Hülse zu gebrauchen. Zu dem Zweck darf der Verband nicht zu dick sein. Er ist dann nach dem Aufschneiden so nachgiebig und dehnbar, dass keine weiteren gelenkartigen Vorrichtungen zu seinem Aufklappen notwendig sind. Man schneidet den abnehmbar zu gestaltenden Gipsverband in der Regel in der Mittellinie des Körpers exakt auf, nimmt ihn vorsichtig vom Körper ab, befestigt an ihm durch Aufnähen mit festen Fäden Vorrichtungen zum Schnüren oder Schnallen, legt ihn

wieder an und fixiert ihn durch Zuziehen der Schnüre oder Schnallen. So kann man nicht nur einfache Hülsen für die Extremitäten anfertigen (Tafel 122a), sondern auch kompliziertere, z. B. für das Becken, bei welch letzterem man dann die Ränder der Hülse in der Regel noch mit einer Lederpolsterung versieht. (Tafel 122b.)

Zum Anlegen des Gipsverbandes an der unteren Extremität, namentlich am Unterschenkel und am Becken, ist eine besondere Lagerungsweise des Patienten notwendig. Um nämlich das Becken bequem stützen zu können, bedient man sich in solchen Fällen der Beckenstützen, die in ihrer bekanntesten Form von v. Esmarch (Tafel 122c) und v. Bardeleben (Tafel 122d) angegeben worden sind.

Zwei Gehilfen müssen dann beim Anlegen der Verbände die Beine in der gewünschten Stellung erhalten. (Tafel 122e.)

Damit man diese Gehilfen auch entbehren kann, hat man neuerdings besondere Extensionsvorrichtungen angegeben, von denen die einfachste und beste von Bruns beschrieben und empfohlen worden ist. (Taf. 123a.)

Recht praktisch ist auch ein vor kurzem von Grosse angegebener Lagerungsapparat. Derselbe hat den Vorteil, dass er leicht für jede Grösse einzustellen ist. Er wird von der Firma Baumgärtel in Halle a/S. geliefert.

Eine eigene Art der Lagerung zur Anlegung von Beckengipsverbänden ist die Dittel'sche. Dittel legt seine Patienten auf zwei runde Eisenstäbe von etwa $1^1/_3$ cm Durchmesser so, dass Kopf und Rumpf auf der Stange auf dem Operationstisch liegen, während die Beine auf den beiden Stangen in der Weise ruhen, dass ein Gehilfe die beiden Malleoli interni gegen die Stangen andrückt. Der Gipsverband wird dann angelegt. Nach Trocknen des Verbands werden die vorher gut eingeölten Stangen aus dem Verband herausgezogen. Das Verfahren ist sehr praktisch und wird von mir sehr oft geübt. (Tafel 123b.)

Einen ausserordentlich praktischen Tisch zur Anlegung von Gipsverbänden in der Beckengegend bei

Tafel 122.



a) Abnehmbare Gipshülle für das Bein. b) Abnehmbare Gipshülle für die Hüfte. c) Beckenstütze nach v. Esmarch. d) Beckenstütze nach v. Bardeleben. e) Lagern des Patienten auf der Beckenstütze.

Tafel 123.

a



b

c

a) Lagerung des *Patienten* auf dem Bruns'schen Apparat.
b) „ „ „ „ den Dittel'schen Stangen.
c) *Albers'sche Kragenschiene.* d) Breigers'sche Gipswattesch



Tafel 124.



Extensionstisch von Schede.

gleichzeitiger Extension an den Beinen hat Schede konstruiert. (Tafel 124.) Ein Lagerungstisch ist durch eine vertikale Achse, die zugleich Trägerin der Kontraextension ist, mit zwei langen Extensionsrahmen verbunden, derart, dass diese letzteren um die Axe nach beiden Seiten horizontale Kreisbogen beschreiben können, mit einem Ausschlage von zusammen über 180°. Die Enden der Rahmen sind von den kraftübertragenden Zahnradkurbeln gekrönt.

Der Tisch hat die Einrichtung, dass sein vorderer Teil in Form zweier Klappen heruntergelassen werden kann, und damit die Beckengegend auch dorsal frei zugänglich wird; die Verschieblichkeit der Tischplatte erlaubt, diesen freien Raum der Grösse des Patienten anzupassen. Die Höhe des Tisches ist eine ziemlich beträchtliche, um ein kraftvolles Einstemmen der Faust hinter die Trochantergegend zu gestatten.

Die Kontraextension sollte starken Zug in unschädlicher Art auf das Perineum, resp. die Vulva übertragen und war daher ein technisch schwieriges Problem. Es ist hier in der Weise zu lösen gesucht, dass ein gummigepolsterter Lederzug als Sehne in einen Eisenbügel gehängt ist, der, in einem Nussgelenk der Vertikalaxe artikulierend, in beliebiger Stellung durch Schraube fixiert werden kann.

Dem starken Zug gegenüber wird die gewünschte Neigung des Bügels durch einen Haltegurt gewährleistet, der oberhalb des Patienten über einen Aufsatz am Kopfende des Tisches hinweggeleitet hier festgeschnallt wird. Den seitlich gerichteten manuellen Manövern gegenüber bewahrt die Kontraextension eine gewisse Nachgiebigkeit unbeschadet der Stärke des Zuges. Die ganze Kontraextension ist leicht abzunehmen.

Die Extensionsrahmen laufen auf Rollen und gestatten, von der Parallelstellung ausgehend, jede beliebige Abduktion der von ihnen geführten Extremitäten; ein Führungskreisbogen erlaubt, jede beliebige Stellung in messbarem Winkel sofort festzustellen. Die Endpfeiler tragen das doppelte Zahnradgetriebe mit Drehkurbeln,

welch letztere ein ganz allmähliches Anziehen und Nachlassen der Zugschnüre durch Auf- und Abdrehen übermitteln. Die Zugschnüre finden paarweise ihren Angriffspunkt an Fusskappen, die sich in einer nach Fussrücken und Hacke auseinanderweichenden Form am besten bewährt haben und in 2 oder 3 Grössen für alle Fälle genügen.

Den Angriffspunkt des Zuges oberhalb des Kniegelenks zu legen, scheiterte leider an der Schwierigkeit, für starken Zug den nötigen Halt zu finden; schwächeren Zug vertragen Gelenk, Nerven und Muskeln ohne nennenswerte Schädigung.

Zur genauen Regulierung des Zuges ist in jedes der Schnurpaare ein Dynamometer eingeschaltet, das ein fortwährendes Ablesen und Korrigieren der Zuggrösse bis auf Kilogramme gestattet; häufigere Benutzung macht allerdings von Zeit zu Zeit eine Revision seiner Angaben notwendig.

Ausser der horizontalen Ausschlagsfähigkeit ist noch eine gewisse Winkelwirkung um eine quere Axe möglich, da sowohl der Stützpunkt des Beckens auf der Kontraextension wie auch die kurbeltragenden Pfeiler erheblich höher gestellt werden können und dadurch in beschränkter Weise eine Flexion wie Hyperextension im Hüftgelenk anwenden lassen.

Der Schede'sche Tisch hat folgende vorzügliche Eigenschaften:

1. Doppelseitige Extension:
   a) in beliebiger Grösse,
   b) mit genauer Dosierung und Kontrolle,
   c) in beliebiger Spreizstellung,
   d) mit gewisser Variation der Beugung — Streckung im Hüftgelenk,
   e) mit Freiheit der Rollung des Beines,
   f) mit Freiliegen der Hüftgegend zu manuellen Einwirkungen.
2. Rasche Ausschaltung der Extension; Anwendung manueller Manöver auf demselben Tisch.



Schede's Tisch zur Anlegung von Gipsverbänden am Rumpf.



Verband- und Extensionstisch von Heuser.

3. Aeusserste Bequemlichkeit beim Anlegen des Gipsverbandes in jeder gewünschten Stellung.

Im Anschluss an diesen Schede'schen Extensionsisch für die Anlegung von Beckengipsverbänden will ch gleich noch den ebenso vorzüglichen Schede'schen Tisch zur Anlegung von Gipsverbänden am Rumpf bei leichzeitiger Extension an den Beinen, dem Arm und m Kopf abbilden. Diese Abbildung spricht wohl für ich selbst (Tafel 124 a). Man kann, und das ist ein sehr rosser Vorzug dieser Tische, eine beliebig dosierte eklinationslage auf demselben darstellen. Er eignet ch daher sehr zur Anlegung von Gipsverbänden, zur ehandlung der Spondylitis.

Zu dem gleichen Zweck wie Schede hat auch Heuser einen sehr guten Verband- und Extensionstisch konstruert. Derselbe dient nicht nur zur Einrenkung angeborener Hüftgelenksverrenkungen und zur Anlegung Calot'scher Verbände, sondern auch zur Anlegung chirurgischer Verbände jeder Art, die in Extension angelegt werden müssen. Die Patienten befinden sich dabei in Brusthöhe, sodass man auch die Unterseite des Verbandes bequem übersehen kann. Der Gebrauch des Tisches erhellt unmittelbar aus der Abbildung (Tafel 125).

Beely'schen Gipshanfschienen. (Tafel 126.)

Die in früheren Jahren vielfach gebrauchten Methoden des Gipsverbandes, der Gipsstreifenverband und der Gipsumschlag, sind heutzutage völlig verlassen worden. Man verwendet dagegen jetzt vielfach die

Gipsschienen,

lie in verschiedener Art und Weise hergestellt werden Wir haben da zunächst die Gipshanfschienen.

Zum Anlegen einer Gipshanfschiene braucht nan Wasser und Gipspulver zum Anrühren eines Gipsreies, gut ausgehechelten Hanf von etwa 50—80 cm Länge, ein Stück Leinwand zum Auflegen auf die Haut und einige Rollbinden. Aus dem Hanf macht man sich kleinere Bündel, die locker hingelegt, eine Breite von

etwa 3—4 cm, bei 1 cm Dicke, haben. Das Anlegen der Schiene geschieht nun in der Weise, dass man dem Gliede zunächst die gewünschte Stellung gibt; dann legt man direkt auf die Haut, um das Ankleben des Gipses an die Haare zu verhüten, das gut angefeuchtete Leinwandstück, dessen Ränder man vorher in Abständen von etwa 5 cm eingeschnitten hat und das etwas länger und breiter sein muss als die gewünschte Schiene; dann macht man sich den Gipsbrei, nimmt eines der vorher präparierten Hanfbündel, zieht dasselbe durch den Brei hindurch, wobei man dafür sorgt, dass der Brei auch ordentlich zwischen die einzelnen Fasern eindringt, streift den überflüssigen Brei durch Hindurchziehen des Bündels zwischen Zeige- und Mittelfinger der linken Hand ab und legt nun das Bündel parallel der Längsachse des Gliedes auf die Leinwand auf. In derselben Weise wie dieses erste Bündel werden nun auch die anderen Bündel getränkt und dann dicht neben einander gelegt oder so, dass sie sich zum Teil noch gegenseitig decken. So fährt man fort, bis eine Schiene von gewünschter Länge und Breite und in der Mitte etwa 2 cm Dicke gebildet ist. Nach den Seitenrändern zu lässt man die Schiene allmählich an Stärke abnehmen. Im allgemeinen sollen die Schienen etwa halb oder ein drittel so breit sein, wie die Circumferenz des betreffenden Gliedes. Ist die Schiene fertig, so werden die überstehenden Teile der Leinwandunterlage über die Ränder der Schiene herübergeschlagen und mit etwas Gipsbrei verstrichen. Ist der Gipsbrei erstarrt, so wird der Verband dadurch vervollständigt, dass man die Schiene mittelst einer Rollbinde an die Extremität anbandagiert.

Will man die Gipshanfschiene so einrichten, dass sie auch zur Suspension des Gliedes dienen kann, so kann man sehr leicht einige Eisenringe mit eingipsen. Man nimmt dann einen dünneren Hanfstreifen, zieht über diesen so viel Ringe, als benötigt sind, und legt dann diesen mit Ringen versehenen Streifen in die Mitte der schon aufgelegten Gipshanfstreifen, schiebt die Ringe an die betreffenden Stellen, wo sie hinkommen sollen.

Tafel 126.



Beely'sche Gipshanfschienen.
a, b) für die obere Extremität;
c, d) für die untere *Extremität* (mit Suspensionsvorrichtung).

und tränkt dann diesen Streifen mit noch gut bindendem, aber ja nicht zu dickem Brei. Dicht neben ihn und zum Teil auch auf ihn legt man dann noch weitere Gipshanfstreifen. (Tafel 126 c, d.)

Die Beely'schen Gipshanfschienen haben den Vorzug, dass sie sich der Körperform genau anschmiegen, dass sie leicht abgenommen werden können und dass sie sehr haltbar sind. Sie haben den Nachteil, dass ihre Anfertigung nur dann möglich ist, wenn man guten Hanf, guten Gips und gute Assistenz zur Seite hat.

Diesen Nachteil hat Braatz beseitigt, indem er das Beely'sche Prinzip beibehaltend, die Gipsschienen für die Praxis dadurch bequemer und zugänglicher machte, dass er die Schienen statt aus Hanf aus Baumwollentrikot oder auch wohl aus einem sackleinwandartigen Gewebe, sogen. Formleinen oder Halbleinen herzustellen lehrte.

Eine technische Vereinfachung für die Herstellung von Gipshanfschienen hat Turner angegeben: Ein Bündel gut durchhechelten Hanfes von der Länge der gewünschten Schiene wird an seinen Enden mit der Schere gerade durchschnitten und dann in toto — also nicht, wie Beely rät, in einzelnen Strähnen — in warmen Gipsbrei von der Konsistenz gewöhnlicher Sahne getaucht, wo es mit den Fingern sorgfältig durchgeknetet wird, um ein möglichst vollkommenes Eindringen des Gipsbreies in alle Zwischenräume zwischen den einzelnen Fasern zu erreichen. Das so zubereitete nasse gipsgetränkte Hanfbündel wird weiterhin gleichsam wie in ein Futteral durch einen Trikotschlauch hindurchgezogen, wie solche als sogenannte gewebte Binden in dieser Breite käuflich sind. Ist auf diese Weise der vorher ebenfalls entsprechend zurechtgeschnittene Trikotschlauch über das Gipshanfbündel hinübergezogen, so wird der nunmehr im Trikotschlauch befindliche Gips-Hanf nochmals durchgeknetet, bis man eine gleichmässig dicke und an den Rändern gut abgerundete Schiene erhält. Hierzu ist es natürlich erforderlich, dass das Hanfbündel und der Trikotschlauch einander in ihrer Breite

entsprechen. Das Gewebe des letzteren wird dabei ein wenig ausgezogen, um jegliche Faltenbildung in der Schiene zu vermeiden.

Die so hergestellte Schiene wird nass an die Extremität angelegt, derselben genau adaptiert und mit einer trockenen Binde in der erforderlichen Stellung befestigt. Nach 3—5 Minuten ist die Schiene erstarrt und behält nun genau die ihr mitgeteilte Form (als Hohlrinne, spiralig gewundene oder winkelig abgebogene Schiene etc.) bei.

Die Schiene lässt sich leicht in 5—6 Minuten herstellen. Sie erstarrt schnell unter Beibehaltung der ihr verliehenen Form. Ihre Oberfläche ist eine gleichmässig glatte und dabei weniger harte als die der typischen Beely'schen Schiene, die Ränder nicht scharf, sondern gut abgerundet, was das lästige Ausbröckeln des Gipses an den Rändern der Schiene verhindert. Endlich geht die ganze Prozedur verhältnismässig rascher und ohne unnötiges Beschmutzen der Körperoberfläche des Kranken mit Gipsbrei vor sich.

Braatz's Gipstrikotschienen.

Die Technik der Gipstrikotschienen ist folgende: Man schneidet aus dem Trikot Streifen, welche etwas breiter sind, als die eigentliche Schiene werden soll, taucht diese Streifen in Gipsbrei, entfernt den überflüssigen Gipsbrei durch Abwärtsstreichen und legt nun diese Streifen auf die in richtiger Stellung gehaltene Extremität. Bei Erwachsenen legt man sie auf die blosse Haut auf; bei Kindern legt man vorher zur Polsterung einen feuchten, an den Rändern gekerbten Leinwandstreifen unter. Dann folgt als zweite Schicht noch ein solcher Gipstrikotstreifen. Genügt dieser noch nicht, so wird noch ein dritter Streifen hinzugefügt. Diese Schichten werden durch kräftiges Streichen mit der flachen Hand ohne Mühe zu einer gleichmässigen Schiene vereinigt. Es resultieren auf diese Weise Schienen von einer Schönheit und Gleichmässigkeit, wie sie bei den Beely'schen Gipshanfschienen kaum gelingen. Wünscht man einen

ganz besonders haltbaren Verband, so schaltet man nach der ersten Lage Gipstrikot einen schmalen Streifen eines dünnen Drahtnetzes ein.

Die **Albers'schen Kragenschienen.**

Viel einfacher noch als die Anlegung der eben beschriebenen Schienen ist die der Albers'schen Schienen, die man auch als Gipslonguetten bezeichnen könnte. Als Beispiel beschreiben wir die Albers'sche Kragenschiene zur Behandlung von Oberarmbrüchen. (Tafel 123 c.)

Die Schiene wird aus einfachen Gipsmullbinden von der Breite des Armes gefertigt, die vor dem Anlegen in heisses Wasser eingetaucht werden. Während nun die Repression des gebrochenen Armes stattfindet, werden die angefeuchteten Gipsbinden auf der leicht eingeölten Haut des Patienten in der Weise abgerollt, dass sie in Längstouren von der Mitte des Halses, wo ihr Anfang von einem Assistenten fixiert wird, über die Schulter, das Acromion, die Streckseiten des Ober und Unterarms und den Handrücken hinweg bis an die Köpfchen der Mittelhandknochen läuft. Hier wird die Binde umgeschlagen, fixiert und dann in umgekehrter Richtung wieder bis zur Mitte des Halses hinaufgeführt. In dieser Weise deckt man die Hälfte des Halses, die ganze Schulter, die laterale Hälfte des Oberarms, die Streckseite des Vorderarms und den Handrücken mit 8 bis 10 fachen Gipsbindenlagen zu, und die einzelnen Lagen werden stets gut verstrichen; sind sie fertig angelegt, so drückt man sie mit an der Hand beginnenden und bis zur Achsel hinaufreichenden Zirkel- resp. Achtertouren von Cambricbinden gut gegen die Unterlage an. Ist man an den Hals gekommen, so schlägt man den am Hals in die Höhe geführten Teil der Schiene als Kragen nach aussen um. Indem man diesen durch einige Bindentouren, welche unter der Achselhöhle der gesunden Seite durchlaufen, befestigt, gewinnt die Schiene nach oben einen guten Halt. Ist der Gips erstarrt, so wird die Schiene abgenommen, am Rande etwas aufgebogen, mit einer dünnen Watteschicht ge-

polstert und wieder angelegt. Der ganze Arm kommt schliesslich in eine Mitella.

Anstatt die Albers'schen Schienen durch Abwicklung von Binden herzustellen, gehe ich jetzt so vor, dass ich sowohl diese Schienen als die Gipshanfschiene ersetze durch

Gips-Stärkebindenlonguetten,

die so hergestellt werden, dass einfacher gestärkter Gazebindenstoff in 3 oder 4 Lagen kravattenförmig in entsprechender Breite und Länge zusammengelegt wird. Die Gazebindenlonguetten werden dann einfach durch Gipsbrei hindurchgezogen; den überflüssigen Gipsbrei streicht man ab, indem man die Longuetten zwischen den Fingern durchzieht und legt nun die Longuette auf den betreffenden Körperteil in der gewünschten Stellung auf. Nach ganz kurzer Zeit ist bei Verwendung guten Materiales die Schiene erstarrt und so ist eine sich der betreffenden Körperform auf das genaueste anschmiegende Schiene in kürzester Zeit und am allerbilligsten hergestellt.

Die **Breiger'schen Gipswatteschienen.** (Tafel 123 d.)

Die Breiger'schen Gipswatteschienen oder auch wohl Gipskataplasmen genannt, sind lange Säcke aus geleimter Watte, mit Gips gefüllt, von verschiedener Grösse und Breite. Die Säcke werden in heisses Wasser getaucht und dann mittelst Binden an den betreffenden Körperteil in der erwünschten Stellung angewickelt. Nach dem Trocknen stellen sie genau passende Hohlrinnen dar, die bequem abgenommen und wieder angelegt werden können. Hat man die Säcke vorrätig, so ist der Verband sehr bald angelegt.

Gipsmulldecken

zur Behandlung von Frakturen u. s. f. empfiehlt in neuester Zeit Gretsch. Gewöhnliche weitmaschige Verbandgaze wird in 4—10 facher Lage zusammengelegt. Zwischen die einzelnen Lagen wird Alabastergips gut eingerieben. Die so hergestellte Decke wird mit Cantricmehl umwickelt und trocken auf das betreffende Glied fest umgewickelt, dann mit warmer Alaunlösung befeuchtet. Nach kurzer Zeit ist die Schiene erstarrt.

Der **Wasserglasverband.** (Tafel 127.)

Wasserglas, eine 30%—60% Lösung von kieselsaurem Natron (Natronwasserglas) oder kieselsaurem Kali (Kaliwasserglas) wird zum Kontentivverband gebraucht, wenn man einen billigen, leichten und haltbaren Verband haben will.

Die Verwertung des Wasserglases geschieht in zweifacher Weise. Entweder schüttet man das Wasserglas, wie man es aus der Apotheke bezieht, eine klare, dicke, syrupähnliche, gelbliche Flüssigkeit, die in gut verkorkten Krügen aufgehoben werden muss, in eine Schüssel, legt in die Flüssigkeit die zu verwendenden Mullbinden hinein, lässt sie sich etwa 10 Minuten lang vollsaugen, presst dann das überflüssige Wasserglas aus und wickelt nun die Binden wie die Gipsbinden am Körper an, oder aber man wickelt zuerst die Mullbinden unimprägniert in gewöhnlicher Weise an und tränkt sie erst nach dem Anlegen, indem man das Wasserglas mittelst eines groben Borstenpinsels aufträgt. Das Bepinseln der Binde geschieht am besten in zirkulärer Richtung; streicht man in der Längsrichtung, so rollen sich die Ränder der Binde leicht auf, so dass Falten entstehen. Ist die eine Bindenlage getränkt, so wird eine frische Mullbinde übergewickelt und in derselben Weise behandelt. Vier Lagen genügen in der Regel zu einem haltbaren Verband.

Das Wasserglas hat den Nachteil, dass es sehr langsam, oft erst nach Tagen, erstarrt. Durch Zusätze von Kreide, Dextrin, kohlensaurem Kalk, Kalkhydrat, Calciumphosphat, Dolomit, Magnesit oder Zement kann man jedoch das Erhärten beschleunigen. Namentlich der Magnesitwasserglas-Verband erhärtet recht gut. Man setzt, um ihn zu erhalten, dem Wasserglas unter beständigem Verrühren so lange Magnesit zu, bis ein gleichmässiger, rahmartiger Brei entsteht, mit dem dann die Binden getränkt werden.

Um zu vermeiden, dass sich bei einer reponierten Fraktur die Dislokation wiederherstellt oder eine redressierte Deformität wieder zurückgeht, während der Wasserglasverband erstarrt, legt man vielfach über dem

Wasserglasverband noch einen Gipsverband an, den man dann nach dem Erhärten des Wasserglases wieder abnimmt.

Der einmal erhärtete Wasserglasverband besitzt auch ohne Verstärkungsschienen eine ausserordentliche Festigkeit. Dabei ist er leicht, sehr elastisch und dauerhaft: er ist nicht spröde wie Gips, bröckelt daher nicht ab und lässt sich deshalb sehr gut bearbeiten, d. h. mit Riemen, Schnallen, Gurten, Gelenkschienen u. s. w. versehen. Die Wasserglasverbände sind daher sehr brauchbar für orthopädische Zwecke. Besonders erwähnenswert sind die sog. „artikuliert-mobilen" Gelenkverbände von Kappeler und Hafter, in denen die Beweglichkeit der Gelenke nicht mit Hilfe von Gelenkschienen, sondern durch zweckmässig angebrachte Ausschnitte hergestellt wird, in der Weise, wie das die Abbildungen zeigen. (Tafel 127.)

Der Leimverband. (Tafel 128.)

Der Leimverband wird in neuerer Zeit wieder vielfach verwendet. Die Technik ist die, dass die beste Qualität des Leimes, der sog. „Kölner" Leim zunächst geruchlos gemacht wird. Zu dem Zweck wird er zwei Tage in Wasser geweicht, bis er zu einer sulzigen Masse aufquillt. Hierauf wird das überflüssige Wasser abgegossen und der Leim bis zur Schaumbildung aufgekocht. Nach dem Abkühlen wird dieselbe Prozedur noch zweimal wiederholt.

Zum Anlegen des Verbandes werden Leinwandstreifen mit dem Leim bestrichen. Diese Leinwandstreifen sind schmal und besitzen gerade die Länge, dass sie, ohne eingeschlagen werden zu müssen, sich dem Umfang des Gliedes faltenlos anpassen. Der auf diese Streifen aufzutragende Leim muss erst heiss gemacht werden. Dies geschieht am besten in einem doppelwandigen Kessel, damit die Temperatur nicht über 100 Grad hinausgeht Ist der Leim heiss, so wird er auf die Leinwandstreifen aufgestrichen und diese dann, jeder einzelne für sich, in der Weise aufgelegt, dass man mit

Tafel 127.



Artikuliert-mobile Wasserglasverbände nach Kappeler und Hafter.

Tafel 128.

a) Anlegung eines Leimverbandes.
b, c, d) Verschiedene Rohrgeflechte.

dem ersten Streifen an der Peripherie beginnt, dann den nächsten Streifen diesen ersten zu etwa einem Drittel decken lässt, und so fort, bis man von der Peripherie zu dem zentralen Ende des Verbandes gelangt ist. Je heisser der Leim ist, desto besser klebt er, und desto schneller trocknet und erhärtet er. Das Erhärten ist nach kurzer Zeit, etwa nach 3—5 Stunden, beendet. Der so erzielte Verband ist leicht, übt eine gute Kompression aus und kann ebenfalls abnehmbar gemacht werden.

Der Holzleimverband.

An die Leimverbände schliessen sich unmittelbar die Holzleimverbände Walltuch's an. Das zu diesem Verband notwendige Material sind Hobelspäne, die etwa 6 cm lang, 5 cm breit und 0,5 mm dick sich von selbst zu Holzbinden aufrollen. Diese Holzbinden werden durch „Kölner Leim" miteinander verbunden. Der Leim wird wiederum etwa 8—10 Stunden in kaltem Wasser erweicht und dann in einem Wasserbade aufgekocht. Er muss so dick sein, dass man beim Führen des Pinsels einen Widerstand spürt. Durch Zusatz von etwa 5 % Glycerin — etwa 3—4 Esslöffel zu einem Liter Leimlösung — wird der Leim nach dem Eintrocknen recht elastisch und durch Zusatz von 8—10 Kaffeelöffel von doppeltchromsaurem Kali auf ein Liter Leimlösung widerstandsfähig gegen Durchnässung. Die Holzverbände müssen immer über einem Gipsmodell gemacht werden, das man vorher zweckmässig mit Trikot überzieht. Um das Anlegen der Holzstreifen auf so unregelmässigen Flächen, wie es die Körperformen sind, zu ermöglichen, spaltet man die einzelnen Streifen von beiden Seiten her auf kurze Strecken mit einem Messer auseinander. Die einzelnen Streifen decken sich wieder zu etwa einem Drittel und werden in der mannigfaltigsten Weise zirkulär, diagonal und spiralig angelegt. Drei Schichten genügen in der Regel. Der vom Modell abgenommene erhärtete Verband wird zur Erlangung grösserer Festigkeit von innen und aussen noch mit Rohleinwand überzogen.

Die Holzverbände sind ausserordentlich haltbar und

leicht und gewinnen daher immer mehr an Verbreitung. Namentlich die Holzkorsette bei Skoliosen sind ausserordentlich beliebt.

Der geleimte Cellulose-Verband.

Als Ersatz für den eben beschriebenen Walltuchschen Holzleimverband empfiehlt Hübscher die geleimte Cellulose. Die im Handel in breiten Rollen zu beziehende Cellulose wird entsprechend zugeschnitten und in lauwarmem Wasser so lange durchfeuchtet, bis sich beim Reiben der Celluloseplatten zwischen den Fingern kleine Teilchen von der Oberfläche abrollen lassen. Dann wird die feuchte Platte dem Modell genau angepasst und auf demselben getrocknet Die Cellulose, die jetzt völlig die Form ihrer Unterlage angenommen hat, wird nun von derselben abgenommen, mittelst eines Borstenpinsels reichlich mit dünnflüssigem Leim auf beiden Seiten bestrichen und sofort wieder an ihre frühere Stelle auf das Gipsmodell aufgelegt, wobei die Ränder übereinander geleimt werden. Nun wird noch eine zweite dünnere Schicht Cellulose entweder sofort oder noch besser nach dem Trocknen der ersten aufgeleimt, dann ist der Verband fertig. Zum Tragen wird er noch gefüttert und mit einer Schnürvorrichtung versehen. Die Verbände sind sehr leicht, elastisch und widerstandsfähig. (Bezugsquelle der Cellulose, die Cellulosefabrik von Simonius in Kelheim.)

Der Rohrgeflecht-Leimverband. (Tafel 128 b, c, d.)

Ein aus Rohrgeflecht und Leim zusammengesetzter Verband ist schon seit längerer Zeit in der chirurgischen Klinik zu Leipzig im Gebrauch. Weit- oder engmaschiges Rohrgeflecht oder ein Geflecht, in dem die Rohrstäbchen in der Hauptsache in der Längsrichtung angeordnet sind, wird in schmälere oder breitere Streifen geschnitten und in heissem Wasser geschmeidig gemacht. Die betreffende Extremität wird zunächst mit einem Trikotstrumpf überzogen. Ueber denselben wird eine Mullbinde gewickelt, die dann mittelst eines Borstenpinsels mit Leim getränkt wird. Nun legt man die Rohrmatten auf, die ebenfalls

erst durch den dünnflüssigen Leim hindurchgezogen werden. Auf diese Einlagen folgen wieder zwei Leim-Mulllagen, und damit ist der Verband vollendet. Nach 12 Stunden ist er soweit getrocknet, dass die Kapsel nach der Abnahme die Form bewahrt. Dieselbe ist fest, dauerhaft, leicht und geschmeidig. (Bezugsquelle des Rohrgeflechtes: A. Ehrich in Leipzig, Dufourstr. 15.)

Der Kleisterverband

wurde 1840 von Seutin erfunden.

Man rührt Stärkemehl mit kaltem Wasser zu einem Brei an und giesst dann unter stetem Umrühren so viel kochendes Wasser hinzu, bis ein klarer, dickflüssiger Schleim entsteht. Man braucht nun zum Kleisterverband Kleisterbinden und Kleisterschienen. Die Kleisterbinden erhält man, wenn man gewöhnliche Baumwollbinden durch den frischen Kleister hindurchzieht und sie dann aufrollt. Die Kleisterschienen stellt man dar, indem man einfache Pappstreifen rasch durch heisses Wasser zieht und dann auf beiden Seiten dick mit Kleister beschmiert.

Zur Anlegung des eigentlichen Verbandes wird das Glied zunächst sorgfältig mit einer feuchten Flanellbinde eingewickelt, nachdem man die Vertiefungen an den Gelenken mit Watte ausgepolstert hat. Nun wickelt man die Kleisterbinden darüber, legt auf diese die weichen Kleisterschienen, wickelt über diese wieder Kleisterbinden und umhüllt das Ganze schliesslich mit einer Mullbinde.

Es dauert 2—3 Tage, bis ein Kleisterverband ganz trocken und hart wird. Ist er aber trocken, so kann er ebenso wie die andern Kontentivverbände abnehmbar gemacht werden.

Dextrinverbände, Tripolithverbände, Kittverbände, Gummi-Kreideverbände, Paraffin-Stearinverbände, Cementverbände, die in früheren Jahren hie und da als Kontentivverbände gebraucht wurden, werden heutzutage kaum mehr in Verwendung gezogen.

Dagegen haben Landerer-Kirsch in den allerletzten Tagen zu Hülsenverbänden einen

Celluloid-Mullverband empfohlen.

Der Celluloid-Mullverband besteht aus Mullbinden, gestärkt mit einer Auflösung von Celluloid in Aceton. Man schneidet mit einer starken Schere Celluloidplatten in kleine Schnitzel, thut diese in eine grosse, weithalsige Flasche bis zu etwa 1/4 der Höhe derselben und giesst dann das Aceton auf, bis die Flasche voll ist. Die Flasche muss einen guten, luftdichten Verschluss haben, da sonst zu viel verdunstet. Von Zeit zu Zeit wird geöffnet und mit einem Stäbchen umgerührt.

Zur Herstellung einer Hülse überzieht man das Gipsmodell zunächst mit einem dicken Stück Flanell, Filz oder wickelt auch wohl nur eine einfache Mullbinde über, so dass sich die Touren etwa zur Hälfte decken. Auf diese Mullschicht wird nun die inzwischen fertiggestellte Celluloidgelatine eingerieben. Da sie an den Finger sehr fest klebt und nur mit Aceton abzuwaschen ist, wird die Hand am besten mit einem Lederhandschuh geschützt. Auf die Celluloidschicht wird nun wieder eine Mullbinde gewickelt, und es wechseln nun diese Schichten — Celluloidlösung und Mullbinde — weiterhin so lange ab, bis der Verband die nötige Stärke erreicht hat. Bei kleinen Hülsen genügen 4—6 Lagen; bei einem Stützkorsett sind mindestens 10 Lagen erforderlich. Die äusserste Schicht stellt nicht die Mullbinde, sondern die Celluloidgelatine dar: dadurch erhält der Verband ein schönes, glänzendes Aussehen.

Die so hergestellten Hülsen sind sehr leicht, elastisch, sehr hart und undurchlässig: sie erhärten in 3—4 Stunden.

Nimmt man anstatt des Mulles Tricot, so erhält man den Celluloid-Tricotverband (Joseph).

Celluloidplattenverbände.

Celluloidplattenverbände sind neuerdings von amerikanischen Kollegen (Phelps) und bei uns von Lorenz vielfach verwendet worden. Die Technik ist eine sehr einfache. Entsprechend zugeschnittene Celluloidplatten werden in heisses Wasser getaucht und aus diesem schnell auf die Modelle gebracht und sorg-

fältig eingewickelt. Grössere Apparate wie Korsette macht man aus zwei Teilen, die dann durch Schnürung verbunden werden.

Zugverbände

(Extensions-Distraktionsverbände).

a) Heftpflastergewichtsextensionsverband.

Unstreitig der beste und der am meisten verwendete Zugverband ist der Heftpflasterzugverband, der von Gordon Buck in Amerika ursprünglich angegeben und in Deutschland besonders von R. v. Volkmann eingeführt wurde.

Seine Technik ist die, dass an das zu extendierende Glied eine Heftpflasterschlinge angeklebt wird, an welcher wiederum eine das Gewicht tragende Schnur befestigt ist. Die Heftpflasterschlinge muss so fest an dem Gliede kleben, dass sie auch bei einem starken und anhaltenden Zuge nicht abgleitet. Man muss deshalb gutes Heftpflaster verwenden, am besten dasjenige, welches wir unter dem Namen „Sparadrap of Mead“ aus Amerika beziehen.

Von diesem Pflaster reisst man sich je nach der Breite der Glieder einen $1^1/_2$—3 Finger breiten und $^1/_2$—$1^1/_2$ Meter langen Streifen ab und klebt diesen in der Richtung, in welcher der Zug ausgeübt werden soll, an die beiden Seitenflächen des zu extendierenden Gliedes in der Weise an, dass die beiden Streifenenden gegen das centrale Ende des Gliedes zu liegen kommen, während die Mitte des Streifens als offene Schlinge das periphere Ende des Gliedes überragt. Beim Ankleben der Heftpflasterstreifen muss man darauf achten, dass keine hervorstehenden Knochenvorsprünge gedrückt werden. Zu dem Zweck wird auch die offene Schlinge durch ein schmales Brettchen, das sogen. „Spreizbrettchen“ angespreizt gehalten, so dass sie in eine Art Steigbügel verwandelt wird. An dieses Brettchen befestigt man die Schnur und leitet diese dann über Rollen, so dass die Gewichte frei herabhängen.

Wollen wir nun z. B. einen Extensionsverband an der unteren Extremität bei einem Erwachsenen anlegen, so wird ein 6—8 cm breiter Streifen zu beiden Seiten des Beines möglichst hoch oben vom Oberschenkel herab angeklebt bis etwa handbreit über die Malleolen. Hier weichen die Streifen auseinander, indem die Schlinge durch Einlegen des Spreizbrettchens in den Steigbügel verwandelt wird. (Taf. 129a.) Beim Anlegen des äusseren Streifen muss man darauf achten, dass der Nervus peroneus auf dem Capitulum fibulae nicht gedrückt wird. Bei mageren Personen legt man nun auf die Tibiakante und um die Malleolen herum etwas Watte auf und wickelt nun das ganze Bein, von den Malleolen angefangen bis zur Leiste hin, mit einer Cambric- oder Flanellbinde ein. (Taf. 129b.) Den Fuss selbst umwickelt man dann gut mit Watte und legt ihn in eine ebenfalls wohlgepolsterte v. Volkmann'sche oder noch besser in eine Bruns'sche Schiene hinein, da bei dieser letzten ein Druck auf die Ferse sicher vermieden wird. Nimmt man die Bruns'sche Schiene, so gleiten die halbkreisförmig abgebogenen Arme leicht auf der Unterlage, so dass die Reibung eine geringe und eine weitere Vorrichtung nicht notwendig ist. Wählt man dagegen die v. Volkmann'sche Schiene, so ist der von der Bettunterlage gebildete Reibungswiderstand so gross, dass ein besonderes sog. Schleifbrett unter die Schiene gelegt werden muss, um das leichte Gleiten der Schiene zu ermöglichen. (Taf. 130b.) Das Schleifbrett ist ein dünnes Brett von etwa $^1/_4$ Meter Länge, und 15—20 cm Breite, an dessen beiden Rändern zwei dreiseitige Prismen mit Leim angeklebt sind. Die Schiene gleitet dann auf den Kanten dieser Prismen. Beim Anlegen der v. Volkmann'schen Schiene muss der Fuss genau rechtwinklig stehen und die Ferse gut in den Fersenausschnitt hineinpassen. Man muss dabei das Fussblech der Schiene fest gegen die Fusssohle andrängen und den Fuss fest an die Schiene anbandagieren; sonst entfernt sich die Ferse vom Fussblech, und die Patienten bekommen unausstehliche Schmerzen und Decubitus an der Ferse. (Tafel 129c.)

Hat man nun die v. Volkmann'sche oder Bruns-

Tafel 129.



Heftpflastergewichtsextensionsverband.

sche Schiene angelegt, so befestigt man weiterhin die Schnur mittelst eines Hakens an dem Spreizbrettchen und leitet schliesslich die Schnur über die am Bett angebrachten Rollen hinweg. Diese Rollen werden entweder an die hölzerne Bettstelle angeschraubt oder an die eiserne Bettstatt durch besondere Rollvorrichtungen befestigt. Die beste und einfachste dieser Rollenvorrichtungen ist die von Koch in Neuffen angegebene, deren Konstruktion und Anwendungsweise wohl unmittelbar aus der Abbildung ersichtlich ist. (Tafel 129 c.)

Recht praktisch ist auch der Rollenträger von Ebert (Tafel 131). Ein 100 cm langes und 20 cm breites Brett ist in seinem unteren Teile auf der Fläche abgeschrägt; in seinem oberen Teile ist es durch eine 2 cm breite und 72 cm lange Spalte der Länge nach in zwei gleiche Hälften geteilt. In dieser Spalte laufen die zwei Rollenträger, welche — aus Bandeisen gebogen — durch Stifte in jeder gewünschten Höhe festgestellt werden können.

Die Grösse des Gewichtes ist je nach der Krankheit, dem Alter und der Muskelkraft des Patienten verschieden und schwankt zwischen 3 bis 30 Pfund. Als Gewichte braucht man entweder eiserne Gewichte, oder man füllt sich Sandsäcke mit der entsprechenden Menge Sand und verbindet dann die Sandsäcke mit der Schnur.

An dem eben beschriebenen ursprünglichen v. Volkmann'schen Extensionsverbande sind nun eine Reihe von Modifikationen vorgenommen worden, die aber alle nicht von wesentlicher Bedeutung sind. Wir erwähnen nur, dass, um den Druck auf die Ferse zu vermeiden, Koenig einen Schleifbügel angegeben hat: eine dorsale Schiene mit zwei seitlich angebrachten Eisenbügeln, welche das Bein in der Schwebe erhalten. Wir haben schon erwähnt, dass sich für den gleichen Zweck am meisten die Bruns'sche Schiene eignet.

Würde man nun den Patienten in das Bett legen und die Gewichte wirken lassen, so würden diese den Patienten im Bett einfach herunterziehen. Man muss also eine Vorrichtung anbringen, welche es verhindert,

dass der Körper des Kranken dem Zuge folgt. Man hat da im allgemeinen drei verschiedene Verfahren. Erstlich kann man zwischen den gesunden Fuss und das Fussende des Bettes ein festes Kissen oder eine Fussbank legen, gegen die sich der Fuss anstemmen kann. Dieser Gegenhalt geht natürlich verloren, sobald der Patient sein gesundes Knie beugt. Es ist daher wohl besser, sich gleich des zweiten Verfahrens zu bedienen. Dieses besteht darin, dass man das Fussende des Bettes durch unterschobene Holzklötze hochstellt. So verwandelt man das Bett in eine schiefe Ebene, auf der der Körper das Bestreben hat, nach abwärts zu gleiten. (Tafel 130a.)

Das beste und sicherste Verfahren, den Gegenhalt zu gewinnen, ist entschieden die Anlegung eines Gegenzuges, der sogenannten Kontraextension. Dieser Gegenzug wird so hergestellt, dass man einen wohlgepolsterten Gummischlauch oder besser eine weiche Schafwollquele zwischen den Beinen des Patienten hindurchführt und entweder am Kopfende der Bettstelle anknüpft oder ebenfalls über eine Gewichtsrolle hinüberführt. Der Gummischlauch oder die Quele erhält dann den Gegenhalt am Damme des Patienten. (Tafel 130a.)

Jeder Extensionsverband muss täglich mehrmals genau kontrolliert werden, und es muss dafür gesorgt werden, dass an keiner Stelle Decubitus entsteht.

Mittelst des Heftpflasters kann der Zug nicht nur in der Längsachse des Gliedes bewirkt werden. Man kann vielmehr auch ganz bequem seitliche Züge anbringen und so z. B. gegen die Wiederkehr der Dislokation bei einem schweren Bein- oder Armbruch erfolgreich vorgehen oder eine winklige Gelenkkontraktur bequem korrigieren. Bei der Frakturenbehandlung hat namentlich Bardenheuer das Heftpflaster in ergiebigster Weise zu verwenden gelehrt; das zeigt zum Beispiel ein Blick auf Tafel 132, welche die Bardenheuer'sche Methode zur Behandlung eines Oberschenkelbruches mit der gewöhnlichen Dislokation, d. h. mit dem nach aussen sehenden Winkel der Fragmente, darstellt. Wir haben da erstens den Längszug, zweitens einen Querzug an der Spitze des Winkels und drittens



a) Anwendung der Kontraextension (mittelst einer Quehle und Hochstellung des Bettes am Fussende).
b) v. Volkmann'sches Schleifbrett.



Rollenträger von Eberth.

Tafel 132.



a) Verwendung mehrerer Heftpflasterzüge nach Bardenheuer.
b) Schedes' Methode der vertikalen Extension bei Kindern.

Tafel 133.



a) Schede's Methode zur Streckung von Kniegelenksverkrümmungen un
uhülfenahme mehrfacher Heftpflasterzüge. b) v. Volkmann's Steckn
verband. c) Zug vermittelst des Contentivverbandes.

einen Querzug am Becken, der das Becken ruhig stellen und ein Ausweichen desselben nach der gesunden Seite hin verhindern soll. (Tafel 132a.)

Als Beispiel der mehrfachen Verwertung des Heftpflasters zur Streckung von Kontrakturen diene die Schede'sche Methode zur Streckung einer mit Subluxation der Tibia nach hinten und aussen einhergehenden Beugekontraktur des Kniegelenkes. Ausser dem Längszug am Unterschenkel sehen wir da einen das Knie nach abwärts und einen die Tibia nach aufwärts führenden Zug in leicht verständlicher Weise angebracht. (Tafel 133a.)

Da Kinder das Heftpflaster leicht beschmutzen, kann man bei ihnen die Extension sehr bequem mit der Suspension verbinden. So verwertet man heutzutage die vertikale Suspension nach dem Vorgange von Schede namentlich zur Behandlung von Oberschenkelbrüchen kleiner Kinder. (Tafel 132b.)

Wir haben bisher als Zugmittel stets das Heftpflaster im Auge gehabt. Manche Patienten sind nun aber gegen das Heftpflaster sehr empfindlich; sie bekommen leicht Eczeme, und es muss dann der Verband alsbald abgenommen werden. Bei solchen Patienten thut man gut, zunächst zwei Streifen von Unna'schem Zinkpflastermull aufzukleben und das Heftpflaster dann erst auf diese Streifen aufzulegen. Die Haltbarkeit des Verbandes bleibt so durchaus gewahrt; Eczeme aber vermeidet man fast sicher.

Eine andere Art und Weise, um bei empfindlichen Patienten doch einen Extensionsverband anbringen zu können, ist die v. Volkmann'sche Methode des „Stecknadelverbandes". Dieser ist sehr leicht anzulegen. Man legt, gerade wie sonst die Heftpflasterstreifen, jetzt eine einfache Leinwandbinde schlingenförmig an dem Bein an, wickelt diese Schlinge nun mit einer anderen Binde sorgfältig an, indem man über den Malleolen beginnt und die Binde kunstgerecht bis zur Leiste hin führt, und steckt schliesslich jede einzelne Tour dieser letzteren Binde an der unterliegenden Lein-

wandbinde mit je einer Stecknadel fest. Hat man gut gewickelt, so kann man schon einen sehr kräftigen Zug ausüben, ohne dass sich der Verband auch nur im mindesten verschiebt. (Tafel 133 b.)

Heusner hat einen sehr praktischen Ersatz des Heftpflasters angegeben. Er verwendet Filzstreifen und aufgelöste Heftpflastermasse (Cerae flavae, Resinae Damarae, Colophon. āā 10,0. Terebinth. 1,0. Aether, Spirit. Ol. therebinth. āā 55,0, filtra!). Diese letztere Flüssigkeit kann in einer verstöpselten Medizinflasche beliebig lange aufbewahrt werden. Mit Hilfe eines eingesetzten Zerstäubungsröhrchens, welches mit dem Munde angeblasen wird, bestäubt man ganz leicht das Bein oder den Arm zu beiden Seiten, legt dann über den bestäubten Teilen je einen etwa handbreiten Filzstreifen, welcher an der Aussenseite mit fester Leinwand übernäht ist, und wickelt nun die beiden Filzstreifen mit einer gewöhnlichen Mullbinde fest an. Ueber diese Mullbinde wickelt man dann noch eine steife Gazebinde recht fest und gleichmässig. Unmittelbar nach Fertigstellung des Verbandes kann man die Extension mit den schwersten Gewichten ausüben. Eczeme entstehen nicht, wenn man die Heftpflastermasse nicht zu dick aufstäubt. Braucht man den Verband nicht mehr, so lassen sich die Filzstreifen ganz leicht vom Gliede abziehen; sie können dann gut noch einigemal verwertet werden. Die geringe Klebrigkeit der Haut, die nach dem Abnehmen der Filzstreifen bestehen bleibt, beseitigt man leicht durch Abseifen und Waschen der Haut.

Der weiche, etwa $^3/_4$ cm dicke Filz, sog. Klavierfilz ist zu beziehen aus der Dittersdorfer Filz- und Kratzentuchfabrik in Dittersdorf (Sachsen).

In den letzten Jahren hat man die permanente Extension mit Erfolg auch zur Behandlung der Frakturen an der oberen Extremität verwertet. Als Beispiel bilden wir die Methode von Clarc ab und beschreiben die Anlegung eines Zugverbandes am ganzen Arme.

Tafel 134.

Clarc's permanenter Extensionsverband.

Tafel 135.

Clarc's permanenter Extensionsverband.

Clarc geht in folgender Weise vor: (Tafel 134.) Während der Kranke geht, steht oder sitzt, ruht der rechtwinklig gebeugte Vorderarm in einer Schlinge, und vom Oberarm hängt frei herab das mit zwei Längsstreifen und einer Bindeneinwickelung befestigte Gewicht, welches bei horizontaler Lage über eine Rolle geführt wird.

Will man, wie bei den Brüchen im oberen Drittel des Humerus, den ganzen Arm extendieren, dann legt man den Patienten ins Bett, befestigt an der Volar- und Dorsalseite des Armes in horizontaler Lage einen Heftpflasterstreifen, der in einer Schlinge endigt und über ein Brettchen geht, an dem das Gewicht, welches über eine Rolle geführt wird, befestigt wird. Die Streifen werden durch circuläre Touren am Arme befestigt, ganz nach Analogie der Extension am Oberschenkel. Die Gegenextension findet am Thorax statt. (Tafel 135.)

b) Zug mittelst eines zusammengelegten Tuches.

Man kann mittelst eines in Kravattenform angeordneten Tuches eine sehr kräftige Extension ausüben. Hennequin hat eine Methode ausgebildet und für die Extensionsbehandlung von Oberschenkelbrüchen verwertet. Die Methode wird in Frankreich sehr häufig verwertet. Man geht in der Weise vor, dass man das Kniegelenk in einem Winkel von etwa 40 Grad beugen lässt. Dann umwickelt man das untere Ende des Femur mit Watte und legt nun über die Watte ein kravattenförmig gefaltetes Tuch in der Weise an, dass die Mitte desselben auf die Vorderseite des Oberschenkels oberhalb des Knies zu liegen kommt. Die beiden Zipfel führt man dann kreuzweise über die Kniekehle und den hinteren oberen Teil des Unterschenkels hinweg nach vorn und knotet sie vor dem Unterschenkel. Von der geknoteten Stelle aus kann man nun, ohne dass das Tuch abgesteckt oder in der Kniekehle drückt, einen sehr kräftigen Zug ausüben. (Tafel 136.) Um den Zug permanent zu machen, wird an die Knotenstelle eine Schnur gebunden; diese wird über eine Rolle geleitet,

Tafel 136.

Extensionsmethode nach Henneguin.

Tafel 137.

Gehverband von F. Krause.

Tafel 138.

Gehverband von F. Krause.

Tafel 139.



a) Taylor'sche Extensionsschiene. b) Thomas'sche Schiene.
c) Heusner'sche Schiene.

Schraubenschienen verlängern lassen und es dadurch ermöglichen, die beiden Gipskapseln allmählich mehr und mehr von einander zu entfernen. (Tafel 133c.)

Zu den hierher gehörigen Verbänden ist auch der sog.

Gehverband

zu zählen.

Die Gehverbände sind zu dem Zwecke angegeben worden, um bei frischen Frakturen den Patienten sofort das Gehen zu gestatten. Wir beschreiben die Methode nach den Vorschriften von F. Krause, der die Methode bei uns eingeführt und im wesentlichen ausgebildet hat.

Soll der Patient sich schon wenige Tage nach der Verletzung auf sein gebrochenes Glied stützen können, so muss sich der Gipsverband so genau allen Umrissen des Gliedes anschmiegen, dass er dessen Form wie im Modell wiedergibt. Dabei darf er natürlich nirgends drücken; es erfordert daher das Anlegen eines solchen Verbandes grosse Uebung.

Man wickelt die Gipsbinde ohne jede Wattepolsterung am besten über einem Trikotstrumpf oder auch wohl über einer doppelten Lage einer Mullbinde an dem Beine exakt an. Während des Abwickelns der feuchten Gipsbinden muss jedes Zerren vermieden werden, auch thut man gut, bei mangelnder Uebung mit den ersten Binden keine Umschläge zu machen, sondern sie lieber durchschneiden zu lassen. Durch sorgsames Streichen werden während des Umlegens der Binden die einzelnen Schichten in innige Berührung gebracht und genau der Form des Gliedes angeschmiegt. Der Sohlenteil wird durch Longuetten verstärkt. Der Verband reicht bei leichten Malleolarbrüchen von den Köpfchen der Mittelfussknochen bis zu den Condylen der Tibia, an denen er eine gute Stütze findet. (Tafel 137.) Bei Oberschenkelbrüchen muss das Becken in den Verband mit hereingenommen werden, es ist darum notwendig, den Verband besonders an der Gesässfalte gut zu modellieren, so dass der Patient gewissermassen mit dem Tuber ischii auf dem Verbande reitet. (Tafel 138.)

Der Verband gibt genau die Contouren des Beines wieder, wenn man nur überall ungefähr gleich viele Bindenschichten übereinander legt. Man kann daher auch im Verbande sehr gut die Stellung der Bruchenden beurteilen.

Das Kniegelenk wird in ganz leichte Beugung gebracht, der Fuss muss vollkommen rechtwinklig zum Unterschenkel, eher noch in leichter Dorsalflexion, ferner in mittlerer Lage zwischen Supination und Pronation, eher etwas mehr in Supination stehen.

Der Verletzte bleibt nach Anlegung des Gipsverbandes noch ein- bis zweimal 24 Stunden im Bett. Wird nach dieser Zeit alles in Ordnung befunden, so soll der Kranke unter Benutzung des verletzten Gliedes umhergehen. Hierbei ist man etwas von der Willenskraft der betreffenden Leute abhängig. Manche gehen sofort an zwei Stöcken oder am v. Volkmann'schen Bänkchen, andere müssen zunächst gestützt werden, bis sie das ängstliche Gefühl, sie könnten sich Schaden thun, überwunden haben. Mit Geduld bringt man aber auch schüchterne Gemüter dazu, fest auf das zerbrochene Glied aufzutreten. Ein Zwang wird nie ausgeübt, die Kranken überzeugen sich sehr schnell selbst von den Vorteilen der ambulanten Behandlung.

Empfinden sie an der Bruchstelle überhaupt Schmerzen, so sind diese gewöhnlich nicht erheblich und pflegen bei fortgesetztem Gebrauch abzunehmen. Die Zehen schwellen beim Umhergehen zuweilen an, was ja auch schon beim Hängenlassen des verletzten Beines eintreten kann; das Oedem verschwindet aber, wenn der Verband nirgends einen schädlichen Druck ausübt, in der Rückenlage. Daher wird der Kranke angewiesen, jedesmal wenn er sich setzt, das verletzte Glied hochzulegen. Man kann die Leute mit einem passenden Schuh sehr gut auch auf der Strasse umhergehen lassen.

Bei einfachen Frakturen könnte der Verband bis zur vollendeten Konsolidation liegen bleiben. Es ist indessen doch besser, ihn in etwa 14 Tagen zu wechseln um zu sehen, ob die Dislokation auch wirklich völlig usgeglichen ist.

Tafel 140.



a) Liermann'sche Schiene.
b) Hessing'scher Schienenhülsenapparat.

c) Der Zug durch Schienen und Apparate.

Der Zug unter Vermittlung von Schienen wurde von den älteren Autoren dadurch hergestellt, dass dieselben Schienen, in der Regel aus Holz, an der Aussenseite der Extremität anbrachten und nun mit Hilfe von Schlingen das centrale und periphere Ende des Gliedes gegen die das Glied an beiden Enden überragende Schiene heranzuziehen suchten. Diese Extensionsschienen belästigen dann in der Regel den Patienten durch schmerzhaften Druck mehr als sie nützten. Erst in neuerer Zeit sind fortschreitende mit der Technik brauchbare Schienen konstruiert worden, und diese finden denn jetzt auch, namentlich bei der Behandlung der Frakturen und Deformitäten der unteren Extremitäten ausgiebige Verwendung.

Die bekanntesten der Extensionsschienen sind folgende:

1. Die Taylor'sche Schiene; sie besteht aus einem Beckengürtel mit zwei Perinealgurten und einer äusseren Extensionsschiene, welche durch ein Zahnrad und Triebschlüssel beliebig verlängert werden kann. Die Extensionsschiene biegt sich unten rechtwinklig um und bildet so eine, von der Fusssohle 2—3 cm abstehende, mit Gummi gepolsterte Gehfläche. An dieser Gehfläche befinden sich zwei Riemen. Man klebt nun zu beiden Seiten des zu extendierenden Beines handbreite, unten mit einer Schnalle versehene Heftpflasterstreifen an, schnallt den Beckengürtel und die Perinealgurte zu und schnallt schliesslich die Riemen der Gehfläche des Apparates an die Schnallen des Heftpflasters an. Den Oberschenkel umfasst eine Lederkappe. Da der Apparat den Fuss überragt, so muss die Stiefelsohle der gesunden Seite entsprechend erhöht werden. (Tafel 139a.)

2. Die Thomas'sche Schiene besteht aus einem Sitzring, auf dem der Patient reitet, und zwei Seitenschienen, die unten rechtwinklig umbiegen und vereinigt einen Steigbügel bilden. Mittelst eines Extensionszügels wird der Fuss gegen die Steigbügelplatte angezogen. (Tafel 139b.)

3. Die Heusner'sche Schiene besteht aus zwei an der Knie- und Knöchelgegend mit Scharnieren versehenen Seitenschienen, die oben einen Sitzring tragen und u[illegible] an einer Sohle aus Stahlblech befestigt sind. [illegible]atte und Seitenspangen sind mit weichem Filz ge[illegible]t. Der Sitzring, ebenfalls mit Filz gefüttert, wird [illegible]eichem Leder überzogen. (Tafel 139c.)

[illegible]. Liermann'sche Schiene ist eine Modi[illegible]ation zuerst von Harbordt erfundenen Schiene. [illegible]die innere Seite des Beines angelegt. Die [illegible]schieht, wie die Abbildung zeigt, mittelst [illegible]hraube, die nach Erzielung des erforderlichen [illegible]ntfernt werden kann. (Tafel 140.)

[illegible]uns'sche Schiene ist diejenige, welche [illegible]m [illegible] wohl die besten Dienste leisten wird. [illegible]e [illegible]h eine Geh- und eine Lagerungsschiene [illegible]nd b[illegible]n wesentlichen aus zwei seitlichen Stäben, einem Sitzringe und einem Steigbügel. Das Bein ruht hinten auf einigen breiten Leinwandstreifen, die zwischen den für verschiedene Grössen einstellbaren Stäben ausgespannt sind; vorne geschieht die Befestigung mit einigen schmalen Gurten. Nehmen wir einen Fall von Schrägbruch im oberen Drittel des Oberschenkels an, so wird die Schiene zunächst als Lagerungsapparat verwendet, um für zwei Wochen zuerst eine kräftige Extension ausüben zu können. An das Bein werden beiderseits Heftpflasterstreifen angeklebt, die durch Vermittlung von Bändern gegen den Steigbügel angezogen werden. An den Steigbügel wird ein Fussbrett befestigt, welches eine Rolle zum Ueberleiten der Gewichte trägt. (Tafel 141.)

Wie bei der gewöhnlichen Gewichtsextension werden nun die Gewichte angehängt. Nach etwa 2 Wochen wird die Extension entfernt, ein leichter Gipsverband von den Zehen bis zur Leiste umgewickelt und darüber die Schiene wieder angelegt. Nunmehr fällt das Fussbrett weg; der Steigbügel wird so gestellt, dass er etwas von der Fusssohle absteht, das Bein wird durch die Streifen gegen den Steigbügel gezogen, die Contra-

Tafel 141.



P. Bruns'sche Extensionsschiene und Lagerungsapparat.

extension aber dadurch gebildet, dass der Sitzring sich gegen das Tuber ischii anstemmt.

Jetzt vermag der Patient in dem Apparat umherzugehen und ist somit nicht mehr an das Bett gefesselt. (Tafel 141.)

6) Unbedingt die beste aller Extensionsschienen ist der Hessing'sche Schienenhülsenapparat. Wie der Name sagt, besteht der Apparat aus Hülsen, die über einem Modelle der Extremität aus Leder gefertigt werden und an den Hülsen festgeschraubten, aber leicht verstellbaren Schienen. Die Extension geschieht dadurch, dass die Schienen erst festgeschraubt werden, während der Ober- und Unterschenkel, sowie der Fuss manuell bei maximaler Streckung in die Hülsen hineingelegt worden sind. Diese Extensionstellung wird dann dadurch aufrecht erhalten, dass sich die wohlgepolsterte Oberschenkelhülle gegen das Tuber ossis ischii anstemmt, während der Fuss durch einen sog. Fersenzug fest gegen den geschmiedeten Sohlenteil der Fusshülse, das Fussblech angedrückt wird. In der Regel verbindet man den Apparat noch mit einem festsitzenden Beckengürtel. (Tafel 140 b.)

d) **Druckverbände.**

Druckverbände werden vielfach verwendet, teils zur Beseitigung chronischer Entzündungsprodukte, teils zur Beseitigung von ödematösen Anschwellungen der Glieder, weiterhin zur Bekämpfung variköser Venenerweiterungen, zur Beseitigung von Gelenkergüssen u. s. w.

Bei jedem Druckverbande soll man mit der Einwickelung des Gliedes an der Peripherie desselben beginnen und die Binden nach dem Centrum hin unter gleichmässigem Druck so anlegen, dass jede folgende Tour die vorhergehende mindestens zu zwei Dritteln deckt.

An der unteren Extremität erstrebt man einen gleichmässigen Druck vielfach durch Verwendung von Gummibinden. Diese Gummibinden soll man nicht auf die blosse Haut legen, sondern vielmehr als Unterlage

stets erst eine Mullbinde benützen. Man darf die Binden ja nicht zu fest anlegen, weil sonst Zirkulationsstörungen entstehen.

Einen gleichmässigen Druck auf die untere Extremität erzielt man auch durch Anwendung von Gummistrümpfen, von denen die von M. Senftleben in Vegesack fabrizierten nahtlosen Patent-Gummistrümpfe unbedingt am meisten zu empfehlen sind, da sie nicht die Hautausdünstung verhindern und beliebig gewaschen werden können.

Druckverbände für die Gelenke zur Beförderung der Resorption von Gelenkergüssen legt man so an, dass man zunächst das ganze Glied von der Peripherie bis zum Centrum einwickelt, dass man dann um das Gelenk selbst zuerst eine Flanellbinde und über diese noch eine Gummibinde anwickelt. Ein starker Druck ist zu vermeiden. Am besten wirkt ein mässiger, aber anhaltender Druck.

Zur forcierten Kompression des Kniegelenkes nach v. Volkmann bei Hydrops des Gelenkes oder bei Bursitis praepatellaris legt man zuerst die schon beschriebene Holzschiene in die Kniekehle und wickelt dann das Knie fest mit Binden ein. Der Verband bleibt 2—3 Tage liegen, auch wenn sich Oedeme bilden und lebhafte Schmerzen entstehen.

Eine andere Art der elastischen Kompression ist die mittelst Schwämmen ausgeübte. Geeignete Badeschwämme von entsprechender Form, die gut gereinigt, in warmes Wasser getaucht und wieder gut ausgedrückt sind, werden auf die zu drückende Körperstelle aufgelegt und mit Binden fest angewickelt. Zur Erzielung eines lokalisierten Druckes kann man wohl auch einen Gipsverband anlegen, an der betreffenden Stelle ein Fenster ausschneiden, dieses mit keilförmigen dicken Schwammstücken ausfüllen und darüber eine Gummibinde fest anwickeln.

Vielfach bedient man sich des Heftpflasters zur Kompression.

Tafel 142.

a) Fricke'scher Heftpflasterverband bei Orchitis und Periorchitis.
b) Baynton'sche Heftpflastereinwickelung bei Ulcus cruris.

So sind z. B. recht beliebt die Fricke'schen Heftpflastereinwickelungen (Taf. 142 a) bei der akuten Orchitis und Periorchitis. Man isoliert den kranken Hoden, indem man ihn mit der linken Hand umfasst und vom anderen Hoden abdrängt, sodass er eiförmig absteht. Nun legt man zunächst an der Wurzel des abgedrängten Hodens unter leichtem Druck einen etwa fingerbreiten Heftpflasterstreifen cirkulär um. Von dieser ersten Tour aus legt man nun etwa fingerbreite Längsstreifen aus Heftpflaster dachziegelförmig so über den Hoden, dass die Streifen von einem Punkt der ersten Tour ab über den Scheitel des Hodens fort- und nach dem entgegengesetzten Punkte der anderen Seite hinlaufen. Man legt so viele Streifen an, dass der Hoden völlig abgeschlossen ist. Schliesslich legt man zur Befestigung der Längsstreifen noch eine Kreistour an der Wurzel des Hodens an, welche die erste Kreistour deckt, Schwillt der Hoden ab, so dass der Verband locker wird; so wird ein neuer Verband angelegt.

Ein weiterer zweckmässiger, mittelst Heftpflasterstreifen (Tafel 142 b) auszuführender Druckverband ist der Baynton'sche Heftpflasterverband, den man braucht, wenn man die Ueberhäutung granulierender Flächen, namentlich von Geschwüren, beschleunigen will. Man legt die Mitte eines etwa fingerbreiten Heftpflasterstreifens an der dem Geschwür entgegengesetzten Seite des Gliedes an, nimmt jedes Ende des Streifens in eine Hand und kreuzt nun beide Streifen mit kräftigem Zuge auf der granulierenden Fläche. Die Streifen müssen etwa $1^1/_2$mal so lang sein als der Umfang des Gliedes beträgt. Man beginnt am peripheren Rand der zu deckenden Fläche und legt nun die einzelnen Streifen aufsteigend so an, dass sie sich dachziegelförmig zu etwa $^2/_3$ decken. Der Verband muss die ganze Geschwürsfläche decken; über das Heftpflaster legt man eine dicke Schicht Watte und wickelt das ganze mit einer Binde gut ein. Der Verband bleibt 2—5 Tage liegen und wird dann erneuert; er ist von ausgezeichneter Wirkung.

Schliesslich wollen wir noch den ebenfalls recht brauchbaren Gibney'schen Heftpflasterverband zur Behandlung der Distorsionen des Fussgelenkes beschreiben. (Tafel 143.)

Nehmen wir an, es handelte sich um eine schwere Distorsion des Fussgelenkes mit Zerreissung der Bänder an der äusseren Seite des Gelenkes, so wird der Verband, wenn der Patient alsbald nach geschehener Verletzung in unsere Behandlung tritt, sofort angelegt. Ist dagegen schon längere Zeit nach dem Unfall verstrichen und ist dann, wie gewöhnlich, eine stärkere Schwellung vorhanden, so wickelt man den Fuss mit einer Flanellbinde ein, legt über diese noch eine Gummibinde und lässt den Fuss dann für 24 Stunden hochlegen. Man verbindet also während dieser Zeit die Elevation des Fusses mit einer elastischen Kompression desselben. Nach Ablauf der genannten Frist wird der Heftpflasterverband angelegt, nachdem vorher noch, wenn nötig, die restierende Schwellung durch Massage nach Möglichkeit beseitigt worden war.

Zur Anlegung des Verbandes bedient man sich des gut klebenden, als Mead's adhesive plaster bekannten, amerikanischen Heftpflasters. Von dem Heftpflaster schneidet man sich zweierlei Streifen, längere und kürzere, zurecht und hängt die Streifen, damit sie nicht aneinander kleben, über eine Stuhllehne. Die Streifen erst während des Anlegens des Verbandes abschneiden zu wollen, ist unzweckmässiger, weil man dann später einen Assistenten mehr braucht. Das Mass für die längeren Streifen bestimmt man so, dass man eine Schnur von der Grenze des mittleren und oberen Drittels des Unterschenkels aus, an der Aussenseite desselben herab und über die Fusssohle auf den Fussrücken bis zur Höhe des gegenständigen Knöchels hinführt. Das Mass für die kürzeren Streifen nimmt man ebenfalls mit der Schur, indem man dieselbe von der Basis der kleinen Zehe längs des äusseren Fussrandes und längs des inneren Fussrandes bis zur Basis der grossen Zehe verlaufen lässt. Die Breite der Streifen

Tafel 143.

Gibney'scher Heftpflasterverband bei Distorsio pedis.

ist etwa die eines erwachsenen Daumens. Man braucht im Durchschnitt 10 längere und 10 kürzere Streifen.

Sowohl beim Nehmen des Masses als namentlich beim Anlegen des Verbandes selbst, ist es unbedingt notwendig, den Fuss genau in rechtwinkliger Stellung zum Unterschenkel halten zu lassen; denn nur bei einer solchen Steilung des Fusses kann der Patient gut gehen; jede auch noch so geringe Spitzfussstellung hindert die Abwickelung des Fusses vom Boden und verursacht damit dem Patienten Beschwerden.

Das Anlegen der Streifen geschieht nun folgendermassen: Während ein Gehilfe den Fuss, wie gesagt, genau in rechtwinkliger Beugestellung hält, klebt man selbst den ersten langen Streifen entsprechend der Tibiakante an der Grenze des oberen und mittleren Drittels des Unterschenkels an. Der Patient selbst drückt mit seinen Fingern den Streifen fest an. Der Arzt aber spannt den Streifen an, führt ihn längs der Tibiakante gerade herunter, legt ihn steigbügelartig um die Fusssohle herum und klebt sein Ende auf der inneren Seite des Fussrückens in der Höhe des Malleolus internus und fingerbreit vor demselben, entsprechend etwa der Sehne des Extensor hallucis longus an. Der Streifen muss straff angespannt sein, wenn er richtig liegt. Dieser erste lange Streifen wird nun am Fuss durch den ersten kurzen Streifen fixiert. Dieser letztere wird am äusseren Fussrand, an der Basis der kleinen Zehe beginnend, angeklebt, um die Hacke herum und bis zu der Basis der grossen Zehe hingeführt und fest angeklebt. Liegen diese zwei ersten Streifen, so werden nun die übrigen langen und kurzen Streifen diesen beiden ersten Streifen parallel so angelegt, dass sich die einzelnen Streifen stets dachziegelförmig zur Hälfte decken, so lange bis die ganze Gegend des Malleolus externus bis zum Rande der Achillessehne mit dem Heftpflaster bedeckt ist.

Wenn man dann will, kann man zur Verstärkung des Verbandes noch einige diagonale Streifen anlegen. Unbedingt nötig ist dies aber nicht.

Handelt es sich um eine Distorsion, bei der vorzüglich die inneren Bänder des Fussgelenkes gelitten haben, so werden die Heftpflasterstreifen anstatt auf der äusseren, auf der inneren Seite angelegt, bis die Gegend des inneren Malleolus völlig bedeckt ist.

Ist vorzugsweise das Mittelfussgelenk geschädigt, so legt man die Touren entsprechend dem früher beschriebenen Steigbügelverband an. Man beginnt z. B. mit dem ersten Streifen an der Innenseite der Hacke, führt denselben um die Hacke herum auf und quer über den Fussrücken bis zur Basis der grossen Zehe, wo er an der Sohlenfläche endigt. Der zweite Streifen geht in derselben Weise vom Malleolus externus um die Hacke herum zur Kleinzehenseite und endigt an der Sohle unterhalb der kleinen Zehe. Die folgenden Streifen werden dann, ebenfalls sich zur Hälfte dachziegelförmig deckend, angeklebt, bis der ganze Fuss von den Zehengelenken bis zur oberen Grenze des unteren Drittels des Unterschenkels eingewickelt ist.

In jedem Falle muss eine völlige cirkuläre Einschnürung des Fussrückens vermieden werden; Cirkulationsstörungen dürfen nicht entstehen. Liegen alle Streifen, so wickelt man über dieselben eine Cambricbinde und eventuell noch eine steife Gazebinde, und nun kann der Patient sofort seinen Strumpf und Schuh überziehen. Die Patienten können nun selbst mit schweren Distorsionen direkt vom Fleck weggehen, indem sie sich anfangs noch eines Stockes bedienen. Bald aber lernen sie es, auch ohne jede Unterstützung ihren Berufsgeschäften wieder nachzugehen.

Bei leichten Distorsionen genügt ein Verband. Man lässt denselben etwa 8 Tage liegen und massiert dann noch für weitere 8 Tage täglich einmal den Unterschenkel.

Bei schweren Distorsionen mit starker Schwellung und Ecchymosenbildung erneuert man den Verband nach 6—8 Tagen, lässt diesen zweiten Verband noch weitere 8—10 Tage liegen und massiert dann ebenfalls noch etwa eine Woche lang. Einen dritten Verband anzulegen ist man wohl nie genötigt.

Register.

A.

Abwickeln der Binden 8.
Achtertour, Petit'sche 28.
Ahornfourniere 71.
Albers'scheKragenschienen103.
Alembrotsalz 51.
Aluminiumschienen 80.
Antiseptische Verbände 53.
Armtuch s. Mitella.
Aseptische Verbände 50.
Aufwickeln der Binden 2.
Augen, Bindenverbände für die 25
— , Tuchverband für die 46.

B.

Baynton'scher Heftpflasterverband 125.
Becken, Wundverbände am 59.
Beckenstützen 96.
Beely'sche Gipshanfschienen 99.
Beinschiene, Sharp'sche 72.
Billroth battist 60, 62.
Binden, Abwickeln der 8.
— , Aufwickeln der 2.
— blaue 1.
— einköpfige 1.
— zweiköpfige 3,
Bindenverbände 1.
Bindenwickelmaschiene 2.
Binoculus 26.
Blechschienen 75.
Blechstreifenschienen, Port'sche 77.
Bleischienen 78.
Blutschorf, feuchter nach Schede 55.
Bonnet'sche Drahthose 67.
Borlint 51.
Braatz'sche Gipstricotschienen 102.
Breiger'sche Gipswatteschienen 104.
Bruns'sche Extensionsschiene 122.
— Schiene 76.
Brust Bindenverbände an der 28.
— , Tuchverband 46.
— Wundverbände 57.

C.

Calicotbinden 1.
Cambricbinden 1.
Capistrum duplex 24.
— simplex 22.
Capitium quadrangulare 45.
— triangulare 44.
Carr'sche Vorderarmschiene 69.
Celluloid-Mullverband 110.
Celluloid-Plattenverbände 110.
Celluloid-Tricotverband 110.
Cellulose-Verband, geleimter 108.
Chirotheca completa 11.
— dimidia 10.
Clarc's Extensionsverband 117.
Clavikularfrakturen, Verbände für 32.
Collodium-Verbände 47.
Compressen 3.
Compressionsverbände 123.
Compressorium mammae 31.
Contentivverbände 67.
Contraextension 114.
Cooper'sche Vorderarmschiene 69.
Cramer'sche Drahtschienen 82.

D.

Damm, Wundverbände am 61.
Dauerverbände 53.
Daumen, Verbände am 10.

Desault'sches Kissen 63.
— 'scher Verband 32.
Distraktionsverbände 111.
Dolabra currens 5.
— repens s. serpens 5.
— reversa 7.
Dorsalschiene, Roser'sche 69.
— , Volkmann'sche 70.
Drahthosen 67.
Drahtrinnen 67.
Drahtrollbinden, Port'sche 83.
Drahtschienen 82.
Drahtsiebstoffschienen 83.
Druckverbände 123.
Dulles'scher Verband 37.
Dumreicher'sches Kissen 64.
Dupuytren'sche Schiene 70.

E.

Ebert's Rollenträger 113.
Eisenblechschienen, englische 77.
Ellbogen, Bindenverbände am 12.
— , Tuchverband am 40.
Erhärtende Verbände 87.
Esmarch'sche Drahtsiebstoffschienen 83.
— Resektionsschienen 72.
— scher schneidbarer Schienenstoff 72.
Extension durch Contentivverbände 118.
Extensionsschienen 121.
Extensionsverbände 112.

F.

Fascia circularis 4.
— nodosa 27.
Fialla'sche Stäbchenbeinlade 65.
Fiberschienen 86.
Filzschienen 84.
Filzstreifen, Heusner'sche zu Extensionsverbänden 116.
Fingerling, Neff'scher 47.
Fingerverbände 10.
Flanellbinden 1.
Flügelschiene, Volkmann'sche 70.
Fricke'sche Heftpflastereinwicklung 125.
Funda 3.
— maxillae 28.
— nasi 28.
Fuss, Bindenverbände am 17.
— , Tücherverbände am 43.
Fussgelenk, Gibney'scher Verband bei Distorsionen im 126.

G.

Gaze 54.
— sterile 55.
Gazebinden 55.
— appretirte 1.
Geh-Verband 119.
Gelenkschienen 95.
Gelenkverbände, artikulirt-mobile 106.
Geranium 9.
Gibney'scher Heftpflasterverband 126.
Gipsbindenmaschiene, Beely'sche 89.
Gipsbindenverband 88.
Gipshanfschiene, Beely'sche 99.
Gipshülsen, abnehmbare 95.
Gipskataplasmen 104.
Gipslonguetten 103.
Gipsmesser 94.
Gipssägen 94.
Gipsscheren 94.
Gipsschienen 99.
Gipstricotschienen, Braatz'sche 102.
Gipsverband 87.
— abnehmbarer 95.
— gefensterter 95.
— unterbrochener 95.
Gipswatteschienen, Breiger'sche 104.
Gooch'sche Spaltschienen 71.
Grosse Beckenstützapparat 96.
Gummibinden 123.
Gummistrümpfe 124.
Guttapercha-Schienen 85.
Gips 87.

H.

Hand, Tuchverband für die 40.

Stäbchen-Beinlade, Fialla'sche 62.
Stapes 16.
Stecknadelverband 103.
Stehbett, Phelps'sches 61, 66.
Steigbügel s. Stapes.
Stella dorsi 28.
— pectoris 29.
— pectoris et dorsi 29.
Sterilisierung der Verbandstoffe 53.
Sternbinde 29.
Stirnband 45.
Stromeyer'sche Armschiene 70.
— Handbretter 69.
— Kissen 63.
Stützhäkchen 39.
Sublimatgaze 53.
Supinationsschiene, v. Volkmann'sche 72.
Suspensionsschiene, v. Volkmann'sche 72.
Suspensorium mammae 30.
— — duplex 31.

T.

Tamponade 53.
Taylor'sche Schiene 121.
T-Binde 3.
Testudo calcanei 16.
— cubiti 12. 13.
— genu 18. 19.
Thomas'sche Schiene 121.
Thymol 50.
Torfmoos 55.
Tuchschienen, Schnyder'sche 71.
Tücherverbände 39.
Triangel, Middeldorpf'scher 63.
Trikotschlauch 90.
T-Schiene, v. Volkmann'sche 75.

U.

Universalzange 78.
Unna'scher Pflastermull 115.
— Zinkleimverband 49.
Untere Extremität, Bindenverbände für die 16.
— Tücherverbände für die 42.
Unterleib, Wundverband am 58.
Unterschenkel, Verbände am 18.

V.

Verbände, antiseptische 53.
— aseptische 50.
— einfache 1.
— erhärtende 87.
Verbandgaze 3.
Verbandschere 8.
Verbandwatte 3.
Velpeau'scher Verband 34.
Volk'scher Holzspangipsverband 97.
v. Volkmann'sche Dorsalschiene 70.
— Flügelschiene 70.
— Kniceschiene 70.
— Supinationsschiene 72.
— T-Schiene 75.
Vorderarm, Verbände am 4.

W.

Wange, Tuchverband für die 45.
Wasserglasverband 104.
Watson'sche Schiene 70.
Wickelverband für den Kopf 57.
Wolfsmaul 94.
Wundverbände 46. 50. 56. 62.

Z.

Zinkblechschienen 76.
Zinkleimverband 49.
Zirkeltour 4.
Zugverbände 111.

Verlag von J. F. LEHMANN in MÜNCHEN.

Anatomie.

Arbeiten aus dem anatomischen Institute zu München. Herausgeg. von **K. v. Kupffer und N. Rüdinger.** (Münchener med. Abhandlungen VII. Reihe). 8°.

Heft 1: Utschneider, A., Lendennerven der Affen und der Menschen. 1892. 32 S. Mit 1 Tafel. M. **1.**—.
Heft 2: Tettenhammer, Ueber offene Schlundspalten bei e. menschl. Embryo. 34 S. Mit 12 Abbild. M. **1.**—.
Heft 3: Höfer, W., Vergleichend-anatomische Studien über die Nerven des Armes und der Hand beim Menschen und bei dem Affen. 1892. 106 S. Mit 6 Tafeln. M. **4.**—.
Heft 4: Kupffer, K. v., Ueber die Entwicklung von Milz und Pankreas. 17 S. Mit 7 Abbildungen. 1892. M. **1.**—.
Heft 5: Kupffer, K. v., Ueber das Pankreas bei Ammocoetes. 24 S. mit 7 Abbildungen. 1893. M. **1.**—.
Heft 6: Kuithan, D. W., Die Entwicklung des Kleinhirns bei Säugetieren. 1895. 40 S., 24 Abbild. M. **1.60.**

Boegle, K., Die Entstehung und Verhütung von Fussabnormitäten. 139 Seiten. Mit 39 Abbildungen. 1893. Broschiert. M. **4.**—.

Boegle, Dr. K., Die Entstehung organischer Formen. 1895, gr. 8°. 18. S. Text mit 40 Abbildungen. M. **1. 20.**

Herzog, Privatdoc. Dr. W., Die Rückbildung d. Nabels und d. Nabelgefässe, m. Berücksichtig. d. Pathogenese der Nabelhernien. Eine anat.-histol. Unters. 1892. gr. 8°. 48 S. m. 8. T. M. **6.**—

Kupffer, K. v., Studien zur vergleichenden Entwicklungsgeschichte des Kopfes der Kranioten.

Lieferung 1.: Entwicklung des Accipenser Sturio. Mit 10 lithograph. Tafeln. Gr. 8°. 1893. Broschiert. M. **10.**—.
Lieferung 2.: Entwicklung des Kopfes von Ammocoetes Planeri. Mit 12. lithogr. Tafeln. Gr. 8°. 1894. M. **10.**—.
Lieferung 3.: Entwicklung der Kopfnerven von Ammocoetes Planeri. 1895. Mit 48 Abbild. Gr. 8°. M. **8.**—.

Das ganze Werk erscheint in zwanglosen Heften; jährlich gelangt 1 Heft zur Ausgabe.

Jedes Heft bildet für sich ein abgeschlossenes Ganzes.

Plessen, J. v. und **J. Rabinowicz,** Die Kopfnerven von Salamandra maculata im vorgerückten Embryonalstadium. 20 Seiten mit 4 colorierten Tafeln. 4°. 1891. Brosch. M. **5.**—.

Schäffer, O., Untersuchungen über die normale Entwicklung der Dimensionsverhältnisse des fötalen Menschenschädels mit bes. Berücksichtig. d. Schädelgrundes und s. Gruben. 4°. 1893. 51 S. mit 50 Abbildungen und Tabellen. M. **7.**—.

Dr. A. Schmitt, Die Fascienscheiden und ihre Beziehungen zu Senkungsabscessen. 122 S. 8°. 2 Tafeln. M. **4.**—.

Verlag von J. F. LEHMANN in MÜNCHEN.

Hygiene.

Arbeiten aus dem hygienischen Institut in München. Herausgegeben von **Geheimrat Prof. Dr. Max v. Pettenkofer.** (Münchener medicin. Abhandlungen V. Reihe). gr. 8°.

Heft 1: Die Schwemmkanalisation in München. Von Max von Pettenkofer. 16 Seiten. 1891. ℳ 1.—.

Heft 2: Die Fehlböden (Zwischendecken). Ihre hygien. Nachteile u. deren Vermeidung. v. Dr. Heinzelmann. 1891. ℳ 1.—.

Heft 3: Acht Thesen gegen die Münchener Schwemmkanalisation. Bespr. von Max v. Pettenkofer. 1892. 22 S. ℳ 1.—.

Heft 4: Ueber Cholera m. Berücksichtig. d. jüngst. Choleraepidemie in Hamburg. Von Max v. Pettenkofer. 1892. ℳ 1.—.

Heft 5: Cholera-Explosionen und Trinkwasser von Max v. Pettenkofer. 3 Bogen Text u. 6 graph. Tafeln. ℳ 1.—.

Bouček, Dr. B., Die Cholera im Bodebrader Bezirke. Eine epidemiologische Studie. Mit 48 S. Text u. 41 Plänen. ℳ 2.—.

Däubler, Dr. C., Die Grundzüge der Tropenhygiene. 123 Seiten Text und sieben Original-Abbildungen. 1895. ℳ 4.—.

Einleitung der Fäkalien Münchens in die Isar. Protokoll der Sitzung d. erweiterten Obermedicinal-Ausschusses. 73 S. 1892. gr. 8°. ℳ 1.20.

Prausnitz, Dr. W., Zur Einführung der Schwemmkanalisation in München. Offener Brief an Prof. Alex. Müller in Berlin. 1891. gr. 8°. 18 S. ℳ —.60.

Ripperger, A., Die Influenza. Ihre Geschichte. Epidemiologie. Aetiologie, Symptomatologie und Therapie, sowie ihre Komplikationen u. Nachkrankheiten. 338 S. M. 4 Taf. 1892. Brosch. ℳ 10.—.

Soxhlet, Die chemischen Unterschiede zwischen Kuh- u. Frauenmilch und die Mittel zu ihrer Ausgleichung. 1893. 16 S. 8°. ℳ —.60.

Soxhlet, Prof. D. Ueber Margarine. Bericht an das General-Comité des landwirtschaftl. Vereins Bayern. 197 S. 1895. ℳ 2.40.

Annalen der Städtischen Allgemeinen Krankenhäuser zu München.

Im Verein mit den Aerzten dieser Anstalten herausgegeben von **Prof. Dr. von Ziemssen,** Director des Städt. Allgem. Krankenhauses I/I.

Band VI. 1890—92. 477 Seiten mit 19 Abbildungen. Preis M. **10.**—.
Band VII. 1893. 366 Seiten mit 4 Abbildungen. Preis M. **10.**—.
„ VIII. 1894. 369 „ „ 16 „ „ M. **10.**—.
„ IX. 1895. 455 „ „ 3 „ „ M. **12.**—.
„ X. 1896. 382 „ „ 12 „ „ M. **10.**—.

Diese Bände erscheinen in wesentlich veränderter Form. Während früher die statistischen Tabellen den Hauptinhalt bildeten, enthalten diese Bände vorzugsweise allgemein interessierende kasuistische Beiträge und lehnen sich in ihrer Einrichtung ganz *an die* Charité-Annalen an.

Verlag von J. F. LEHMANN in MÜNCHEN.

Lehmann's medicinische Handatlanten,

nebst kurzgefassten Lehrbüchern.

Herausgegeben von

Prof. Dr. **O. Bollinger**, Dr. **G. Brühl**, Doz. Dr. **H. Dürck**, Dr. E. **Golebiewski**, Dr. **L. Grünwald**, Prof. Dr. **O. Haab**, Prof. Dr. **H. Helferich**, Prof. Dr. **A. Hoffa**, † Prof. Dr. **E. von Hofmann**, Prof. Dr. **Chr. Jakob**, Prof. Dr. **K. B. Lehmann**, Doz. Dr. **Lüning**, Doz. Dr. **G. Marwedel**, Prof. Dr. **Mracek**, Dr. **R. Neumann**, Dozent Dr. **O. Schäffer**, Doz. Dr. **Schulthess**, Prof. Dr. **Schultze**, Doz. Dr. **J. Sobotta**, Doz. Dr. **W. Weygandt**, Doz. Dr. **O. Zuckerkandl**, u. a. m.

Bücher von hohem wissenschaftlichen Werte, in bester Ausstattung, zu billigem Preise.

Urteile der Presse:

Therapeutische Monatshefte.

Es ist entschieden als ein glücklicher Gedanke des Verlegers zu bezeichnen, das, was in der Medizin bildlich darzustellen ist, in Form von Handatlanten zu bringen, die infolge ihres ausserordentlich niedrigen Preises jedermann leicht zugänglich sind.

Medico.

Es ist als verdienstvolles Unternehmen der Lehmann'schen Verlagsbuchhandlung zu bezeichnen, dass sie in einer Serie von gut ausgeführten und doch billigen Handatlanten einen Ersatz für die, dem grossen Kreise der Interessenten, wegen der meist sehr erheblichen Anschaffungskosten kaum zugänglichen grösseren Werke, geschaffen hat. Denn bildliche Darstellungen sind für das Verständnis ein kaum zu entbehrendes Hilfsmittel.

Wiener medicinische Wochenschrift.

Sowohl der praktische Arzt als der Student empfinden gewiss vielfach das Bedürfnis, die Schilderung des Krankheitsbildes durch gute, bildliche Darstellung ergänzt zu sehen. Diesem allgemeinen Bedürfnisse entsprechen die bisherigen Atlanten und Bildwerke wegen ihrer sehr erheblichen Anschaffungskosten nicht. Das Unternehmen des Verlegers verdient daher alle Anerkennung. Ist es doch selbst bei eifrigem Studium kaum möglich, aus der wörtlichen Beschreibung der Krankheitsbilder sich allein eine klare Vorstellung von den krankhaften Veränderungen zu machen. Der Verleger ist somit zu der gewiss guten Idee zu beglückwünschen, ebenso glücklich war die Wahl der Fachmänner, unter deren Aegide die bisherigen Atlanten erschienen sind.

Verlag von J. F. LEHMANN in MÜNCHEN.

Lehmann's medicin. Hand-Atlanten

II. Band:

Atlas und Grundriss

der Lehre vom Geburtsakt

und der operativen

Geburtshilfe

dargestellt in 249 Tafeln in Lepurellinart

nebst kurzgefasstem Lehrbuche

von **Dr. O. Schäffer,**

Privatdocent an der Universität Heidelberg.

124 zweifarbige Bilder auf 56 grossfolige Tafeln.

V. gänzlich umgearbeitete Auflage.

Preis elegant gebunden Mk. 8.—.

Die Wiener medicinische Wochenschrift schreibt:

— — Die diesem Atlas beigegebenen so guten Bilder geben in Verein mit dem knappen aber sehr verständlichen Text Alles, was man in der Praxis wissen muss.

Die beste Ergänzung dieses Atlas ist:

Dr. O. Schaeffer's

Geburtshilfliches Phantom

Becken aus Metallguss nebst Leder-Puppe mit Spiral-Drahtgestell und Kopf aus weichem Gummi.

⅓ natürl. Grösse.

Preis in Kistchen Mk. 20.—.

Zum ersten Male wird hier ein Phantom geliefert, das in Folge seines billigen Preises und seiner vorzüglichen Verwendbarkeit allen Anforderungen entspricht und dem Studenten wie dem Praktiker das Studium ausserordentlich erleichtert. Vermittelst der elastischen Lederpuppe, die durch Gummibänder in jeder Stellung festgehalten werden kann, lassen sich alle vorkommenden Lagen darstellen. Auch alle verschiedenen Schädelformen sind, da der Kopf aus weichem Gummi besteht, sehr gut bei der Durchführung durch das Becken hervorzubringen. Das Becken wird an die Tischplatte angeschraubt, sodass sich bequem arbeiten lässt. Als Text und Vorlagenwerk gilt O. Schäffer, Atlas und Grundriss der Lehre vom Geburtsakt. 5. Auflage. (Verlag von J. F. Lehmann. Preis Mk. 8.—.)

Verlag von J. F. LEHMANN in MÜNCHEN.

Lehmann's medicin. Handatlanten.

Band II:

Anatomischer Atlas

der

geburtshilflichen Diagnostik und Therapie.

Mit 160 meist farbigen Abbildungen auf Tafeln nach Originalien von den Malern A. Schmitson und C. Krapf, zahlreichen Text-Illustrationen und 318 Seiten Text

von

Dr. Oskar Schaeffer,

Privatdozent an der Universität Heidelberg.

Zweite vollständig umgearbeitete und erweiterte Auflage.

Preis eleg. geb. Mk. 12.—

Prof. Fritsch, Bonn, schreibt: (Centralbl. f. Gynaekologie 1895. No. 39)

Als Gegengewicht gegen die quantitative Vermehrung des Lernstoffes hat man vielfach die Lehrmittel verbessert. Es sind kurze Compendien, instruktive Abbildungen eingeführt, ungefähr so, wie im Elementarunterricht das Lernen durch den sogen. Anschauungsunterricht erleichtert wird.

Diese Tendenz verfolgen auch die bei Lehmann erschienenen Atlanten. Einer der besten ist jedenfalls der von S. Ich möchte den Studenten mehr diesen Atlas als eines der modernen Compendien empfehlen. Alle Zeichnungen sind einfach, übersichtlich und jedenfalls so hergestellt, dass der Lernende auf den ersten Blick das sieht, was er sehen soll.

Es wäre sehr zu wünschen, dass diese Atlanten von den Lehrern überall warm empfohlen würden.

Münchener medicinische Wochenschrift 1894 Nr. 10

Ein Atlas von ganz hervorragender Schönheit der Bilder zu einem überraschend niedrigen Preise. Auswahl und Ausführung der meisten Abbildungen ist gleich anerkennenswert, einzelne derselben sind geradezu mustergiltig schön. Verfasser, Zeichner und Verleger haben sich um diesen Atlas in gleicher Weise verdient gemacht.

Der Text bietet mehr, als der Titel verspricht: er enthält — abgesehen von den geburtshilflichen Operationen — ein vollständiges Compendium der Geburtshilfe. Damit ist dem Praktiker und dem Studierenden Rechnung getragen, welche in dem Buche neben einem Bilderatlas auch das finden, was einer Wiedergabe durch Zeichnungen nicht bedarf.

Das Werkchen wird wohl mehrere Auflagen erleben. Als Atlas betrachtet, dürfte das Buch an Schönheit und Brauchbarkeit alles übertreffen, was an Taschen-Atlanten überhaupt und zu so niedrigem Preise im besonderen geschaffen wurde.

Verlag von J. F. LEHMANN in MÜNCHEN.

Band III:

Handatlas u. Grundriss der Gynäkologie.

Mit 90 farbigen Tafeln, 65 Textillustrationen und 308 Seiten Text.

Von **Dr. O. Schäffer,** Privatdozent an der Universität Heidelberg.

II. vollständig umgearbeitete und erweiterte Auflage.

Preis elegant geb. *M.* **14.—.**

Urteile der Presse:

Medicinisch-chirurg. Central-Blatt. Der vorliegende Band der von uns schon wiederholt rühmlich besprochenen Lehmann'schen medicinischen Atlanten bringt eine Darstellung des gesamten Gebietes der Gynaekologie. Die trefflich ausgeführten Abbildungen bringen Darstellungen von klinischen Fällen und anatomischen Präparaten, wobei besonders hervorzuheben ist, dass jeder einzelne Gegenstand von möglichst vielen Seiten, also aetiologisch, in der Entwickelung, im secundären Einfluss, im Weiterschreiten und im Endstadium oder der Heilung dargestellt ist, und dass die Abbildungen von Präparaten wieder durch schematische und halbschematische Zeichnungen erläutert sind. Der Text zerfällt in einen fortlaufenden Teil, der von rein praktischen Gesichtspunkten bearbeitet ist und in die Erklärung der Tafeln, welche die theoretischen Ergänzungen enthält. Ausführliche Darlegungen über den Gebrauch der Sonde, der Pessarien werden vielen Praktikern willkommen sein. Eingehende Berücksichtigung der Differentialdiagnose, sowie Zusammenstellung der in der Gynaekologie gebräuchlichen Arzneimittel, sowie deren Anwendungsweisen erhöhen die praktische Brauchbarkeit des Buches.

Therapeutische Monatshefte: Der vorliegende Band reiht sich den Atlanten der Geburtshilfe desselben Autors ebenbürtig an. Er entspricht sowohl den Bedürfnissen des Studierenden wie denen des Praktikers. Der Schwerpunkt des Werkes liegt in den Abbildungen. In den meisten Fällen sind diese direkt nach der Natur oder nach anatomischen Präparaten angefertigt. Manche Zeichnungen sind der bessern Uebersicht wegen mehr schematisch gehalten. Auch die einschlägigen Kapitel aus der Hystologie (Tumoren, Endometritisformen etc.) sind durch gute Abbildungen vertreten. Besonders gelungen erscheinen uns die verschiedenen Spiegelbilder der Portio. Jeder Tafel ist ein kurzer begleitender Text beigegeben. Der 2. Teil des Werkes enthält in gedrängter Kürze die praktisch wichtigen Grundzüge der Gynaekologie; übersichtlich sind bei jedem einzelnen Krankheitsbilde die Symptome, die differentiell-diagnostisch wichtigen Punkte u. s. w. zusammengestellt. Feis (Frankfurt a. M.).

Verlag von J. F. LEHMANN in MÜNCHEN.

Lehmann's medic. Hand-Atlanten.

Band IV:

Atlas der Krankheiten der Mundhöhle, des Rachens und der Nase.

In 69 meist farbigen Bildern auf Tafeln und erklärendem Text von

Dr. L. Grünwald.

Preis eleg. geb. ℳ 6.—.

Der Atlas beabsichtigt, eine Schule der semiotischen Diagnostik zu geben. Daher sind die Bilder derart bearbeitet, dass die einfache Schilderung der aus denselben ersichtlichen Befunde dem Beschauer die Möglichkeit einer Diagnose bieten soll. Dem entsprechend ist auch der Text nichts weiter, als die Verzeichnung dieser Befunde, ergänzt, wo notwendig, durch anamnestische u. s. w. Daten. Wenn demnach die Bilder dem Praktiker bei der Diagnosenstellung behilflich sein können, lehrt anderseits der Text den Anfänger, wie er einen Befund zu erheben und zu deuten hat.

Von den Krankheiten der Mund- und Rachenhöhle sind die praktisch wichtigen sämtlich dargestellt, wobei noch eine Anzahl seltenerer Krankheiten nicht vergessen sind. Die Bilder stellen möglichst Typen der betreffenden Krankheiten im Anschluss an einzelne beobachtete Fälle dar.

Band V.

Atlas und Grundriss

der

Hautkrankheiten

mit 65 farbigen Tafeln nach Originalaquarellen des Malers Arthur Schmitson und zahlreichen schwarzen Abbildungen

von **Prof. Dr. Franz Mracek in Wien.**

Dieser Band, die Frucht jahrelanger wissenschaftlicher und künstlerischer Arbeit enthält neben 65 farbigen Tafeln von ganz hervorragender Schönheit noch zahlreiche schwarze Abbildungen, und einen reichen, das gesamte Gebiet der Dermatologie umfassenden Text. Die Abbildungen sind durchwegs Originalaufnahmen nach dem lebenden Materiale der Mracek'schen Klinik, und die Ausführung der Tafeln übertrifft die Abbildungen aller, selbst der teuersten bisher erschienenen dermatologischen Atlanten.

Der Preis des Buches beträgt eleg. geb. Mk. 14.—

Verlag von J. F. LEHMANN in MÜNCHEN.

Lehmann's medic. Handatlanten.

Band VI:

Atlas der Syphilis

und der

venerischen Krankheiten

mit einem

Grundriss der Pathologie und Therapie derselben

mit 71 farbigen Tafeln nach Originalaquarellen

von Maler A. SCHMITSON und 16 schwarzen Abbildungen

von

Professor Dr. **Franz Mracek** in Wien.

Preis des starken Bandes eleg. geb. Mk. 14.—

Nach dem einstimmigen Urteile der zahlreichen Autoritäten, denen die Originale zu diesem Werke vorlagen, übertrifft dasselbe an Schönheit Alles, was auf diesem Gebiete nicht nur in Deutschland, sondern in der gesamten Weltliteratur geschaffen wurde.

Die **Ungarische medicinische Presse** Nr. 41 vom 19. XI. 1897 schreibt:

„Es wird wohl genügen den Titel dieses Werkes niederzuschreiben den Autor und Verleger zu nennen, um in den weitesten Kreisen lebhaftes Interesse für dasselbe zu erregen. Bei der Besprechung des Werkes hört eigentlich jede Kritik auf und die beschreibende Schilderung tritt in ihr Recht. Mit dieser Bemerkung wollen wir aber unsere Schwäche eingestehen und die Unmöglichkeit anerkennen, die durchaus lehrreichen, frappant schönen und naturgetreuen Abbildungen durch Beschreibung vor den Augen der Leser auch nur annähernd begreiflich zu machen. Alles, was die bunten und zahlreichen syphilitischen Erkrankungsformen Lehrreiches nur bieten können, ist in diesem schönen Werke klassisch dargestellt, in einem leicht fassbaren System gruppiert. Die meisterhafte Hand des Malers spricht klar und decidiert zu dem Studierenden, so dass man durch diese Tafeln thatsächlich all das spielend erlernen kann, was man sich sonst auf diesem Gebiete nur durch viel Mühe, Zeit und Erfahrung anzueignen im stande wäre. Um alles zu sehen, was man sehen *muss*, dient noch der erläuternde Text über Syphilis, aus welchem *nicht nur der* heutige Stand der Lehre, sondern gleichzeitig auch *eine rationelle* Therapie herauszulesen ist." N.

Verlag von J. F. LEHMANN in MÜNCHEN.

Lehmann's medic. Handatlanten.

Band VII:

Atlas und Grundriss

der

Ophthalmoscopie u. ophthalmoscop. Diagnostik.

Mit 6 Text- und 138 farbigen Abbildungen auf 88 Tafeln.

Von Professor **Dr. O. Haab**, Direktor der Augenklinik in Zürich.

III. vielfach erweiterte Auflage.

Preis eleg. geb. **M. 10.—.**

Urteile der Presse:

Schmidt's Jahrbücher 1895, S. 211: Endlich wieder einmal ein Buch, das für den praktischen Arzt von wirklichem, dauerndem Nutzen, *für den im Ophthalmoscopieren auch nur einigermassen Geübten geradezu ein Bedürfnis ist. Das Buch enthält im I. Teil eine kurze vortreffliche Anleitung zur Untersuchung mit dem Augenspiegel. Was der Mediciner wissen muss und was er sich auch merken kann, das ist alles in diesen praktischen Regeln zusammengestellt. Der II. Teil enthält auf 64 Tafeln die Abbildungen des Augenhintergrundes in normalem Zustande und bei den verschiedenen Krankheiten. Es sind nicht seltene Fälle berücksichtigt, sondern die Formen von Augenerkrankungen, die am häufigsten und unter wechselndem Bilde vorkommen. Der grossen Erfahrung Haab's und seiner bekannten grossen Geschicklichkeit im Zeichnen ist es zu danken, dass ein mit besonderen Schwierigkeiten verbundener Atlas in dem vorliegenden Werke in geradezu vorzüglicher Weise zu stande kam.*

(Lamhofer, Leipzig.)

Correspondenzblatt f. schweiz. Aerzte: *Ein prächtiges Werk. Die mit grosser Naturtreue wiedergegebenen Bilder des kranken und gesunden Augenhintergrundes bilden eine vorzügliche Studie für den ophthalmologischen Unterricht sowohl als für die ophthalmologische Diagnose in der Praxis.*

Eine vorzügliche Ergänzung zu diesem Atlas bildet das:

Skizzenbuch

zur Einzeichnung von Augenspiegel-Bildern.

Von Professor **Dr. O. Haab,**

Professor an der Universität und Direktor der Augenklinik in Zürich.

Preis in Mappe M. 3.—.

II. Auflage.

Jeder Käufer des Haab'schen Atlas wird auch gern das Skizzenbuch erwerben, da er in diesem mit geringer Mühe alle Fälle, die er in seiner Praxis zu untersuchen hat, naturgetreu darstellen kann.

Verlag von J. F. LEHMANN in München.

Lehmann's medicin. Handatlanten.

Band IX.

ATLAS

des gesunden u. kranken Nervensystems

nebst Grundriss der Anatomie, Pathologie und Therapie desselben

von

Dr. Christfried Jakob,

Vorstand d. patholog. Institutes f. Gehirn- u. Geisteskrankheiten a. d. Universität Buenos-Ayres,
s. Z. I. Assistent der medizin. Klinik in Erlangen.

Mit einer Vorrede von *Prof. Dr. Ad. v. Strümpell*, Direktor der medicin. Klinik in Erlangen.

II. vollständig umgearbeitete Auflage.

Mit 105 farbigen und 120 schwarzen Abbildungen sowie 284 Seiten Text und zahlreichen Textillustrationen.

Preis eleg. geb. **Mk. 14.—**

Prof. Dr. Ad. von Strümpell schreibt in seiner Vorrede zu dem vorliegenden Bande: Jeder unbefangene Beurteiler wird, wie ich glaube, gleich mir den Eindruck gewinnen, dass die Abbildungen Alles leisten, was man von ihnen erwarten darf. Sie geben die thatsächlichen Verhältnisse in deutlicher und anschaulicher Weise wieder und berücksichtigen in grosser Vollkommenheit fast alle die zahlreichen und wichtigen Ergebnisse, zu denen das Studium des Nervensystems in den letzten Jahrzehnten geführt hat. Dem Studierenden sowie dem um diesem Zweige der medicinischen Wissenschaft noch nicht näher vertrauten praktischen Arzt, ist somit die Gelegenheit geboten, sich mit Hilfe des vorliegenden Atlasses *verhältnismässig* leicht ein klares Bild von dem jetzigen Standpunkte der gesamten *Neurologie* zu machen.

Verlag von J. F. LEHMANN in MÜNCHEN.

Lehmann's medicin. Handatlanten.

Band X.

Atlas und Grundriss der Bakteriologie

und

Lehrbuch der speciellen bakteriolog. Diagnostik.

Von Prof. **Dr. K. B. Lehmann** und **Dr. R. Neumann** in Würzburg.

Bd. I Atlas mit ca. 700 farb. Abbildungen auf 69 Tafeln, Bd. II Text 496 Seiten mit 30 Bildern.

II. vielfach erweiterte Auflage.

Preis der 2 Bände eleg. geb. Mk. 16.—

Münch. medic. Wochenschrift 1896 Nr. 23. Sämtliche Tafeln sind mit ausserordentlicher Sorgfalt und so naturgetreu ausgeführt, dass sie ein glänzendes Zeugnis von der feinen Beobachtungsgabe sowohl, als auch von der künstlerisch geschulten Hand des Autors ablegen.

Bei der Vorzüglichkeit der Ausführung und der Reichhaltigkeit der abgebildeten Arten ist der Atlas ein wertvolles Hilfsmittel für die Diagnostik, namentlich für das Arbeiten im bakteriologischen Laboratorium, indem es auch dem Anfänger leicht gelingen wird, nach demselben die verschiedenen Arten zu bestimmen. Von besonderem Interesse sind in dem 1. Teil die Kapitel über die Systematik und die Abgrenzung der Arten der Spaltpilze. Die vom Verfasser hier entwickelten Anschauungen über die Variabilität und den Artbegriff der Spaltpilze mögen freilich bei solchen, welche an ein starres, schablonenhaftes System sich weniger auf Grund eigener objektiver Forschung, als vielmehr durch eine auf der Zeitströmung und unerschütterlichem Autoritätsglauben begründete Voreingenommenheit gewöhnt haben, schweres Bedenken erregen. Allein die Lehmann'schen Anschauungen entsprechen vollkommen der Wirklichkeit und es werden dieselben gewiss die Anerkennung aller vorurteilslosen Forscher finden. — —

So bildet der Lehmann'sche Atlas nicht allein ein vorzügliches Hilfsmittel für die bakteriologische Diagnostik, sondern zugleich einen bedeutsamen Fortschritt in der Systematik und in der Erkenntnis des Artbegriffes bei den Bakterien. Prof. Dr. Hauser.

Allg. Wiener medicin. Zeitung 1896 Nr. 28. Der Atlas kann als ein sehr sicherer Wegweiser bei dem Studium der Bakteriologie bezeichnet werden. Aus der Darstellungsweise Lehmann's leuchtet überall gewissenhafte Forschung, leitender Blick und volle Klarheit hervor.

Pharmazeut. Zeitung 1896 S. 471/72. Fast durchweg in Originalfiguren zeigt uns der Atlas die prachtvoll gelungenen Bilder aller für den Menschen pathogenen, der meisten tierpathogenen und sehr vieler indifferenter Spaltpilze in verschiedenen Entwickelungsstufen.

Trotz der Vorzüglichkeit des „Atlas" ist der „Textband" die eigentliche wissenschaftliche That.

Für die Bakteriologie hat das neue Werk eine neue, im Ganzen auf botanischen Prinzipien beruhende Nomenklatur geschaffen und diese muss und wird angenommen werden. C. Mez - Breslau.

Verlag von J. F. LEHMANN in MÜNCHEN.

Lehmann's medic. Hand-Atlanten.

Band XI/XII:

Atlas u. Grundriss der patholog. Anatomie.

In 120 farbigen Tafeln nach Originalen von Maler **A. Schmitson.**

Preis jeden Bandes eleg. geb. Mk. 12.—

Von Obermedicinalrat Professor **Dr. O. Bollinger.**

Prof. Bollinger hat es unternommen, auf 120 durchwegs nach Original-Präparaten des pathologischen Institutes in München aufgenommenen Abbildungen einen Atlas der pathologischen Anatomie zu schaffen und diesem durch Beigabe eines concisen, aber umfassenden Grundrisses dieser Wissenschaft, auch die Vorzüge eines Lehrbuches zu verbinden.

Von dem glücklichen Grundsatze ausgehend, unter Weglassung aller Raritäten, nur das dem Studierenden wie dem Arzte wirklich Wichtige, das aber auch in erschöpfender Form zu behandeln, wurde hier ein Buch geschaffen, das wohl mit Recht zu den praktischsten und schönsten Werken unter den modernen Lehrmitteln der medizinischen Disziplinen zählt. Es ist ein Buch, das aus der Sektionspraxis hervorgegangen und daher wie kein anderes geeignet ist, dem secierenden Arzte und Studenten Stütze resp. Lehrer bei der diagnostischen Sektion zu sein.

Die farbigen Abbildungen auf den 120 Tafeln sind in 15 fachem Farbendruck nach Originalaquarellen des Malers A. Schmitson hergestellt und können in Bezug auf Naturwahrheit und Schönheit sich dem besten auf diesem Gebiete Geleisteten ebenbürtig an die Seite stellen. Auch die zahlreichen Textillustrationen sind von hervorragender Schönheit. Der Preis ist im Verhältnis zum Gebotenen sehr gering.

Excerpta medica (1896. 12): Das Werk birgt lauter Tafeln, die unsere Bewunderung erregen müssen. Die Farben sind so naturgetreu wiedergegeben, dass man fast vergisst, nur Bilder vor sich zu haben. Auch der Text dieses Buches steht, wie es sich bei dem Autor von selbst versteht, auf der Höhe der Wissenschaft, und ist höchst präcis und klar gehalten.

Korrespondenzblatt f. Schweizer Aerzte 1895 24: Die farbigen Tafeln des vorliegenden Werkes sind geradezu mustergiltig ausgeführt. Die komplizierte Technik, welche dabei zur Verwendung kam (15 facher Farbendruck nach Original-Aquarellen) lieferte überraschend schöne, naturgetreue Bilder, nicht nur in der Form, sondern eben namentlich in der Farbe, so dass man hier wirklich von einem Ersatz des natürlichen Präparates reden kann. Der praktische Arzt, welcher erfolgreich seinen Beruf ausüben soll, darf die pathol. Anatomie, „diese Grundlage des ärztl. Wissens und Handelns" (Rokitansky) zeitlebens nie verlieren. — Der vorliegende Atlas wird ihm dabei ein ausgezeichnetes Hilfsmittel sein, dem sich zur Zeit, namentlich wenn man den geringen Preis berücksichtigt, nichts Aehnliches an die Seite stellen lässt. Die Mehrzahl der Tafeln sind reine Kunstwerke; der verbindende Text aus der bewährten Feder Prof. Bollinger's gibt einen zusammenhängenden Abriss der für den Arzt wichtigsten path.-anat. Prozesse. — Verfasser und Verleger ist zu diesem prächtigen Werke zu gratulieren.

E. Haffter
(Redacteur d. Korr.-Bl. f. Schweizer Aerzte).

Verlag von J. F. LEHMANN in MÜNCHEN.

Lehmann's medicin. Handatlanten.

Band XIV:

Atlas

und

Grundriss

der

Kehlkopfkrankheiten.

Mit 44 farbigen Tafeln und zahlreichen Textillustrationen nach Orginalaquarellen des Malers Bruno Keilitz

von **Dr. Ludwig Grünwald in München.**

Preis elegant geb. Mk. 8.—.

Dem oft und gerade im Kreise der praktischen Aerzte und Studierenden geäusserten Bedürfnisse nach einem farbig illustrierten Lehrbuch der Kehlkopfkrankheiten, das in knapper Form das anschauliche Bild mit der im Text gegebenen Erläuterung verbindet, entspricht das vorliegende Werk des bekannten Münchener Laryngologen. Weit über hundert praktisch wertvolle Krankheitsfälle und 30 mikroskopische Präparate, nach Naturaufnahmen des Malers Bruno Keilitz, sind auf den 44 Volltafeln in hervorragender Weise wiedergegeben, und der Text, welcher sich in Form semiotischer Diagnose an diese Bilder anschliesst, gehört zu dem Instruktivsten, was je über dieses Gebiet geschrieben wurde.

Verlag von J. F. LEHMANN in MÜNCHEN

Lehmann's medicin. Hand-Atlanten.

Band XV.

ATLAS

der klinischen

Untersuchungsmethoden

nebst

Grundriss der klinischen Diagnostik

und der

speziellen Pathologie und Therapie der inneren Krankheiten

von

Prof. Dr. Christfr. Jakob,

s. Z. I. Assistent der medizinischen Klinik in Erlangen.

Mit 182 farbigen Abbildungen auf 68 Tafeln und 250 Seiten Text mit 64 Textabbildungen.

Preis eleg. geb. 10 Mark.

Während alle anderen Atlanten sich meist nur an Specialisten wandten, bietet dieser Band für **jeden praktischen Arzt und für jeden Studenten ein geradezu unentbehrliches Vademecum.**

Neben einem vorzüglichen Atlas der klinischen Mikroskopie sind in dem Bande die **Untersuchungsbefunde aller inneren Krankheiten** in instruktivster Weise in 50 vielfarbigen schematischen Bildern zur Darstellung gebracht. Nach dem Urteil eines der hervorragendsten Kliniker, ist das Werk für den Studierenden ein Lehrmittel von unschätzbarem Werte, für den praktischen Arzt ein Repetitorium, in dem er sich sofort orientieren kann und das ihm in der täglichen Praxis vorzügliche Dienste leistet.

Wandtafeln für den neurologischen Unterricht.

Herausgegeben von

Prof. Dr. Ad. v. Strümpell in Erlangen und **Prof. Dr. Chr. Jakob** in Buenos Ayres.

13 Tafeln im Format von 80 cm zu 100 cm.

Preis in Mappe **Mk. 50.—.** Auf Leinwand aufgezogen Mk. 70.—

Der Text in den Bildern ist lateinisch.

Verlag von J. F. LEHMANN in MÜNCHEN.

Lehmann's medicinische Handatlanten.

Band XVI.

Atlas
und
Grundriss
der

chirurgischen Operationslehre

von

Dr. Otto Zuckerkandl

Privatdozent an der Universität Wien.

24 farbige Tafeln nach Originalaquarellen des Malers Bruno Keilitz.

217 schwarze Abbildungen meist auf Tafeln. 27 Bog. Text.

Preis eleg. geb. M. 10.—

Geheimrat Prof. Dr. **Helferich** schreibt in der M. M. W. Nr. 45 vom 9. XI. 1897:

Der vorliegende stattliche Band enthält 24 farbige Tafeln und 217 Abbildungen im Text bei 400 Seiten Text. Auf 41 Seiten mit einer Tafel (Darmnaht) und 38 Textbildern wird zunächst im allgemeinen die Durchtrennung und die Wiedervereinigung der Gewebe zur Darstellung gebracht. Auf Seite 42 bis 400 wird die spezielle Operationslehre (Extremitäten, Kopf und Hals, Rumpf und Becken) dargestellt.

Das Werk wird den Anfängern das Studium und Verständnis der z. T. recht schwierigen Kapitel in hohem Grade erleichtern. Es existirt unseres Wissens kaum ein Werk, welches mit so zahlreichen und vorzüglichen Abbildungen ausgestattet und mit so bündig klarem Text, den neuesten Fortschritten Rechnung tragend, im ganzen zur Einführung wie zur Fortbildung in der operativen Chirurgie so gut geeignet wäre, wie das vorliegend. Der Erfahrenere wird sich freuen, manches speziell nach der Technik und Anschauung der Wiener Schule dargestellt zu finden. Die Abbildungen sind zum grössten Teil vorzüglich und fast auf den ersten Blick klar und verständlich. Nur einige wenige sind weniger klar und eventuell bei einer zweiten Auflage zu erneuern. Dass eine solche nicht lange *auf sich warten* lassen wird, darf bei der vielseitigen Brauchbarkeit des Werkes erwartet werden.

Verlag von J. F. LEHMANN in MÜNCHEN.

Lehmann's medic. Handatlanten.

Band XVII.

ATLAS

der

gerichtlichen Medicin

nach Originalen von Maler **A. Schmitson**

mit erläuterndem Text

von

Hofrat Professor Dr. E. Ritter von Hofmann,

Direktor des gerichtlich medizin. Instituts in Wien.

Mit 56 farbigen Tafeln und 193 schwarzen Abbildungen.

Preis eleg. geb. **M. 15.—.**

Zum ersten Male wird hier **Medizinern** und **Juristen** ein Werk geboten, das durch seinen ausserordentlichen Bilder-Reichtum und die mustergiltige Ausführung der farbigen Tafeln eine vorzügliche Einführung in die gerichtliche Medizin bietet. Dabei ist der **Preis von 15 Mark** im Vergleich zu dem Gebotenen ganz erstaunlich billig, wenn man bedenkt, dass der Atlas von Lesser, der weniger und kaum so gut ausgeführte Tafeln enthält, M. 180.— kostet. Der Hofmann'sche Atlas bildet eine Ergänzung zu sämtlichen Lehrbüchern der gerichtlichen Medizin. Für **Gerichtsärzte, Physici, Staatsanwälte, Untersuchungsrichter** etc. ist das Werk **unentbehrlich,** aber auch jedem Arzte und jedem Juristen bietet es eine Fülle von Anregungen und dem Studierenden erleichtert es das Verständnis und das Eindringen in die schwierige Disziplin ganz ungemein.

Das Amtsblatt des k. Staatsministerium des Innern vom 23. XI. 1897 sagt in seinem Erlasse an sämtliche k. Regierungen, Kammern des Innern und amtliche Aerzte:

Nach dem Gutachten des k. Obermedizinal-Ausschusses bildet der vorliegende Atlas eine Ergänzung des vortrefflichen Lehrbuches des Verfassers, der auf dem Gebiete der gerichtlichen Medizin unbestritten die erste Autorität in deutschen Landen war.

Die technische Wiedergabe der dargestellten Veränderungen ist eine vorzügliche und die von Künstlerhand ausgeführten farbigen Tafeln verdienen namentlich alles Lob.

Es ist hienach Anlass gegeben, auf den bezeichneten Atlas in den einschlägigen Kreisen aufmerksam zu machen und ihn zur Anschaffung zu empfehlen.

München, 13. November 1898. *Frhr. v. Feilitzsch.*

Verlag von J. F. LEHMANN in MÜNCHEN.

Lehmanns' medicin. Handatlanten

Band XVIII.

Atlas und Grundriss

der

äusseren Erkrankungen des Auges.

Von

Prof. Dr. O. Haab,
Direktor d. Augenklinik in Zürich.

Mit 67 farb. Abbildungen auf 40 Taf., zahlreichen Textillustrationen und 16. Bogen Text.

Preis eleg. geb.
M. 10.—.

Dieses neue Werk des rühmlichst bekannten Züricher Ophthalmologen ist wie wenige geeignet, ein wahres Handbuch in der Bücherei eines jeden praktischen Arztes zu werden. Der gleiche, so hervorragend lehrhafte Vortrag, wie er in der Ophthalmoskopie (Atlas, Bd. VII) desselben Autors bewundert wurde, und diesem Werke zu einem ausserordentlichen Erfolge verhalf, ist auch eine glänzende Eigenschaft dieses neuen Buches. Die oft so komplicierten Verhältnisse der Erkrankungen des äusseren Auges sind wohl nie noch klarer und fasslicher veranschaulicht worden, als in dem vorliegenden Werke, in welchem die bildliche mit der verbalen Darstellung sich in schönster Weise ergänzen.

Der Preis des Buches ist mit Hinblick auf das Gebotene ein auffallend niedriger.

Verlag von J. F. LEHMANN in MÜNCHEN.

Lehmann's medicin. Handatlanten

Band XIX.

Atlas und Grundriss der

Unfallheilkunde.

Mit 40 farbigen Tafeln, 142 Textabbildungen
nach Originalen von Maler Johann Fink
und 38 Bogen Text von **Dr. Ed. Golebiewski** in Berlin.
Preis eleg. geb. Mk. **15.—**.

In diesem Werke werden zum ersten Mal systematisch die Nachkrankheiten der Unfallverletzungen beschrieben und durch zahlreiche Abbildungen illustriert.

Die grosse Erfahrung des Verfassers auf dem Gebiete der Unfallheilkunde, das reiche, sehr sorgfältig gesammelte Beobachtungsmaterial, das ihm seit Ende 1886 zur Verfügung steht, berechtigen zu der Annahme, dass in dem vorliegenden Atlas und Grundriss ein Werk entstanden ist, mit welchem einem dringenden Bedürfnis abgeholfen wird.

Das Eigenartige dieses Werkes beruht nicht allein auf den zahlreichen, sehr instruktiven und naturgetreuen Abbildungen — es enthält 40 prachtvoll ausgeführte, nach Original-Aquarellen vom Maler Fink hergestellte Tafeln und 142 schwarze Abbildungen, darunter viele Röntgenbilder — sondern auch in der Behandlung des Textes selbst.

Der Schwerpunkt der textlichen Darstellung liegt in der Beschreibung der functionellen Störungen, wie sie bei den Nachkrankheiten der Unfallverletzungen noch zurückzubleiben pflegen.

Zum genaueren Verständnis dieser funktionellen Störungen ist den einzelnen Abschnitten eine zumeist auf eigenen Studien an der Leiche und am Lebenden basierende Erklärung der normalen anatomisch-funktionellen Verhältnisse vorausgeschickt.

Der Verfasser hat auch bei jedem Kapitel die Anzahl der von ihm selbst beobachteten einschlägigen Fälle angegeben.

Eine sehr reiche und nach vielen Richtungen hin interessante Casuistik ist ferner zur Ergänzung dem Text beigegeben.

Das Werk ist unentbehrlich für jeden Arzt, welcher mit Unfallverletzungen zu thun hat, für Studierende in den höheren Semestern, für Chirurgen, es ist aber auch wegen der zahlreichen Abbildungen und der grossen Casuistik nicht minder *nützlich für nicht ärztliche Organe der Unfallversicherung.*

Verlag von J. F. LEHMANN in MÜNCHEN.

Lehmann's medic. Handatlanten.

Band XX/XXI:

Atlas und Grundriss der pathologischen Histologie.

Spezieller Teil.

120 farbigen Tafeln nach Originalen
des Universitätszeichners **C. Krapf** und reicher Text.

Von Privatdozent **Dr. Hermann Dürck.** Prosektor an d. Krankenhause München l. J.

2 Bände. Preis geb. je M. 11.—.

Dürck hat in diesem Werke dem Lernenden einen zuverlässigen Führer für das weite und schwierige Gebiet der pathologischen Histologie geschaffen. Es wurde dies durch sorgfältige Auswahl der Abbildungen mit Berücksichtigung aller praktisch wichtigeren Gewebsveränderungen und durch Beigabe eines Textes, welcher in gleicher Weise Rücksicht nimmt auf die Bedürfnisse des Studierenden und des Praktikers erreicht. Vor allem sucht der Verfasser den Anfänger darüber aufzuklären, wie das makroskopisch wahrnehmbare Bild einer Organerkrankung durch die Veränderungen an dessen elementaren Bausteinen bedingt wird und warum die krankhaften Prozesse diejenigen Formen hervorbringen mussten, welche uns am Sektionstisch vor Augen treten.

Die sämtlichen Abbildungen sind lithographisch in den Originalfarben der Präparate reproduziert. Bei möglichster Naturtreue wurde der grösste Wert auf Klarheit und korrekte Zeichnung gelegt; in dieser Beziehung sind die Bilder den vielfach auftauchenden photographischen Reproduktionen mikroskopischer Objekte weit überlegen, da sie den Beschauer nicht durch die Massenhaftigkeit der gleichzeitig dargestellten Details und die hiedurch bedingte Unschärfe verwirren, sondern demselben die Kontrole über die Bedeutung jedes einzelnen Striches und Punktes an der Hand der Figurenerklärung und des Textes ermöglichen. Besonders der Anfänger vermag nur aus klaren, eindeutigen Abbildungen klare Vorstellungen zu gewinnen. Der Text schliesst sich den Figuren eng an, beide ergänzen einander und tragen in erster Linie der didaktischen Tendenz des Buches Rechnung. Aus diesem Grunde ist der Besprechung jedes Organs resp. jeder Organgruppe eine kurze, präcise, aber erschöpfende Rekapitulation der normal-histologischen Verhältnisse vorausgeschickt

So stellt das Buch ein wichtiges Hilfsmittel für das Studium der pathologischen Anatomie dar, deren Verständnis undenkbar ist ohne genaue Kenntnis in dem Gebiete der pathologischen Histologie.

Atlas und Grundriss der pathologischen Histologie. Allgemeiner Teil nebst einem Anhange über patholog. histolog Technik von **Privatdozent Dr. Hermann Dürck wird als Band XXII der Atlanten im Sommer 1900** erscheinen.

Verlag von J. F. LEHMANN in MÜNCHEN.

Grundzüge der Hygiene

Pro

Für Studierende

Aer

Vierte

Mit 523 Se

P

Stere

Pathol

In 50 I

Kgl. säch

Preis mit

Dieser

Herzens in au

stellung. Die

das natürliche

Gebu

Mit

VI und 19 S

bewegliche

Geburt

von Dr. Arth

Kleine A

und

M113 Hoffa, A.J. 106119
H69 Atlas und Grundriss
1900 der Verbandlehre.

NAME DATE DUE